Appendix 7. Model Archival Summary for Sodium Concentration at U.S. Geological Survey Site 06887500, Kansas River at Wamego, Kansas, during July 2012 through September 2019

This model archival summary summarizes the sodium (Na; U.S. Geological Survey [USGS] parameter code 00930) concentration model developed to compute 15-minute Na concentrations from July 2012 onward. This model supersedes all previous models.

Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Site and Model Information

Site number: 06887500

Site name: Kansas River at Wamego, Kansas

Location: Lat 39°11'54", long 96°18'19" referenced to North American Datum of 1927, in SW 1/4 NW 1/4 SE 1/4 sec.9, T.10 S.,

R.10 E., Pottawatomie County, Kans., hydrologic unit 10270102.

Equipment: A YSI 6600 water-quality monitor equipped with sensors for water temperature, specific conductance (SC), dissolved oxygen, pH, and turbidity was installed from August 2012 through June 2014. A Xylem YSI EXO2 water-quality monitor equipped with sensors for water temperature, SC, dissolved oxygen, pH, turbidity, and chlorophyll and phycocyanin fluorescence was installed during June 2014 through September 2019. The monitor was housed in a 4-inch-diameter galvanized steel pipe. Readings from the water-quality monitor were recorded every 15 minutes and transmitted by way of satellite, hourly.

Date model was created: March 30, 2020

Model calibration data period: July 19, 2012, through September 23, 2019

Model application date: July 19, 2012, onward

Model-Calibration Dataset

All data were collected using USGS protocols (Wagner and others, 2006; U.S. Geological Survey, variously dated) and are stored in the National Water Information System (U.S. Geological Survey, 2020) database and available to the public. Ordinary least squares analysis was used to develop regression models using R programming language (R Core Team, 2020). Potential explanatory variables that were evaluated individually and in combination included streamflow, water temperature, SC, dissolved oxygen, pH, turbidity, and chlorophyll and phycocyanin fluorescence. The maximum time span between two continuous data points used for interpolation was 2 hours (in order to preserve the sample dataset, field monitor averages obtained during sample collection were used for model development data if no continuous data were available or if gaps larger than 1 hour in the continuous data record resulted in missing interpolated data). Seasonal components (sine and cosine variables) were also evaluated as potential explanatory variables.

The final selected regression model was based on 101 concurrent measurements of Na concentration and sensor-measured SC during July 19, 2012, through September 23, 2019. Samples were collected throughout the range of continuously observed hydrologic conditions. No samples had concentrations below laboratory detection limits. Summary statistics and the complete model-calibration dataset are provided below. Potential outliers were identified using the methods described in Rasmussen and others (2009). Additionally, studentized residuals from the final model were inspected for values greater than three or less than negative three. Values outside of that range were considered potential outliers and were investigated. One of the sensor-measured SC samples, from July 28, 2014, was deemed an outlier and removed from the model calibration dataset. The removed sensor-measured SC value was significantly lower than the field monitor and laboratory result during this sample. All other potential outliers were not found to have errors associated with collection, processing, or analysis and were therefore considered valid.

This model is specific to the Kansas River at Wamego, Kans., during this study period and cannot be applied to data collected from other sites on the Kansas River or data collected from other waterbodies.

Sodium Sampling Details

Cross-section samples typically were collected either from the downstream side of the bridge or instream within 100 feet of the bridge. The equal-width-increment collection method was used (although multiple vertical, single vertical, and grab samples were occasionally collected), and samples typically were composited for analysis (U.S. Geological Survey, variously dated). During July 2012 through June 2017, cross-section samples were collected every 2 weeks during March through October, once a month during November through February, and during selected reservoir release and runoff events. During July 2017 through September 2019, cross-section samples were collected on a monthly to bimonthly basis, depending on flow conditions. A FISP US DH–81, DH–95,

D-95, D-96a, or D-96 depth integrating sampler was used. Additional detail on sample collection is available in Foster and Graham (2016) and Graham and others (2018). Samples were analyzed for Na concentration at the USGS National Water Quality Laboratory in Lakewood, Colorado.

Model Development

Ordinary least squares regression analysis was done using R programming language (R Core Team, 2020) to relate discretely collected Na concentration to sensor-measured SC. The distribution of residuals was examined for normality, and the plots of residuals (the difference between the measured and computed values) were examined for homoscedasticity (departures from zero did not change substantially over the range of computed values). Previously published explanatory variables were also strongly considered for continuity.

SC was selected as a good surrogate for Na based on residual plots, coefficient of determination (R^2), and model standard percentage error. Values for all the aforementioned statistics were computed and are included below along with all relevant sample data and additional statistical information.

Model Summary

The following is a summary of final regression analysis for Na concentration at USGS site 06887500:

Na concentration-based model:

$$\log N\alpha = 1.52 \times \log SC - 2.57$$

where

log = logarithm base 10;

Na =sodium concentration, in milligrams per liter; and

SC = specific conductance, in microsiemens per centimeter at 25 degrees Celsius.

SC makes physical and statistical sense as an explanatory variable for Na because of its positive correlation with charged ionic species (Hem, 1992).

The logarithmically (log) transformed model may be retransformed to the original units so that Na can be calculated directly. The retransformation introduces a bias in the calculated constituent. This bias may be corrected using Duan's bias correction factor (BCF; Duan, 1983). For this model, the calculated BCF is 1.01. The retransformed model, accounting for BCF is as follows:

$$Na = 1.01 \times (SC^{1.52} \times 10^{-2.57})$$

Previous Models

Start Year	End Year	Model Equation	Reference
2012	2019	logNa = 1.52logSC - 2.55	Foster and Graham (2016)
1999	2003	logNa = 1.46logSC - 2.39	Rasmussen and others (2005)

Model Statistics, Data, and Plots


Model


logNa = + 1.52 * logSC - 2.57


Variable Summary Statistics

	logNa	Na	logSC	SC
Minimum	0.991	9.79	2.42	262
1st Quartile	1.560	36.40	2.74	553
Median	1.800	63.80	2.87	749
Mean	1.760	66.70	2.85	762
3rd Quartile	1.990	97.20	2.99	988
Maximum	2.130	134.00	3.11	1280


Box Plots







Exploratory Plots



Red line shows the locally weighted scatterplot smoothing (LOWESS).

The x- and y-axis labels for a given bivariate plot are defined by the intersecting row and column labels.

Basic Model Statistics

Number of Observations	101
Standard error (RMSE)	0.0581
Average Model standard percentage error (MSPE)	13.4
Coefficient of determination (R ²)	0.951
Adjusted Coefficient of Determination (Adj. R ²)	0.95
Bias Correction Factor (BCF)	1.01

Explanatory Variables

	Coefficients	Standard Error	t value	Pr(> t)
(Intercept)	-2.57	0.0989	-25.9	6.02e-46
logSC	1.52	0.0346	43.8	1.31e-66

Correlation Matrix

	Intercept	E.vars
Intercept	1.000	-0.998
E.vars	-0.998	1.000

Outlier Test Criteria


Leverage (Cook's D	DFFITS
0.0594	0.1944	0.2814

Flagged Observations

logNa Estimate Residual Standard Residual Studentized Residual Leverage Cook's D DFFITS 201306170830 1.400 1.23 0.1650 2.920 3.040 0.05270 0.23800 0.718 201308050730 0.991 1.10 -0.1090 -1.940 -1.970 0.07680 0.15700 -0.569 201408110920 1.940 1.78 0.1580 2.740 2.830 0.00999 0.03780 0.285 201409080800 1.300 1.41 -0.1050 -1.840 -1.860 0.02870 0.04990 -0.320 201506290820 1.310 1.19 0.1230 2.180 2.220 0.06020 0.15200 0.562 201810110940 1.160 1.14 0.0284 0.506 0.504 0.06960 0.00959 0.138											
201308050730 0.991 1.10 -0.1090 -1.940 -1.970 0.07680 0.15700 -0.569 201408110920 1.940 1.78 0.1580 2.740 2.830 0.00999 0.03780 0.285 201409080800 1.300 1.41 -0.1050 -1.840 -1.860 0.02870 0.04990 -0.320 201506290820 1.310 1.19 0.1230 2.180 2.220 0.06020 0.15200 0.562		logNa	Estimate	Residual	Standard Res	idual	Studentized	Residual	Leverage	Cook's D	DFFITS
201408110920 1.940 1.78 0.1580 2.740 2.830 0.00999 0.03780 0.285 201409080800 1.300 1.41 -0.1050 -1.840 -1.860 0.02870 0.04990 -0.320 201506290820 1.310 1.19 0.1230 2.180 2.220 0.06020 0.15200 0.562	201306170830	1.400	1.23	0.1650		2.920		3.040	0.05270	0.23800	0.718
201409080800 1.300 1.41 -0.1050 -1.840 -1.860 0.02870 0.04990 -0.320 201506290820 1.310 1.19 0.1230 2.180 2.220 0.06020 0.15200 0.562	201308050730	0.991	1.10	-0.1090	-	1.940		-1.970	0.07680	0.15700	-0.569
201506290820 1.310	201408110920	1.940	1.78	0.1580		2.740		2.830	0.00999	0.03780	0.285
	201409080800	1.300	1.41	-0.1050	-	1.840		-1.860	0.02870	0.04990	-0.320
201810110940 1.160 1.14 0.0284 0.506 0.504 0.06960 0.00959 0.138	201506290820	1.310	1.19	0.1230		2.180		2.220	0.06020	0.15200	0.562
	201810110940	1.160	1.14	0.0284		0.506		0.504	0.06960	0.00959	0.138

201902050950 1.940	2.06 -0.1230	-2.140	-2.180 0.02400 0.05620 -0.341
201904150950 1.990	2.12 -0.1290	-2.260	-2.310 0.03040 0.08030 -0.409

Statistical Plots




First row (left): Residual Na related to regression computed Na with local polynomial regression fitting, or locally estimated scatterplot smoothing (LOESS), indicated by the blue line.


First row (right): Residual Na related to the corresponding normal quantile of the residual with simple linear regression, indicated by the blue line.

Second row: Residual Na related to date (left) and regression computed Na multiplied by the BCF (right) with LOESS, indicated by the blue line.

Third row: Observed Na related to regression computed Na.


Fourth row: Residual Na related to SC with LOESS, indicated by the blue line.





Cross-Validation

Cross-validation



Fold - equal partition of the data (10 percent of the data).

Large symbols - observed value of a data point removed in a fold.

Small symbols - recomputed value of a data point removed in a fold.

Recomputed regression lines - adjusted regression line with one fold removed.


Minimum MSE of folds: 0.00153

Mean MSE of folds: 0.00341

Median MSE of folds: 0.00267

Maximum MSE of folds: 0.00788

(Mean MSE of folds) / (Model MSE): 1.01000



Red line - Model MSE

Blue line - Mean MSE of folds

Model-Calibration Dataset

IVIOC	iei-Calibratio	on Dat	aset								
	Date	logNa	logSC	Na	SC	Computed	Computed	Residual		Censored	
0						logNa	Na		Quantiles	Values	
1	2012-07-19	1.9	2.93	79.9	842	1.87	74.4	0.0349	0.635		
2	2012-07-30	1.8	2.89	63.8	770	1.81	65	-0.004	-0.199		
3	2012-08-13	2	2.99	99.3	970	1.96	92.2	0.0362	0.665		
4	2012-08-27	1.93	2.95	85.5	884	1.9	80.1	0.0323	0.575		
5	2012-09-10	1.93	2.93	84.7	854	1.88	76	0.0509	1.06		
6	2012-09-24	1.94	2.94	88	872	1.89	78.4	0.0538	1.11		
7	2012-10-15	1.9	2.92	79.2	828	1.86	72.5	0.0421	0.795		
8	2012-10-29	2.02	2.99	104	971	1.96	92.3	0.0556	1.15		
9	2012-11-19	1.88	2.92	76.5	837	1.86	73.7	0.0199	0.302		
10	2012-12-17	2.06	3.05	114	1110	2.05	113	0.00739	0.0248		
11	2013-01-14	2.13	3.11	134	1280	2.14	140	-0.0162	-0.328		
12	2013-02-11	2.11	3.07	130	1180	2.09	124	0.0242	0.381		
13	2013-03-11	2.07	3.04	117	1090	2.04	110	0.0306	0.518		
14	2013-04-08	2.11	3.06	129	1150	2.07	119	0.0378	0.696		
15	2013-05-06	1.82	2.84	66.5	684	1.73	54.3	0.0919	1.69		
16	2013-05-20	1.8	2.86	63.3	726	1.77	59.4	0.0313	0.546		
17	2013-06-03	1.44	2.68	27.6	477	1.49	31.4	-0.0523	-0.901		
18	2013-06-17	1.4	2.51	24.9	320	1.23	17.2	0.165	2.52		
19	2013-07-01	1.34	2.61	21.8	404	1.38	24.5	-0.0464	-0.829		
20	2013-07-15	1.79	2.82	61.7	666	1.71	52.1	0.077	1.44		

	21 2013-08-05		2.42 9.7		1.1	12.7	-0.109	-1.6	
	22 2013-08-19	1.4	2.62 25.		1.41	25.7	-0.00193	-0.124	
	23 2013-09-09	1.99	2.98 98.		1.95	89.6	0.0441	0.864	
	24 2013-09-23	2.03		7 1030	2	101	0.0291	0.489	
	25 2013-10-21	2		6 1000	1.98	96.5	0.0174	0.225	
	26 2013-11-18	1.99	2.99 9		1.97	94.8	0.0183	0.25	
	27 2013-12-16	1.68	2.79 48.		1.66	46.4	0.0191	0.276	
	28 2014-01-13	2.03		6 1120	2.06	115	-0.0301	-0.434	
	29 2014-02-10	1.95	3.01 89.		1.99	99	-0.0401	-0.696	
	30 2014-03-10	2.04		0 1160	2.08	121	-0.0371	-0.605	
	31 2014-04-07	2.01	3 10		1.97	95.2	0.0337	0.605	
	32 2014-05-05	1.93	2.97 85.		1.94	87	-0.00444	-0.225	
	33 2014-05-19	2		1 1040	2.01	102	-0.00233	-0.149	
	34 2014-06-02	1.75	2.83 56.		1.72	53.1	0.0278	0.434	
	35 2014-06-11	1.33	2.61 21.		1.38	24.4	-0.0499	-0.864	
	36 2014-06-30	1.53	2.69 33.		1.51	32.6	0.0202	0.328	
	37 2014-07-14	1.8	2.87 63.		1.79	62	0.0149	0.149	
	88 2014-08-11	1.94	2.87 8		1.78	61	0.158	2.15	
	39 2014-08-25	1.88	2.9 75.		1.84	69.2	0.0409	0.761	
	10 2014-09-08	1.3	2.62 20.		1.41	25.8	-0.105	-1.51	
	12014-09-22	1.68	2.75 48.		1.6	39.8	0.0857	1.6	
	2014-10-06	2.05		3 1110	2.05	113	0.00356	-0.0248	
	3 2014-10-20	1.82	2.87 66.		1.78	60.9	0.04	0.728	
	14 2014-11-17	1.93	2.91 84.		1.85	71.1	0.0796	1.51	
	15 2014-12-15	1.74	2.77 54.		1.63	42.9	0.109	1.8	
	16 2015-01-12 17 2015-02-09	2	3 10		1.98	96 116	0.0257	0.407	
	18 2015-02-09 18 2015-03-09	2.07 2.06		8 1130 5 1050	2.06 2.01	116 103	0.0106	0.0992	
	19 2015-04-06	1.97	2.99 93.		1.96	92.5	0.0498 0.0103	1.02 0.0744	
	50 2015-05-04	1.79	2.86 62.		1.77	59.7	0.0226	0.354	
	51 2015-05-18	1.27	2.53 18.		1.77	18.6	0.0220	0.0496	
	52 2015-06-01	1.41	2.65 25.		1.45	28.3	-0.0363	-0.518	
	32 2015-00-01 33 2015-06-15	1.29	2.57 19.		1.32	21.3	-0.0302	-0.462	
	54 2015 06 19 54 2015-06-29	1.31	2.48 20.		1.19	15.5	0.123	1.95	
_	55 2015-07-13	1.34	2.55 2		1.3	20	0.045	0.939	
	66 2015-07-27	1.53	2.73 3		1.57	37.6	-0.0402	-0.728	
	57 2015-08-10	1.76	2.87 57.		1.78	61.3	-0.0219	-0.381	
	8 2015-08-24	1.7	2.85 50.		1.75	57.2	-0.0529	-0.978	
	9 2015-09-08	1.81	2.89 64.		1.81	65.9	-0.00725	-0.276	
	0 2015-09-21	1.84	2.93 69.		1.88	76.8	-0.0371	-0.575	
	51 2015-10-05	1.8	2.91 62.		1.84	69.9	-0.0453	-0.795	
	52 2015-10-19	2.1		6 1120	2.06	115	0.0449	0.901	
6	3 2015-11-16	1.97	2.99 92.		1.97	93.9	-0.0018	-0.0992	
6	54 2015-12-14	1.43	2.71 27.		1.55	35.6	-0.113	-1.95	
6	55 2016-01-11	1.51	2.73 32.	1 537	1.57	37.6	-0.0651	-1.15	
6	66 2016-02-08	1.72	2.88 52.	2 767	1.81	64.6	-0.0886	-1.37	
6	7 2016-03-03	1.84	2.93 68.	9 855	1.88	76.1	-0.0395	-0.665	
6	8 2016-04-04	1.98		1 1010	1.99	98	-0.00466	-0.25	
6	9 2016-05-02	1.56	2.8 36.	4 626	1.67	47.5	-0.112	-1.8	
7	0 2016-05-16	1.45	2.7 28.	4 498	1.52	33.6	-0.0687	-1.25	
7	1 2016-06-06	1.54	2.7 34.	3 496	1.52	33.4	0.0159	0.199	
	2 2016-06-20	1.34	2.6 22.	1 402	1.38	24.3	-0.0367	-0.546	
7	73 2016-07-11	1.75	2.89 56.	5 777	1.81	65.9	-0.0627	-1.11	
7	4 2016-07-25	1.9	2.95 79.	2 887	1.9	80.5	-0.0032	-0.174	
7	75 2016-08-08	1.59	2.76 39.	1 573	1.61	41.5	-0.0226	-0.407	

76	2016-08-22	1.79	2.9 61.4	787	1.82	67.1	-0.035	-0.489	
77	2016-09-12	1.5	2.72 31.4	523	1.55	36.1	-0.0573	-1.06	
78	2016-09-26	1.83	2.87 67.6	742	1.78	61.4	0.0455	0.978	
79	2016-10-11	1.81	2.85 65.2	713	1.76	57.8	0.056	1.25	
80	2016-10-24	2.05	3.05 112	1110	2.05	113	-0.000302	-0.0744	
81	2016-11-07	1.99	3.04 98	1100	2.04	112	-0.0523	-0.939	
82	2016-12-12	1.69	2.8 49.1	629	1.68	47.8	0.0154	0.174	
83	2017-01-09	1.86	2.97 72.7	925	1.93	85.8	-0.0682	-1.2	
84	2017-02-06	1.99	3.03 97.2	1080	2.03	108	-0.0438	-0.761	
85	2017-03-06	2.03	3.04 108	1100	2.04	112	-0.0101	-0.302	
86	2017-04-10	1.52	2.77 32.9	585	1.63	42.8	-0.111	-1.69	
87	2017-05-08	1.62	2.76 41.5	573	1.61	41.5	0.0037	0	
88	2017-05-22	1.71	2.79 51	619	1.66	46.6	0.0428	0.829	
89	2017-06-05	1.76	2.84 57	696	1.74	55.8	0.0132	0.124	
90	2017-06-19	1.8	2.84 62.7	684	1.73	54.3	0.0664	1.31	
91	2018-10-11	1.16	2.44 14.6	277	1.14	13.8	0.0284	0.462	
92	2018-11-28	1.45	2.61 28.1	409	1.39	24.9	0.0557	1.2	
93	2018-12-17	1.66	2.74 45.8	553	1.59	39.3	0.0698	1.37	
94	2019-02-05	1.94	3.05 86.4	1130	2.06	116	-0.123	-2.15	
95	2019-03-18	1.74	2.87 54.4	749	1.79	62.3	-0.0547	-1.02	
96	2019-04-15	1.99	3.09 98.4	1240	2.12	134	-0.129	-2.52	
97	2019-05-09	1.21	2.54 16.3	346	1.28	19.3	-0.0706	-1.31	
98	2019-06-25	1.39	2.63 24.5	431	1.43	27	-0.0378	-0.635	
99	2019-07-15	1.58	2.73 38	542	1.58	38.2	0.00167	-0.0496	
100	2019-08-19	1.61	2.77 40.8	585	1.63	42.8	-0.0173	-0.354	
101	2019-09-23	2	3.07 101	1190	2.09	125	-0.0886	-1.44	

Definitions

Cook's D: Cook's distance (Helsel and others, 2020).

DFFITS: Difference in fits statistic (Helsel and others, 2020).

E.vars: Explanatory variables.

Leverage: An outlier's measure in the x direction (Helsel and others, 2020).

LOESS: Local polynomial regression fitting, or locally estimated scatterplot smoothing (Helsel and others, 2020).

LOWESS: Locally weighted scatterplot smoothing (Cleveland, 1979; Helsel and others, 2020).

MSE: Model standard error (Helsel and others, 2020).

MSPE: Model standard percentage error (Helsel and others, 2020).

Na: Sodium, in milligrams per liter (00930).

Probability(>|t|): The probability that the independent variable has no effect on the dependent variable (Helsel and others, 2020).

RMSE: Root mean square error (Helsel and others, 2020).

SC: Specific conductance, in microsiemens per centimeter at 25 degrees Celsius (00095).

t value: Student's t value; the coefficient divided by its associated standard error (Helsel and others, 2020).

References Cited

- Cleveland, W.S., 1979, Robust locally weighted regression and smoothing scatterplots: Journal of the American Statistical Association, v. 74, no. 368, p. 829-836.
- Duan, N., 1983, Smearing estimate—A nonparametric retransformation method: Journal of the American Statistical Association, v. 78, no. 383, p. 605–610. [Also available at https://doi.org/10.1080/01621459.1983.10478017.]
- Foster, G.M., and Graham, J.L., 2016, Logistic and linear regression model documentation for statistical relations between continuous real-time and discrete water-quality constituents in the Kansas River, Kansas, July 2012 through June 2015: U.S. Geological Survey Open-File Report 2016–1040, 27 p., accessed July 2020 at https://doi.org/10.3133/ofr20161040.
- Graham, J.L., Foster, G.M., Williams, T.J., Mahoney, M.D., May, M.R., and Loftin, K.A., 2018, Water-quality conditions with an emphasis on cyanobacteria and associated toxins and taste-and-odor compounds in the Kansas River, Kansas, July 2012 through September 2016: U.S. Geological Survey Scientific Investigations Report 2018–5089, 55 p. [Also available at https://doi.org/10.3133/sir20185089.]
- Helsel, D.R., Hirsch, R.M., Ryberg, K.R., Archfield, S.A., and Gilroy, E.J., 2020, Statistical methods in water resources: U.S. Geological Survey Techniques and Methods, book 4, chap. A3, 458 p. [Also available at https://doi.org/10.3133/tm4a3.] [Supersedes USGS Techniques of Water-Resources Investigations, book 4, chap. A3, ver. 1.1.]
- Hem, J.D., 1992, Study and interpretation of the chemical characteristics of natural water (3d. ed): U.S. Geological Survey Water-Supply Paper 2254, 264 p. [Also available at https://doi.org/10.3133/wsp2254.]
- R Core Team, 2020, R—A language and environment for statistical computing, version 4.0.3: Vienna, Austria, R Foundation for Statistical Computing, accessed December 2020 at https://www.R-project.org/.
- Rasmussen, P.P., Gray, J.R., Glysson, G.D., and Ziegler, A.C., 2009, Guidelines and procedures for computing time-series suspended-sediment concentrations and loads from in-stream turbidity sensor and streamflow data: U.S. Geological Survey Techniques and Methods, book 3, chap. C4, 53 p. [Also available at https://doi.org/10.3133/tm3C4.]

- Rasmussen, T.J., Ziegler, A.C., and Rasmussen, P.P., 2005, Estimation of constituent concentrations, densities, loads, and yields in lower Kansas River, northeast Kansas, using regression models and continuous water-quality monitoring, January 2000 through December 2003: U.S. Geological Survey Scientific Investigations Report 2005–5165, 117 p. [Also available at https://doi.org/10.3133/sir20055165.]
- U.S. Geological Survey, 2020, USGS water data for the Nation: U.S. Geological Survey National Water Information System database, accessed April 2020 at https://doi.org/10.5066/F7P55KJN.
- U.S. Geological Survey, variously dated, National field manual for the collection of water-quality data: U.S. Geological Survey Techniques of Water-Resources Investigations, book 9, chaps. A1–A9 [variously paged], accessed July 2020 at https://water.usgs.gov/owq/FieldManual/.
- Wagner, R.J., Boulger, R.W., Jr., Oblinger, C.J., and Smith, B.A., 2006, Guidelines and standard procedures for continuous water-quality monitors—Station operation, record computation, and data reporting: U.S. Geological Survey Techniques and Methods, book 1, chap. D3, 51 p. plus 8 attachments. [Also available at https://doi.org/10.3133/tm1D3.] [Supersedes USGS Water-Resources Investigations Report 2000–4252.]