

Prepared in cooperation with the Navajo Nation and Peabody Western Coal Company

Groundwater, Surface-Water, and Water-Chemistry Data, Black Mesa Area, Northeastern Arizona—2018–2019

Open-File Report 2022–1086

U.S. Department of the Interior
U.S. Geological Survey

Cover. Aerial view of Moenkopi Wash looking west. Prominent light colored cliffs are outcrops of Entrada Sandstone located about 20-miles east (upstream) of Tuba City and Moenkopi. Photograph taken in December 2015 by Jon Mason.

Groundwater, Surface-Water, and Water-Chemistry Data, Black Mesa Area, Northeastern Arizona—2018–2019

By Jon P. Mason

Prepared in cooperation with the Navajo Nation and Peabody Western Coal Company

Open-File Report 2022–1086

**U.S. Department of the Interior
U.S. Geological Survey**

U.S. Geological Survey, Reston, Virginia: 2022

For more information on the USGS—the Federal source for science about the Earth, its natural and living resources, natural hazards, and the environment—visit <https://www.usgs.gov/> or call 1–888–ASK–USGS (1–888–275–8747).

For an overview of USGS information products, including maps, imagery, and publications, visit <https://store.usgs.gov>.

Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Although this information product, for the most part, is in the public domain, it also may contain copyrighted materials as noted in the text. Permission to reproduce copyrighted items must be secured from the copyright owner..

Suggested citation:

Mason, J.P., 2022, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2018–2019: U.S. Geological Survey Open-File Report 2022–1086, 47 p., <https://doi.org/10.3133/ofr20221086>.

Contents

Abstract.....	1
Introduction.....	1
Purpose and Scope	4
Previous Investigations.....	6
Description of Study Area	9
Physiography.....	9
Climate	10
Geology.....	10
Geologic Units Below the N Aquifer.....	10
Geologic Units of the N Aquifer.....	10
Moenave Formation	12
Wingate Sandstone	12
Kayenta Formation.....	12
Navajo Sandstone	13
Carmel Formation.....	13
Geologic Units of the D Aquifer	13
Entrada Sandstone	13
Morrison Formation	13
Dakota Sandstone	14
Mancos Shale	14
Geologic Units of the T Aquifer	14
Mesaverde Group.....	14
Bidahochi Formation	14
Hydrologic Data.....	14
Withdrawals from the N Aquifer	16
Withdrawals in Calendar Year 2019 Compared to Previous Years	16
Groundwater Levels in the N Aquifer.....	16
Spring Discharge from the N Aquifer.....	26
Surface-Water Discharge, Calendar Year 2019.....	31
Moenkopi Wash	31
Dinnebito Wash.....	31
Polacca Wash	32
Pasture Canyon Springs	33
Surface-Water Base Flow.....	33
Water Chemistry	33
Water-Chemistry Data for Wells Completed in the N Aquifer.....	35
Water-Chemistry Data for Springs that Discharge from the N Aquifer	35
Summary.....	41
References Cited.....	43

Figures

1. Map showing location of study area, Black Mesa area, northeastern Arizona	3
2. Stratigraphic section showing rock formations and hydrogeologic units of the Black Mesa area, northeastern Arizona.....	5
3. Aerial photograph of Organ Rock monocline and folding strata of Skeleton Mesa near Kayenta, Arizona	9
4. Map showing mean annual precipitation, Black Mesa area, Arizona, 1981–2010.....	11
5. Aerial photograph of Moenave Formation outcropping on Garces Mesas, northeastern Arizona.....	12
6. Plot of annual withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, 1965–2019.....	17
7. Map of well systems monitored for annual withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, calendar year 2019, and confined/unconfined zones of the N aquifer.....	18
8. Map showing water-level changes in wells completed in the N aquifer, Black Mesa area, northeastern Arizona, prestress period (prior to 1965) to 2019.....	20
9. Plots of observed water levels in annual observation well network of the N aquifer, Black Mesa area, northeastern Arizona, 1950–2019	23
10. Plots of observed groundwater levels in continuous-record observation wells BM1–BM6 in the N aquifer, Black Mesa area, northeastern Arizona, 1963–2019.....	27
11. Map of surface-water and water-chemistry data-collection sites in the N aquifer, Black Mesa area, northeastern Arizona, 2019.....	28
12. Plots of discharge from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso, N Aquifer, Black Mesa area, northeastern Arizona, 1987–2019.....	29
13. Plots of daily mean discharge for Moenkopi Wash at Moenkopi, Dinnebito Wash near Sand Springs, Polacca Wash near Second Mesa, and Pasture Canyon Springs, Black Mesa area, northeastern Arizona, calendar year 2019.....	32
14. Plots of median winter discharge for November through February for Moenkopi Wash at Moenkopi, Dinnebito Wash near Sand Springs, Polacca Wash near Second Mesa, and Pasture Canyon Springs, Black Mesa area, northeastern Arizona, winter 1977–2018	34
15. Map showing water chemistry and distribution of dissolved solids at wells and springs in the N aquifer, Black Mesa area, northeastern Arizona, 2019	36
16. Plot of dissolved-solids concentrations for water samples from selected wells completed in the N aquifer, Black Mesa area, northeastern Arizona	37
17. Plots of concentrations of dissolved solids, chloride, and sulfate for water samples from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso, which discharge from the N aquifer, Black Mesa area, northeastern Arizona, 1982–2019.....	42

Tables

1. Withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, 1965–2019.....	6
2. Tabulated list of progress reports for the Black Mesa monitoring program, 1978–2021	7
3. Identification numbers and names of monitoring-program wells used for water-level measurements or for collection of water-chemistry samples, Black Mesa area, northeastern Arizona, 2018–19	15
4. Withdrawals from the N aquifer by well system, Black Mesa area, northeastern Arizona, calendar year 2019	17
5. Total, industrial, and municipal withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, for discrete time periods from 1965 to 2019	17
6. Water-level changes in monitoring program wells completed in the N aquifer, Black Mesa area, northeastern Arizona, prestress period (prior to 1965) to calendar year 2019.....	19
7. Well-construction characteristics, depth to top of N aquifer, and 2019 static water level for wells used in annual water-level measurements and for continuous-record observation wells, Black Mesa area, northeastern Arizona, 2018–19	21
8. Median changes in water levels in monitoring-program wells of the N aquifer, Black Mesa area, northeastern Arizona, prestress period (prior to 1965) to 2019.....	22
9. Discharge from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso, N Aquifer, Black Mesa area, northeastern Arizona, 1952–2019.....	30
10. Streamflow-gaging stations used in the Black Mesa monitoring program, their periods of record, and drainage areas	32
11. Comparison of chemical analyses of environmental and replicate water samples taken from Moenkopi School Spring (360632111131101), Black Mesa area, northeastern Arizona, 2019	35
12. Physical properties and chemical analyses of water samples from selected industrial wells completed in the N aquifer, Black Mesa area, northeastern Arizona, 2019.....	37
13. Specific conductance and concentrations of selected chemical constituents in water samples from selected industrial wells completed in the N aquifer, Black Mesa area, northeastern Arizona, 1968–2019.....	38
14. Physical properties and chemical analyses of water samples from four springs that discharge from the N aquifer, Black Mesa area, northeastern Arizona, 2019	39
15. Specific conductance and concentrations of selected chemical constituents in water samples from four springs that discharge from the N aquifer, Black Mesa area, northeastern Arizona, 1948–2019	40

Conversion Factors

U.S. customary units to International System of Units

Multiply	By	To obtain
Length		
inch (in.)	2.54	centimeter (cm)
inch (in.)	25.4	millimeter (mm)
foot (ft)	0.3048	meter (m)
mile (mi)	1.609	kilometer (km)
Area		
square mile (mi ²)	259.0	hectare (ha)
square mile (mi ²)	2.590	square kilometer (km ²)
Volume		
acre-foot (acre-ft)	1,233	cubic meter (m ³)
acre-foot (acre-ft)	0.001233	cubic hectometer (hm ³)
Flow rate		
acre-foot per year (acre-ft/yr)	1,233	cubic meter per year (m ³ /yr)
acre-foot per year (acre-ft/yr)	0.001233	cubic hectometer per year (hm ³ /yr)
cubic foot per second (ft ³ /s)	0.02832	cubic meter per second (m ³ /s)
gallon per minute (gal/min)	0.06309	liter per second (L/s)

Temperature in degrees Celsius (°C) may be converted to degrees Fahrenheit (°F) as °F = (1.8 × °C) + 32.

Datum

Vertical coordinate information is referenced to the National Geodetic Vertical Datum of 1929 (NGVD 29).

Horizontal coordinate information is referenced to the North American Datum of 1927 (NAD 27).

Supplemental Information

Specific conductance is given in microsiemens per centimeter at 25 degrees Celsius (µS/cm at 25 °C).

Concentrations of chemical constituents in water are given in either milligrams per liter (mg/L) or micrograms per liter (µg/L).

Abbreviations

ADWR	Arizona Department of Water Resources
BIA	Bureau of Indian Affairs
C aquifer	Coconino aquifer
D aquifer	Dakota aquifer
EPA	U.S. Environmental Protection Agency
MCL	Maximum Contaminate Level
N aquifer	Navajo aquifer
NTUA	Navajo Tribal Utility Authority
NWIS	National Water Information System
PWCC	Peabody Western Coal Company
QC	Quality control
SMCL	Secondary Maximum Contaminate Level
T aquifer	Toreva aquifer
USGS	U.S. Geological Survey

Groundwater, Surface-Water, and Water-Chemistry Data, Black Mesa Area, Northeastern Arizona—2018–2019

By Jon P. Mason

Abstract

The Navajo (N) aquifer is an extensive aquifer and the primary source of groundwater in the 5,400-square-mile Black Mesa area in northeastern Arizona. Water availability is an important issue in the Black Mesa area because of the arid climate, past industrial water use, and continued water requirements for municipal use by a growing population. Precipitation in the area typically ranges from less than 6 to more than 16 inches per year depending on location.

The U.S. Geological Survey water-monitoring program in the Black Mesa area began in 1971 and provides information about the long-term effects of groundwater withdrawals from the N aquifer for industrial and municipal uses. This report presents results of data collected as part of the monitoring program in the Black Mesa area from calendar year 2019, and additionally uses streamflow statistics from November and December 2018. The monitoring program includes measurements of (1) groundwater withdrawals (pumping), (2) groundwater levels, (3) spring discharge, (4) surface-water discharge, and (5) groundwater chemistry.

In calendar year 2019, total groundwater withdrawals were estimated to be 3,070 acre-feet (acre-ft), industrial withdrawals were 670 acre-ft, and municipal withdrawals were estimated to be 2,400 acre-ft. Total withdrawals during 2019 were about 58 percent less than total withdrawals in 2005 because of Peabody Western Coal Company's discontinued use of water to transport coal in a coal slurry pipeline after 2005 and cessation of mining operations in 2019.

Water levels measured in 2019 from wells completed in the unconfined areas of the N aquifer within the Black Mesa area showed a decline in 10 of 16 wells when compared with water levels from the prestress period (prior to 1965). The changes in water levels across all 16 wells ranged from +8.2 feet (ft) to -40.0 ft, and the median change was -1.7 ft. Water levels also showed decline in 16 of 18 wells measured in the confined area of the aquifer when compared to the prestress period. The median change for the confined area of the aquifer was -38.8 ft, with changes across all 18 wells ranging from +12.9 ft to -185.0 ft.

Spring flow was measured at four springs in 2019. Flow fluctuated during the period of record for Burro Spring and Pasture Canyon Spring, but a decreasing trend was statistically significant ($p < 0.05$) at Moenkopi School Spring and Unnamed Spring near Dennehotso. Discharge at Burro Spring has

remained relatively constant since it was first measured in the 1980s and discharge at Pasture Canyon Spring has fluctuated for the period of record.


Continuous records of surface-water discharge in the Black Mesa area were collected from streamflow-gaging stations at the following sites: Moenkopi Wash at Moenkopi 09401260 (1976 to 2019), Dinnebito Wash near Sand Springs 09401110 (1993 to 2019), Polacca Wash near Second Mesa 09400568 (1994 to 2019), and Pasture Canyon Springs 09401265 (2004 to 2019). Median winter flows (November through February) of each winter were used as an estimate of the amount of groundwater discharge at the above-named sites. For the period of record, the median winter flows have generally remained constant at Polacca Wash and Pasture Canyon Springs, whereas a decreasing trend was indicated at Moenkopi Wash and Dinnebito Wash.

In 2019, water samples collected from four springs and three wells in the Black Mesa area were analyzed for selected chemical constituents. Results from the four springs were compared with previous analyses from the same springs. Concentrations of dissolved solids, chloride, and sulfate increased at Moenkopi School Spring during the more than 30 years of record at that site. Concentrations of dissolved solids, chloride, and sulfate at Pasture Canyon Spring have not varied significantly ($p > 0.05$) since the early 1980s, and there is no increasing or decreasing trend in those data. Concentrations of dissolved solids, chloride, and sulfate at Unnamed Spring near Dennehotso have varied for the period of record, but there is no statistical trend in the data. Concentrations of dissolved solids and chloride at Burro Spring have varied for the period of record, but there is no statistical trend in the data; however, concentrations of sulfate from Burro Spring now show a trend towards lower concentrations. No statistical trend tests were performed for the three wells sampled in 2019 since less historical water-quality data were available for comparison.

Introduction


The 5,400-square-mile (mi²) Black Mesa study area is enclosed within the Navajo Nation and Hopi Reservation in northeastern Arizona (fig. 1). It contains diverse topography that includes flat plains, mesas, and incised drainages (fig. 1).

2 Groundwater, Surface-Water, and Water-Chemistry Data, Black Mesa Area, Northeastern Arizona—2018–2019



Base from U.S. Geological Survey digital data, 2010;
Lambert Conformal Conic projection, standard parallels
29°30' N. and 45°30' N., central meridian 111°30' W.,
North American Datum of 1927

0 5 10 15 20 25 MILES
0 5 10 15 20 25 KILOMETERS



EXPLANATION




-  **Area of Hopi Tribal Lands within Navajo Nation**
-  **Boundary of Black Mesa**
-  **Boundary of study area**—Based on the mathematical boundary of groundwater model from Brown and Eychaner (1988)



Figure 1. Map showing location of study area, Black Mesa area, northeastern Arizona. Boundary of study area is based on boundary of groundwater model from Eychaner (1983).

Modified from Brown and Eychaner, 1988

Black Mesa, a topographic high at the center of the study area, encompasses about 2,000 mi². It has 2,000-foot (ft)-high cliffs on its northern and northeastern sides, but it slopes gradually down to the south and southwest. Availability of water is an important issue in the study area because of continued groundwater withdrawals, the growing population, and an arid to semiarid climate.

Aquifers that are utilized in the Black Mesa area include the Toreva (T), Dakota (D), and Navajo (N) aquifers (fig. 2). Shallow aquifers composed of surficial sediments or volcanic rock also are used locally to supply small quantities of water. Water from the T and D aquifers are not used in significant quantities in the Black Mesa area. Water from the T aquifer is used locally for livestock watering and to irrigate small plots of land, but it likely cannot produce enough water for municipal or industrial use. Water from the D aquifer is used locally for livestock watering and has contributed to some wells at the Peabody Western Coal Company (PWCC) industrial well field, but water from the aquifer has elevated total-dissolved solids concentrations that make it unsuitable for municipal use. The deeper Coconino (C) aquifer is present throughout the Black Mesa area, but it is deeply buried and likely has total-dissolved solids concentrations above what can be used without treatment. The N aquifer, lying between the D and C aquifers, is the major source of water for industrial and municipal uses in the Black Mesa area. For this reason, groundwater data collected for this report was exclusively from the N aquifer.

According to Eychaner (1983), the N aquifer is composed of three hydraulically connected formations—the Navajo Sandstone, the Kayenta Formation, and the Wingate Sandstone—that function as a single aquifer (fig. 2). However, more recent geologic mapping indicates the Wingate Sandstone is absent from much of the Black Mesa area. Outcrops of sandstone previously mapped as Wingate Sandstone in the Black Mesa area are now considered to be part of the Moenave Formation (Billingsley and others, 2012, 2013). Based on this recent geologic mapping, it is unclear if the Wingate Sandstone is present at all in the Black Mesa area. If present, it would only be in the northeastern part of the study area where it would be deeply buried. The N aquifer is confined under most of Black Mesa, and the overlying stratigraphy limits recharge to this part of the aquifer. The N aquifer is unconfined in most areas surrounding Black Mesa, and most recharge occurs where the Navajo Sandstone is exposed in the area near Shonto (fig. 1) (Lopes and Hoffmann, 1997). From the recharge areas near Shonto, groundwater moves radially to the southwest toward Tuba City, to the south toward the Hopi Reservation, and to the east toward Rough Rock and Dennehotso (Eychaner, 1983).

Within the Black Mesa study area, the Navajo Nation and Hopi Tribe are the principal municipal water users, and the PWCC is the principal industrial water user. Withdrawals from the N aquifer in the Black Mesa area increased fairly consistently from 1965 through 2005 and then decreased markedly in 2006 (table 1). The PWCC began operating a strip mine in the northern part of the study area in 1968 (fig. 1). The PWCC's mining operation consisted of two mines on Black Mesa—the Kayenta mine, which transported coal to the Navajo Generating Station by train, and the Black Mesa mine, which transported coal 275 miles (mi) to the Mohave Generating Station by a water-based coal slurry pipeline.

The PWCC operated both mines on Black Mesa from the 1970s until about 2005, when the Mohave Generating Station ceased operations. On December 31, 2005, the PWCC reduced pumping of the N aquifer by approximately 70 percent as a result of discontinued use of the coal slurry pipeline that delivered water, in addition to coal, to the Mohave Generating Station. The two mines at the PWCC were then combined into the Black Mesa Complex and continued to deliver coal to the Navajo Generating Station by electric train until 2019. In August 2019, coal mining operations at the Black Mesa Complex ceased due to the planned closure of the Navajo Generating Station which was permanently closed in November 2019. The PWCC continued to pump about 1,100 to 1,600 acre-feet (acre-ft) per year between 2006 and 2018, primarily for dust control. In 2019 the PWCC reduced its annual pumping to 670 acre-ft (table 1).

There are four major stream systems that provide surface drainage for the Black Mesa area. They are Moenkopi Wash, Dinnebito Wash, Oraibi Wash, and Polacca Wash. All four stream systems have headwaters high on Black Mesa and eventually drain into the Little Colorado River to the south and southwest of the study area (fig. 1). Most reaches of these streams are ephemeral, flowing only in response to runoff from precipitation events, but a few short reaches flow at least part of each year as a result of groundwater discharge.

The members of the Navajo Nation and the Hopi Tribe have been concerned about the long-term effects of withdrawals from the N aquifer on available groundwater supplies, on stream and spring discharge, and on groundwater chemistry. In 1971, these water-supply concerns led to the establishment of a monitoring program for the water resources in the Black Mesa area by the U.S. Geological Survey (USGS) in cooperation with the Arizona Water Commission, which was the predecessor to the present Arizona Department of Water Resources (ADWR). In 1983, the Bureau of Indian Affairs (BIA) joined the cooperative effort. Since 1983, the Navajo Tribal Utility Authority (NTUA); the PWCC; the Hopi Tribe; and the Western Navajo, Chinle, and Hopi Agencies of the BIA have assisted in the collection of hydrologic data.

Purpose and Scope

This report presents results of groundwater, surface-water, and water-chemistry monitoring in the Black Mesa area from January 2019 to December 2019. Additionally, surface-water statistics are used from November and December 2018. Continuous and periodic groundwater and surface-water data are collected to monitor the possible effects of industrial and municipal withdrawals from the N aquifer on groundwater levels, stream and spring discharge, and groundwater chemistry. Groundwater data include groundwater withdrawals (pumping), water levels, spring-discharge rates, and water chemistry. Surface-water data include discharge rates at four continuous-record streamflow-gaging stations. Together, these data are compared with groundwater and surface-water data from 1965 to 2019 to describe the overall status of and change over time of groundwater conditions in the N aquifer, as well as to provide information on how the aquifer responds to groundwater development stresses. Some statistical analyses of the data are included in this report.




Figure 2. Stratigraphic section showing rock formations and hydrogeologic units of the Black Mesa area, northeastern Arizona (not to scale). The Navajo (N) aquifer is approximately 1,000 feet thick.

Table 1. Withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, 1965–2019.

[Values are rounded to nearest 10 acre-feet. Data for 1965–79 from Eychaner (1983). Total withdrawals in Littin and Monroe (1996) were for the confined area of the aquifer]

Calendar Year	Industrial Withdrawals ^a	Municipal withdrawals ^{b,c}		Total withdrawals	Calendar Year	Industrial Withdrawals ^a	Municipal withdrawals ^{b,c}		Total withdrawals
		Confined areas	Unconfined areas				Confined areas	Unconfined areas	
		(acre-feet)					(acre-feet)		
1965	0	50	20	70	1993	3,700	1,250	1,570	6,520
1966	0	110	30	140	1994	4,080	1,210	1,600	6,890
1967	0	120	50	170	1995	4,340	1,220	1,510	7,070
1968	100	150	100	350	1996	4,010	1,380	1,650	7,040
1969	40	200	100	340	1997	4,130	1,380	1,580	7,090
1970	740	280	150	1,170	1998	4,030	1,440	1,590	7,060
1971	1,900	340	150	2,390	1999	4,210	1,420	1,480	7,110
1972	3,680	370	250	4,300	2000	4,490	1,610	1,640	7,740
1973	3,520	530	300	4,350	2001	4,530	1,490	1,660	7,680
1974	3,830	580	360	4,770	2002	4,640	1,500	1,860	8,000
1975	3,500	600	510	4,610	2003	4,450	1,350	1,440	7,240
1976	4,180	690	640	5,510	2004	4,370	1,240	1,600	7,210
1977	4,090	750	730	5,570	2005	4,480	1,280	1,570	7,330
1978	3,000	830	930	4,760	2006	1,200	^d 1,300	^d 1,600	^d 4,100
1979	3,500	860	930	5,290	2007	1,170	1,460	1,640	4,270
1980	3,540	910	880	5,330	2008	1,210	^e 1,430	^e 1,560	4,200
1981	4,010	960	1,000	5,970	2009	1,390	1,440	1,400	4,230
1982	4,740	870	960	6,570	2010	1,170	^d 1,450	1,420	^d 4,040
1983	4,460	1,360	1,280	7,100	2011	1,390	^d 1,460	1,630	^d 4,480
1984	4,170	1,070	1,400	6,640	2012	1,370	^d 1,380	1,260	^d 4,010
1985	2,520	1,040	1,160	4,720	2013	1,460	^d 1,410	^d 1,110	^d 3,980
1986	4,480	970	1,260	6,710	2014	1,580	^d 1,280	^d 1,310	^d 4,170
1987	3,830	1,130	1,280	6,240	2015	1,340	^d 1,370	^d 1,260	^d 3,970
1988	4,090	1,250	1,310	6,650	2016	1,090	^d 1,380	^d 1,070	^d 3,540
1989	3,450	1,070	1,400	5,920	2017	1,110	1,330	^d 1,270	^d 3,710
1990	3,430	1,170	1,210	5,810	2018	1,170	^d 1,370	1,130	^d 3,670
1991	4,020	1,140	1,300	6,460	2019	670	^d 1,340	^d 1,060	^d 3,070
1992	3,820	1,180	1,410	6,410					

^aMetered pumpage from the confined part of the aquifer by Peabody Western Coal Company.^bDoes not include withdrawals from the wells equipped with windmills.^cIncludes estimated pumpage 1965–73 and metered pumpage 1974–79 at Tuba City; metered pumpage at Kayenta and estimated pumpage at Chilchinbito, Rough Rock, Piñon, Keams Canyon, and Kykotsmobi before 1980; metered and estimated pumpage furnished by the Navajo Tribal Utility Authority and the Bureau of Indian Affairs and collected by the U.S. Geological Survey, 1980–85; and metered pumpage furnished by the Navajo Tribal Utility Authority, the Bureau of Indian Affairs, various Hopi Village Administrations, and the U.S. Geological Survey, 1986–2019.^dMetered data were incomplete; therefore, municipal withdrawals are estimated, and total withdrawals included an estimation in the calculations.^eConfined and unconfined area totals were reported incorrectly in Open-File Report 2010–1038 and 2011–1198.

to examine trends in the data that characterize groundwater conditions in the N aquifer.

Previous Investigations

Progress reports on the Black Mesa area monitoring program have been prepared by the USGS since 1978, and these progress reports are summarized in table 2. The groundwater-level,

surface-water discharge, and water-chemistry data from the Black Mesa area monitoring program are contained in these progress reports and in the USGS National Water Information System (NWIS) database (<http://waterdata.usgs.gov/az/nwis/>). Water-withdrawal data are presented in tables in the progress reports.

Stream-discharge and periodic water-quality data collected from Moenkopi Wash before the 1982 water year were published by the USGS (1963–64a,b; 1965–74a,b; and 1976–83). Stream-discharge data from water years 1983 to 2005 for Moenkopi Wash

Table 2. Tabulated list of progress reports for the Black Mesa monitoring program, 1978–2021.

Year published	Author(s)	Title	USGS report type and number
1978	U.S. Geological Survey	Progress report on Black Mesa monitoring program—1977	Open-File Report 78–459
1985	Hill, G.W.	Progress report on Black Mesa monitoring program—1984	Open-File Report 85–483
1986	Hill, G.W., and Whetten, M.I.	Progress report on Black Mesa monitoring program—1985–86	Open-File Report 86–414
1987	Hill, G.W., and Sottolare, J.P.	Progress report on the ground-water, surface-water, and quality-of-water monitoring program, Black Mesa area, northeastern Arizona—1987	Open-File Report 87–458
1988	Hart, R.J., and Sottolare, J.P.	Progress report on the ground-water, surface-water, and quality-of-water monitoring program, Black Mesa area, northeastern Arizona—1987–88	Open-File Report 88–467
1989	Hart, R.J., and Sottolare, J.P.	Progress report on the ground-water, surface-water, and quality-of-water monitoring program, Black Mesa area, northeastern Arizona—1988–89	Open-File Report 89–383
1992	Sottolare, J.P.	Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1989–90	Water-Resources Investigations Report 92–4008
1992	Littin, G.R.	Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1990–91	Water-Resources Investigations Report 92–4045
1993	Littin, G.R.	Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1991–92	Water-Resources Investigations Report 93–4111
1995a	Littin, G.R., and Monroe, S.A.	Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1992–93	Water-Resources Investigations Report 95–4156
1995b	Littin, G.R., and Monroe, S.A.	Results of ground-water, surface-water, and water-chemistry monitoring, Black Mesa area, northeastern Arizona—1994	Water-Resources Investigations Report 95–4238
1996	Littin, G.R., and Monroe, S.A.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1995	Open-File Report 96–616
1997	Littin, G.R., and Monroe, S.A.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1996	Open-File Report 97–566
1999	Littin, G.R., Baum, B.M., and Truini, M.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1997	Open-File Report 98–653
2000	Truini, M., Baum, B.M., Littin, G.R., and Shingoitewa-Honanie, G.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1998	Open-File Report 00–66
2000	Thomas, B.E., and Truini, M.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1999	Open-File Report 00–453
2002a	Thomas, B.E.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2000–2001, and performance and sensitivity of the 1988 USGS numerical model of the N aquifer	Water-Resources Investigations Report 02–4211
2002b	Thomas, B.E.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2001–02	Open-File Report 02–485
2004	Truini, M., and Thomas, B.E.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2002–03	Open-File Report 03–503
2005	Truini, M., Macy, J.P., and Porter, T.J.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2003–04	Open-File Report 2005–1080

Table 2. Tabulated list of progress reports for the Black Mesa monitoring program, 1978–2019.—Continued

Year published	Author(s)	Title	USGS report type and number
2006	Truini, M., and Macy, J.P.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2004–05	Open-File Report 2006–1058
2007	Truini, M., and Macy, J.P.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2005–06	Open-File Report 2007–1041
2008	Truini, M., and Macy, J.P.	Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2006–07	Open-File Report 2008–1324
2009	Macy, J.P.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2007–2008	Open-File Report 2009–1148
2010	Macy, J.P.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2008–2009	Open-File Report 2010–1038
2011	Macy, J.P., and Brown, C.R.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2009–2010	Open-File Report 2011–1198
2012	Macy, J.P., Brown, C.R., and Anderson, J.R.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2010–2011	Open-File Report 2012–1102
2014	Macy, J.P., and Unema, J.A.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2011–2012	Open-File Report 2013–1304
2016	Macy, J.P., and Truini, M.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2012–2013	Open-File Report 2015–1221
2017	Macy, J.P., and Mason, J.P.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2013–2015	Open-File Report 2017–1127
2018	Mason, J.P., and Macy, J.P.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2015–2016	Open-File Report 2018–1193
2021	Mason, J.P.	Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2016–2018	Open-File Report 2021–1124

at Moenkopi (09401260), Dinnebito Wash near Sand Springs (09401110), Polacca Wash near Second Mesa (09400568), Laguna Creek at Dennehotso (09379180), and Pasture Canyon Springs (09401265) in the Black Mesa area were published in White and Garrett (1984, 1986, 1987, 1988), Wilson and Garrett (1988, 1989), Boner and others (1989, 1990, 1991, 1992), Smith and others (1993, 1994, 1995, 1996, 1997), Tadayon and others (1998, 1999, 2000, 2001), McCormack and others (2002, 2003), Fisk and others (2004, 2005, 2006), and online for year 2006 to present (<http://wdr.water.usgs.gov>). Before the monitoring program, a large data-collection effort in the 1950s resulted in a compilation of well and spring data for the Navajo Nation and Hopi Reservation (Davis and others, 1963).

Many interpretive studies have investigated the hydrology of the Black Mesa area. Cooley and others (1969) made the first comprehensive evaluation of the regional hydrogeology of the Black Mesa area. Eychaner (1983) developed a two-dimensional numerical model of groundwater flow in the N aquifer. Brown and Eychaner (1988) recalibrated Eychaner's model by using a finer grid and by using revised estimates of selected aquifer characteristics. GeoTrans, Inc. (1987) also developed a two-dimensional numerical

model of the N aquifer in the 1980s. In the late 1990s, HSI GeoTrans, Inc., and Waterstone Environmental Hydrology and Engineering, Inc. (1999) developed a three-dimensional numerical model of the N aquifer and the overlying D aquifer.

Kister and Hatchett (1963) made the first comprehensive evaluation of the chemistry of water collected from wells and springs in the Black Mesa area. HSI GeoTrans, Inc. (1993) evaluated the major-ion and isotopic chemistry of the D and N aquifers. Lopes and Hoffmann (1997) analyzed groundwater ages, recharge, and hydraulic conductivity of the N aquifer by using geochemical techniques. Zhu and others (1998) estimated groundwater recharge in the Black Mesa area by using isotopic data and flow estimates from the N-aquifer model developed by GeoTrans, Inc. (1987). Zhu (2000) estimated recharge using advective transport modeling and the same isotopic data from the GeoTrans model. Truini and Longworth (2003) described the hydrogeology of the D aquifer and the movement and ages of groundwater in the Black Mesa area by using data from geochemical and isotopic analyses. Truini and Macy (2005) looked at possible leakage through the confining unit between the D aquifer and the N aquifer as part of an investigation of the Carmel Formation.

Description of Study Area

The availability and chemistry of water resources within the Black Mesa area are directly related to physiography, climate, and geology. Physiography affects the movement of both surface water and groundwater in the area, and climate affects the water budget. The complex geologic history of the area has resulted in the accumulation of abundant coal resources and also influences the movement and chemistry of surface water and groundwater.

Physiography

The Black Mesa area is located in the Colorado Plateaus physiographic province of the Intermontane Plateaus Region (Raisz, 1972). The dominant physiographic feature in the study area is Black Mesa itself, but numerous smaller features play an important role in the movement of surface water and groundwater (fig. 1). Black Mesa is the remnant of a large sedimentary basin that has undergone significant tectonic deformation and uplift during the past 70 million years. Parts of Black Mesa which were once below sea level now rise over 6,000 ft above sea level. As a result of this uplift, the region has gone from a depositional cycle to an erosional cycle. Much of the erosion responsible for

present day topography likely occurred in the past 10 million years (Lazear and others, 2013). Since uplift occurred, Black Mesa has been dissected by streams, resulting in the formation of numerous smaller mesas such as the Hopi Mesas.

The geologic units that comprise the N aquifer occur at or near the land surface in a large extent around the periphery of Black Mesa. In these areas the aquifer is generally unconfined. West of Kayenta, exposed N aquifer units form Skeleton Mesa and the Shonto Plateau. At the southeast edges of these features, the aquifer units are folded by the Organ Rock monocline (fig. 3) and plunge steeply to the southeast below the younger Cretaceous rocks of Black Mesa forming Long House Valley. The N-aquifer units continue to the southeast under Black Mesa eventually reappearing south of the Hopi mesas. The aquifer units pinch out not far from where they reappear. In general, the confined portion of the N aquifer occurs where the aquifer units are deeply buried beneath Black Mesa.

The paths of stream channels also are influenced by physiography. Geologic structural folds, joint patterns, rock type, and topography all affect the flow of surface water in the study area. Major streams of the study area are shown in figure 1. The surface topography of Black Mesa slopes downhill from northeast to southwest. Likewise, the major streams that drain Black Mesa flow from northeast to southwest toward the Little Colorado River.



Figure 3. Aerial photograph of Organ Rock monocline and folding strata of Skeleton Mesa near Kayenta, Arizona. The Navajo Sandstone is truncated in this part of the monocline, forming the flatirons along the lower part of the monocline. Photograph by Jodi Norris.

Climate

The climate in most of the Black Mesa area is broadly classified by Hendricks (1985) as steppe, which is characterized by limited amounts of precipitation. Much of the precipitation in steppe regions evaporates before it can infiltrate to groundwater. As a result, the vegetation cover consists mostly of mesquite, pinyon-juniper, and various grasses (Hendricks, 1985). A small area around Tuba City is classified by Hendricks (1985) as desert, signifying even less annual rainfall and a vegetative cover consisting mostly of creosote bush, cacti, and sagebrush.

Mean annual precipitation for the Black Mesa area was estimated using spatial regression methods that incorporated precipitation data from traditional weather stations and high-altitude meteorological sites (Daly and others, 1994). Based on 30-year averages from 1981–2010, precipitation in the Black Mesa area ranges from less than 6 inches (in.) in the lower elevation regions around the mesa to more than 16 in. at the highest elevations on the mesa (fig. 4; PRISM Climate Group, 2018).

According to Sellers and Hill (1974), about 60 percent of average annual precipitation in northeastern Arizona falls between the months of May and October (primarily in July and August). They report that, on average, the plateaus and mesas of northeastern Arizona are the driest part of the state during the colder half of the year and rarely receive heavy winter precipitation. Using more recent precipitation data, an analysis of 30-year normal precipitation for the period 1981–2010 (PRISM Climate Group, 2018) in the Black Mesa area show that about 55 percent of precipitation occurred from May–October. Much of the groundwater contained in the N aquifer was recharged during the late Pleistocene when the temperature was cooler and precipitation amounts were higher (Zhu and Kipfer, 2010).

Geology

The stratigraphic section (fig. 2) used in the current and previous Black Mesa monitoring reports was modified from Harshbarger and others (1966). The original stratigraphic section showed the Wingate Sandstone between the Chinle Formation and the Kayenta Formation and did not have the Moenave Formation present. More recently, Billingsley and others (2012, 2013) concluded that sandstones in the Black Mesa area formerly mapped in outcrop as Wingate Sandstone are in fact part of the Moenave Formation. It is unclear if the Wingate Sandstone could be present in the subsurface under parts of the Black Mesa area. Since the two geologic units are considered coeval, the Moenave Formation is shown as present and possibly intertongued with the Wingate Sandstone in figure 2. Harshbarger and others (1966) considered the eolian facies of the Wingate Sandstone to be a water-bearing unit of the N aquifer. It is unclear if any of the sandstones now mapped as Moenave Formation could be water bearing.

Rocks of Triassic age and older are not discussed in detail in this report because they are not significant sources

of groundwater in the Black Mesa area. Instead, this section focusses on Jurassic and younger rocks that are part of hydrologic systems used in this area.

The Black Mesa area is the remnant of a large sedimentary basin that has been uplifted and dissected by streams since its original formation. When the sedimentary rock units (fig. 2) in the Black Mesa area were deposited the region had a much lower surface elevation nearer to, and sometimes below, sea level. As the thick sequence of sedimentary rock units was being deposited, the basin was slowly subsiding, allowing more sediments from nearby highlands to be deposited. The entire Colorado Plateau including Black Mesa was then tectonically uplifted 1 mi above sea level during the Tertiary by processes that are still not fully understood. According to Flowers (2010), Colorado Plateau “elevation gain could have occurred in the early Tertiary associated with Sevier-Laramide contraction, the middle Tertiary synchronous with the proposed demise of the Laramide flat slab, [or] the late Tertiary coeval with regional extensional tectonism in adjacent provinces.”

Geologic Units Below the N Aquifer

The geologic units below the N-aquifer system are Triassic and older in age (fig. 2) and generally are not suitable as a water supply in the Black Mesa area and will not be discussed in detail. The Permian Coconino Sandstone and Kaibab Limestone (fig. 2) can produce adequate quantities of water in the Black Mesa area, but they are deeply buried and likely have total-dissolved solids concentrations above what can be used without treatment.

Geologic Units of the N Aquifer

The geologic units associated with the N aquifer are members of the Glen Canyon Group and include the Moenave Formation, Wingate Sandstone, Kayenta Formation, and Navajo Sandstone (fig. 2). The group is named after Glen Canyon of the Colorado River in southeastern Utah where these units are typically exposed (Harshbarger and others, 1957). The Glen Canyon Group was originally thought to be Late Triassic to Early Jurassic in age (Harshbarger and others, 1957), but more recent paleontological and stratigraphic discoveries strongly suggest the group is largely Early Jurassic in age (Peterson and Pipiringos, 1979). According to Blakey and Ranney (2008), when the Glen Canyon Group was deposited, the Black Mesa basin was slightly above sea level, and the climate was windy and dry. This led to widespread deposition of eolian and fluvial deposits (Blakey and Ranney, 2008) that now compose the sandstone units of the N aquifer.

Where the N aquifer is confined it is capped by the Carmel Formation (fig. 2), which is considered part of the San Rafael Group; the Carmel Formation is discussed in this section since it both confines the aquifer in places and hydraulically separates the N aquifer from the overlying D aquifer (where the D aquifer is present).





Figure 4. Map showing mean annual precipitation, Black Mesa area, Arizona, 1981-2010.

Moenave Formation

The Moenave Formation (fig. 2) contains several members with the most prominent one in the Black Mesa area being the Dinosaur Canyon Member. Blakey and Ranney (2008) described the Moenave Formation as being deposited by northwest flowing streams at the same time the Wingate Sandstone was being deposited to the northeast. Billingsley and others (2012) described the lithology of the formation as reddish-brown, thin-bedded, flat-bedded, and crossbedded, fine- to coarse-grained fluvial siltstone and silty sandstone.

The Moenave Formation forms distinctive orange-red cliffs along the southwest edge of the Moenkopi Plateau and west of Oraibi Wash on Garces Mesas (figs. 1, 5). The Moenave Formation is not known to yield economic quantities of water in the Black Mesa area.

Wingate Sandstone

It is uncertain if the Wingate Sandstone is present in the Black Mesa area. Billingsley and others (2012, 2013) considered the Wingate Sandstone to be absent from the Moenkopi Plateau and the Hopi Buttes area and concluded that sandstones in these areas formerly mapped as Wingate are in fact part of the Moenave Formation. The Wingate Sandstone may be present deep in the subsurface of the northeastern part of the Black Mesa area, but there is no corroborating information to verify this. Historically the Wingate Sandstone was divided into two members. The upper unit

was the Lukachukai Member, which consisted mostly of eolian, large-scale crossbedded sandstone, while the lower Rock Point Member mainly consisted of flat-bedded fluvial and lacustrine sediments (McKee and MacLachlan, 1959). More recently, the Rock Point Member has been assigned to the underlying Chinle Formation and the Lukachukai Member has been dropped, leaving the name Wingate Sandstone (Dubiel, 1989). At its type locality near Fort Wingate, New Mexico, Harshbarger and others (1957) described the Wingate Sandstone as “pale-reddish-brown fine- to very fine-grained quartz sandstone.” Harshbarger and others (1966) considered the eolian facies of the Wingate Sandstone to be a water-bearing unit of the N aquifer where present.

Kayenta Formation

According to Blakey and Ranney (2008), the Wingate sand dunes were eventually “overwhelmed everywhere by a sandy, braided fluvial system preserved as the Kayenta Formation” (fig. 2). Imlay (1980) reported that the Kayenta Formation consists of light-gray to reddish-orange sandstone and siltstone. The sandstone layers in the Kayenta Formation tend to form ledges whereas the siltstone layers form slopes. Wilson (1965) described the thickness of the Kayenta Formation in south-central Utah as increasing progressively from east to west in part owing to intertonguing with the overlying Navajo Sandstone. Intertonguing of the Kayenta Formation and Navajo Sandstone can be seen clearly in outcrops of the two units along Moenkopi Wash near



Figure 5. Aerial photograph of Moenave Formation outcropping on Garces Mesas, northeastern Arizona. White caprock on top of the Moenave Formation is silicified sandstone of the Kayenta Formation. Photograph by Jon Mason, U.S. Geological Survey.

Tuba City. The Kayenta Formation does not yield economic quantities of water in the Black Mesa area and is therefore not considered an aquifer.

Navajo Sandstone

The Navajo Sandstone is the principal water-bearing unit of the N aquifer (fig. 2). According to Harshbarger and others (1957) it is an eolian deposit made up of sediments derived in part from fluvial deposits of the underlying Kayenta Formation. Beitler and others (2005) described the Navajo Sandstone as a “subrounded, fine- to medium grained, well-sorted, quartz arenite to subarkose sandstone.” The type and amount of cement in the sandstone varies considerably and includes quartz, calcite, dolomite, kaolinite, goethite, and hematite. It is characterized by high-angle, large scale cross-stratification and striking red to white color variations. The red pigment in Navajo Sandstone comes from thin hematite grain coatings. When these coatings are chemically reduced by hydrocarbons migrating through pore spaces, the sandstone is bleached to a lighter color (Beitler and others, 2003). Bedding features in the Navajo Sandstone are identical to those in modern dunes of the transverse and barchan types. In the Black Mesa area, the Navajo Sandstone contains many lenticular beds of cherty limestone deposited in interdune lakes that can be seen between Tuba City and the Hopi Buttes (Harshbarger and others, 1957).

The thickness of the Navajo Sandstone was reported by Harshbarger and others (1957) as 950 ft near Shonto, 478 ft near Dennehotso, 335 ft at Rock Point, and 15 ft northwest of Chinle. Well log data indicate that the top of the Navajo Sandstone is about 2,500 ft below the Black Mesa Complex and has a thickness in the mine area of around 700 ft. In the Tuba City area, where the Navajo Sandstone and Kayenta Formation are intertongued, well log data suggest the combined thickness of the intertongued portion to be greater than 500 ft. Interpretation of the well log from Black Mesa observation well 3 (BM 3) located in Kayenta indicate the top of the Navajo Sandstone is about 155 ft below land surface and that the unit is about 700 ft thick. In the Keams Canyon area, well logs indicate the top of the Navajo Sandstone is about 900 ft below land surface and has a thickness of around 150 ft. Well logs from Kykotsmobi Village indicate the top of the Navajo Sandstone is around 850 ft below land surface with a thickness of over 200 ft.

Carmel Formation

The Carmel Formation (fig. 2) is part of the San Rafael Group. Harshbarger and others (1957) reported the formation in northeastern Arizona as Middle and Late Jurassic in age and consisting of resistant ledge-forming sandstone beds 1 to 3 ft thick separated by slope-forming siltstone strata 5 to 20 ft thick. They further described the siltstone beds as weakly cemented grayish red, weathering to pale reddish brown in color, and the sandstone beds as light greenish gray, weathering to pale yellow (Harshbarger and others, 1957). In most places in northeastern Arizona the Carmel Formation is 100 to 200 ft thick but is thinner at the limits of its deposition (Harshbarger and others, 1957).

According to Blakey and others (1983), the Carmel Formation was deposited in two major transgressive-regressive cycles of the Jurassic Western Interior Seaway, resulting in varied depositional facies including fluvial, eolian, coastal sabkha, and marine. Where present in the Black Mesa area, the Carmel Formation overlies the Navajo Sandstone, forming a confining layer when the Navajo is fully saturated. In most of the study area where the N aquifer is unconfined, the Carmel Formation is absent.

The Carmel Formation also hydraulically separates the N aquifer from the overlying D aquifer in areas where both aquifers are present. In the southern part of Black Mesa there may be some leakage from the D aquifer through the Carmel Formation into the N aquifer (Truini and Macy, 2005). Because the D aquifer has higher total-dissolved solids concentrations than the N aquifer, leakage between the two could degrade the water quality of the N aquifer.

Geologic Units of the D Aquifer

Entrada Sandstone

The Entrada Sandstone (fig. 2) is part of the San Rafael Group and was deposited during the Middle Jurassic in widespread eolian sand seas that were adjacent and inland from a restricted marine seaway (Blakey, 2008; Peterson, 1988). Harshbarger and others (1957) described two general facies of the Entrada Sandstone in the Black Mesa area. The first is a red silty spheroidally weathered sandstone which often weathers into hoodoos. The second is a clean, sandy facies which weathers into rounded massive cliffs. Where resistant cap rocks are present, the Entrada Sandstone weathers into prominent cliffs. Billingsley and others (2012) described the sediments in the Entrada Sandstone as crossbedded, white and interbedded white and red in color.

Harshbarger and others (1951) named a unit overlying the Entrada Sandstone near the Navajo community of Cow Springs, as the Cow Springs Sandstone. Peterson (1988) reports that the Cow Springs Sandstone is closely related to the Entrada Sandstone and often difficult to differentiate from it, but also that the Cow Springs Sandstone can serve as a useful stratigraphic marker. For this reason, Peterson (1988) reduced the rank of the Cow Springs Sandstone to a member of the Entrada Sandstone. The Entrada Sandstone is a water-bearing unit of the D aquifer in the Black Mesa area.

Morrison Formation

The Morrison Formation (fig. 2) was deposited by streams draining uplands in Nevada and central Arizona during the Late Jurassic (Blakey and Ranney, 2008). Harshbarger and others (1957) described the Morrison Formation as primarily fluvial, consisting of alternating flood-plain and channel deposits. There are several recognized members within the Morrison Formation, but only a general description for the formation will be presented here. The Morrison Formation is commonly very colorful. Cooley and others (1969) reported formation colors

include white, gray, green, red, orange, purple, tan, yellow, and brown. They reported the lithology as having mudstone, siltstone, sandstone, conglomerate, and limestone (Cooley and others, 1969). The extent of the Morrison Formation is not fully known in the Black Mesa area. On the west side of Black Mesa there are areas such as Coal Mine Canyon and Blue Canyon where the adjacent units of Entrada Sandstone and Dakota Sandstone (fig. 2) crop out, but the Morrison Formation is missing. Cooley and others (1969) show the Morrison Formation present in a band along and to the north and northeast of Black Mesa. Where present, sandstone beds in the Morrison Formation can be a water-bearing part of the D aquifer in the Black Mesa area (Cooley and others, 1969).

Dakota Sandstone

According to Aubrey (1992), the Dakota Sandstone (fig. 2) represents a complex variety of continental, marginal-marine, and marine environments, and was deposited during the Late Cretaceous in response to the westward transgression of the Cretaceous Interior Seaway. Blakey and Ranney (2008) described the Dakota Sandstone as being made up of beach and coastal plain deposits. The lithology of the unit is described as “medium-to light-gray, slope-forming, laminated to thin-bedded mudstone, siltstone, and sandstone” by Billingsley and others (2012). Cooley and others (1969) reported that the Dakota Sandstone was the chief unit of the D aquifer system.

Mancos Shale

Kirkland (1991) reported that exposures of Mancos Shale (fig. 2) around Black Mesa represent an open marine environment of the Cretaceous Interior Seaway. According to Blakey and Ranney (2008), the Mancos Shale is drab gray and can form odd, moonlike badlands. A good example of badlands weathering of the Mancos Shale can be seen in Blue Canyon along Moenkopi Wash on the Hopi Reservation. Presumably the canyon takes its name from the blueish-gray hue of the Mancos Shale in this location. The Mancos Shale is a thick aquiclude that separates groundwater in the underlying Dakota Sandstone from that in the overlying sandstone aquifers of the Mesaverde Group (Cooley and others, 1969).

Geologic Units of the T Aquifer

Mesaverde Group

According to Franczyk (1988), units of the Mesaverde Group (fig. 2) in the Black Mesa area (Toreva Formation, Wepo Formation, and Yale Point Sandstone) were deposited during the Late Cretaceous by further transgressions and regressions of the Cretaceous Interior Seaway. The Toreva Formation is likely a fluvial and deltaic deposit laid down as the Cretaceous sea regressed after depositing the Mancos Shale (Franczyk, 1988). The formation has multiple members that represent the different depositional environments associated with coastal deposition.

The lithology of the Toreva Formation is varied. Franczyk (1988) reported the formation includes sandstone, siltstone, mudstone, and shale, with some carbonaceous beds.

Page and Repenning (1958) reported the Wepo Formation is of mostly continental origin and consists of a thick series of intercalated siltstone, mudstone, sandstone, and coal. According to Franczyk (1988), the Wepo Formation was deposited while the Cretaceous Interior Seaway was located to the northeast of Black Mesa. Coal beds mined at the Black Mesa Complex occur in the Wepo Formation.

Molenaar (1983) described the Yale Point Sandstone as a coastal-barrier sandstone deposited during one of the last transgressions of the Cretaceous Interior Seaway. According to O’Sullivan and others (1972), the Yale Point Sandstone is “yellowish gray, weathers grayish orange, and is composed of coarse- to fine-grained subrounded to subangular clear quartz.” Bedding in the formation is lenticular, and individual units are crossbedded (O’Sullivan and others, 1972).

Sandstone units in the Mesaverde Group can be water-bearing units of the T aquifer. Many small contact springs issue from Mesaverde sandstones around the perimeter of Black Mesa and in canyons where the sandstones have been truncated.

Bidahochi Formation

According to a distribution map of the Bidahochi Formation by Repenning and Irwin (1954), the only place the formation is present in the Black Mesa area is around the Hopi Buttes. They described it as consisting of fluvial and lacustrine deposits and basaltic volcanic rock (Repenning and Irwin, 1954). According to Blakey and Ranney (2008), the depositional environment of the Bidahochi Formation is still unresolved; they suggested the formation could have been deposited in a Neogene lake, but the evidence for this deposition is unclear. Harshbarger and others (1966) reported that the lower part of the Bidahochi Formation can be water bearing.

Hydrologic Data

Groundwater data collected for this report were exclusively from the N aquifer. Water from the T and D aquifers are not used in significant quantities in the Black Mesa area. Water from the T aquifer is used locally for livestock watering and to irrigate small plots of land, but it probably cannot produce enough water for municipal or industrial use. Water from the D aquifer is used locally for livestock watering and in the past contributed to some wells at the PWCC industrial well field, but water from the aquifer generally has total-dissolved solids concentrations that make it unsuitable for municipal use.

In 2018–19, activities of the Black Mesa area monitoring program included metered groundwater withdrawals, measurements of groundwater levels, spring-discharge measurements, streamflow gaging, and the collection of

water-chemistry samples from wells and springs. All data were collected by the USGS except withdrawal data from NTUA wells, which were compiled by NTUA personnel. Annual discharge measurements were made at 4 springs, and annual groundwater-level measurements were made at 34 wells. Of the 34 wells, 6 are continuous-recording observation wells that have been outfitted for real-time data telemetry (denoted to as “BM observation well” in table 3 and “BM” in text). The water-level data from these six continuous-recording observation wells are available on the NWIS website (<http://waterdata.usgs.gov/az/nwis/gw>).

Groundwater-withdrawal data were compiled during February 2020. Spring discharges and groundwater levels were measured between February and April 2019. Groundwater samples were collected from 4 springs and 3 wells from April to August 2019 and were analyzed for chemical constituents. Annual groundwater-withdrawal data are collected from 36 well systems within the BIA, NTUA, and Hopi Reservation municipal systems, as well as the PWCC industrial well field. Water-level measurements are attempted from 34 wells and well identification information is shown in table 3. Streamflow data are collected at four USGS gaging stations and are available online (<http://waterdata.usgs.gov/az/nwis/rt>). All annual data reported in this document are for calendar years beginning January 1 and ending December 31. Median winter streamflow is reported as the year in which the winter season began, which for this report is 2018. The period before appreciable groundwater withdrawals began for mining or municipal purposes (about 1965) is referred to in this report as the prestress period.

Kendall's tau trend analyses were applied to streamflow data, spring-discharge measurements, and water-chemistry samples by using R Project for Statistical Computing (R Development Core Team, 2020). The Kendall's tau correlation coefficient was computed between the measured data and time. The null hypothesis was that no correlation existed between time and the measured data; the alternate hypothesis was that time and the measured data were correlated. A significance level of 0.05 was chosen to determine whether the result of the test for significance of Kendall's tau correlation coefficient was statistically significant. A two-sided p-value that was less than or equal to 0.05 indicated that there was a statistically significant correlation in the data, and the null hypothesis was rejected. The sign of the slope indicated whether there was an increasing or decreasing trend if a significant correlation existed. P-values greater than 0.05 indicated that there was no statistically significant correlation between time and concentration, and the null hypothesis was accepted.

In addition, the Theil-Sen slope estimator was calculated and plotted for streamflow data, spring-discharge measurements, and water-chemistry data using R package “NADA” (Lee, 2020). Closely related to Kendall's tau, the Theil-Sen slope estimator provides a slope for trends similar to ordinary least-squares regression, but is less affected by outliers (Helsel and others, 2020).

Table 3. Identification numbers and names of monitoring-program wells used for water-level measurements or for collection of water-chemistry samples, Black Mesa area, northeastern Arizona, 2018–19.

[---, no data]

U.S. Geological Survey identification number	Common name or location	Bureau of Indian Affairs site number
355023110182701	Keams Canyon PM2	---
355230110365801	Kykotsmovi PM1	---
355236110364501	Kykotsmovi PM3	---
355428111084601	Goldtooth	3A-28
355924110485001	Howell Mesa	3K-311
360055110304001	BM observation well 5 ^a	4T-519
360217111122601	Tuba City	3K-325
360614110130801	Piñon PM6	---
360734111144801	Tuba City	3T-333
360904111140501	Tuba City NTUA 1R ^b	---
360918111080701	Tuba City Rare Metals 2	---
360927111142401	Tuba City NTUA 3R ^c	---
360953111142401	Tuba City NTUA 4	3T-546
361225110240701	BM observation well 6 ^a	---
361737110180301	Forest Lake NTUA 1	4T-523
361832109462701	Rough Rock	10T-258
362043110030501	Kits'ili NTUA 2	---
362149109463301	Rough Rock	10R-111
362333110250001	Peabody 9	---
362406110563201	White Mesa Arch	1K-214
362456110242301	Peabody 7	---
362625110223701	Peabody 3	---
362823109463101	Rough Rock	10R-119
362936109564101	BM observation well 1 ^a	8T-537
363013109584901	Sweetwater Mesa	8K-443
363103109445201	Rough Rock	9Y-95
363143110355001	BM observation well 4 ^a	2T-514
363213110342001	Shonto Southeast	2K-301
363232109465601	Rough Rock	9Y-92
363309110420501	Shonto	2K-300
363423110305501	Shonto Southeast	2T-502
363727110274501	Long House Valley	8T-510
363850110100801	BM observation well 2 ^a	8T-538
364034110240001	Marsh Pass	8T-522
364226110171701	Kayenta West	8T-541
364248109514601	Northeast Rough Rock	8A-180
364338110154601	BM observation well 3 ^a	8T-500

^aWell with continuous water-level recorder.

^bWell replaced Tuba City NTUA 1 (360904111140201) in 2018.

^cWell replaced Tuba City NTUA 3 (360924111142201) in 2018.

Withdrawals from the N Aquifer

Total annual withdrawals from the N aquifer are monitored on a continuing basis to help determine the effects from industrial and municipal pumping. Withdrawals from the N aquifer are separated into three categories: (1) industrial withdrawals from the confined area, (2) municipal withdrawals from the confined area, and (3) municipal withdrawals from the unconfined areas. Within the study area there are no industrial withdrawals from the unconfined areas. The industrial category includes eight wells in the PWCC industrial well field in the northern Black Mesa area. The BIA, NTUA, and Hopi Tribe operate about 70 municipal wells that are combined into 36 well systems. Information about withdrawals from the N aquifer is compiled primarily on the basis of metered data from individual wells operated by the BIA, NTUA, and Hopi Tribe (table 4). Meter readings from nine well systems operated by the NTUA were not available when data for this report was compiled. NTUA offices closed in the spring of 2020 owing to the COVID-19 pandemic before all the 2019 meter readings were collected. In the case of well systems with no available 2019 meter readings, the average annual pumpage over the last four years from each of the systems was used as an estimate for 2019 pumpage.

Withdrawals from wells equipped with windmills are not measured in this monitoring program and are not included in total withdrawal values reported here. About 270 windmills in the Black Mesa area withdraw water from the N, D, T, and alluvial aquifers, primarily for livestock. The estimated total withdrawal by the windmills from the N aquifer is about 65 acre-feet per year (acre-ft/yr) (HSIGeoTrans, Inc. and Waterstone Environmental Hydrology and Engineering, Inc., 1999). The total withdrawal by the windmills is less than 1 percent of the total annual withdrawal from the N aquifer.

Withdrawals in Calendar Year 2019 Compared to Previous Years

In 2019 total groundwater withdrawal from the N aquifer was estimated to be about 3,070 acre-ft (table 1). Total withdrawals for municipal use from 2019 was estimated to be about 2,400 acre-ft; municipal withdrawals from the confined area totaled about 1,340 acre-ft, while withdrawals from the unconfined areas totaled about 1,060 acre-ft. Withdrawals for industrial use totaled about 670 acre-ft (tables 1, 5).

Withdrawals from the N aquifer have varied annually from 1965 to the present but generally increased from 1965 to 2005 and decreased from 2006 to 2019. Beginning in 2006, the PWCC reduced their pumping by about 70 percent, a reduction that is reflected by a decrease in total annual withdrawals from 2005 by about 44 percent (tables 1, 5; fig. 6). Total withdrawals for the period of record (1965–2019) was 273,740 acre-ft; industrial withdrawals were 57 percent and municipal withdrawals were 43 percent of total withdrawals (table 5). Total withdrawals in 2019 were 3,070 acre-ft, with 22 percent from industrial withdrawals and 78 percent from municipal withdrawals (table 5). Industrial

pumping decreased about 43 percent between 2018 and 2019. As discussed earlier, the PWCC stopped producing coal from the Black Mesa Complex in August 2019 due to the planned closure of the Navajo Generating Station. Groundwater continues to be used at the Black Mesa Complex for the reclamation process, but at a much-diminished rate.

Groundwater Levels in the N Aquifer

Groundwater levels are monitored at wells that are screened in the N aquifer to help understand the effects of withdrawals on the potentiometric surface of the aquifer. Groundwater in the N aquifer is under confined conditions in the central part of the study area and under unconfined or water-table conditions around the periphery (fig. 7). Because of the different ways groundwater is released from storage, confined and unconfined aquifers respond dissimilarly to groundwater withdrawal. When the same volume of water is withdrawn from both confined and unconfined aquifers by pumping, the water level decline in wells within the confined aquifer will usually be much greater than the decline seen in wells within the unconfined aquifer. This is one reason water levels have generally declined more in wells screened in the confined portion of the N aquifer versus the unconfined portion within the monitoring area. A corollary to this phenomenon is that water levels in confined aquifer wells also tend to recover quicker if pumping is decreased or stopped than they would in unconfined aquifer wells.

Direct comparison between water levels from confined aquifer wells and unconfined aquifer wells is of limited value. The two sets of data are distinct populations that are probably best considered individually. For this reason, a distinction is made between water level changes from confined aquifer wells and unconfined aquifer wells in this section of the report.

Between February and April 2019, water-levels were measured in 34 wells and compared to prestress levels (table 6) to identify long-term changes. Of the 34 wells, 6 are continuous-recording observation wells. Water levels were measured quarterly using an electric tape in these six wells during water year 2019 to verify or update instrument calibration. Only water levels from municipal and stock wells that were not considered to have been recently pumped, affected by nearby pumping, or blocked or obstructed are compared.

The wells used for water-level measurements are distributed throughout the study area (fig. 8). The wells were constructed between 1934 and 2018 and the well depths range from 107 ft near Rough Rock (8A-180) to 2,674 ft at Forest Lake (Forest Lake NTUA 1). Depths to the top of the N aquifer range from 0 ft near Tuba City to 2,205 ft at Kits'iilie on top of Black Mesa (table 7). The Tuba City NTUA 1R and 3R wells were drilled in 2018 to replace wells Tuba City NTUA 1 and 3. The new wells were drilled close to the old wells but are screened at slightly different intervals. The original wells have been plugged and are no longer available for water-level measurements. Water levels from the new Tuba City NTUA 1R and 3R wells are compared to the prestress water levels from the original Tuba City NTUA 1 and 3 wells in table 6.

Table 4. Withdrawals from the N aquifer by well system, Black Mesa area, northeastern Arizona, calendar year 2019.

[Withdrawals are from flowmeter measurements. BIA, Bureau of Indian Affairs; NTUA, Navajo Tribal Utility Authority; USGS, U.S. Geological Survey; PWCC, Peabody Western Coal Company; Hopi, Hopi Village Administrations]


Well system (one or more wells)	Owner	Source of data	2019 withdrawals		Well system (one or more wells)	Owner	Source of data	2019 withdrawals	
			Confined aquifer	Unconfined aquifer				Confined aquifer	Unconfined aquifer
			(acre-feet)					(acre-feet)	
Chilchinbito	BIA	USGS/BIA	4.1		Kayenta	NTUA	USGS/NTUA	^b 360.1	
Dennehotso	BIA	USGS/BIA		4.2	Kits'iili	NTUA	USGS/NTUA	^b 16.6	
Hopi High School	BIA	USGS/BIA	10.1		Piñon	NTUA	USGS/NTUA	^b 359.7	
Hotevilla	BIA	USGS/BIA	23.4		Red Lake	NTUA	USGS/NTUA		39.0
Kayenta	BIA	USGS/BIA	20.3		Rough Rock	NTUA	USGS/NTUA	46.2	
Keams Canyon	BIA	USGS/BIA	58.3		Shonto	NTUA	USGS/NTUA		^b 24.1
Low Mountain	BIA	USGS/BIA	^a 0		Shonto Junction	NTUA	USGS/NTUA		^b 75.4
Piñon	BIA	USGS/BIA	^a 0		Tuba City	NTUA	USGS/NTUA		663.1
Red Lake	BIA	USGS/BIA		3.0	Mine Well Field	PWCC	PWCC	672.8	
Rocky Ridge	BIA	USGS/BIA	3.2		Bacavi	Hopi	USGS/Hopi	15.7	
Rough Rock	BIA	USGS/BIA	9.8		Hopi Civic Center	Hopi	USGS/Hopi	1.1	
Second Mesa	BIA	USGS/BIA	3.4		Hopi Cultural Center	Hopi	USGS/Hopi	4.9	
Shonto	BIA	USGS/BIA		76.6	Kykotsmovi	Hopi	USGS/Hopi	63.1	
Tuba City	BIA	USGS/BIA		66.4	Mishongnovi	Hopi	USGS/Hopi	5.7	
Chilchinbito	NTUA	USGS/NTUA	^b 62.7		Moenkopi	Hopi	USGS/Hopi		65.8
Dennehotso	NTUA	USGS/NTUA		^b 39.8	Polacca	Hopi	USGS/Hopi	150.3	
Forest Lake	NTUA	USGS/NTUA	^b 13.4		Shipaulovi	Hopi	USGS/Hopi	19.0	
Hard Rock	NTUA	USGS/NTUA	44.7		Shungopovi	Hopi	USGS/Hopi	30.2	

^aWell taken out of service.

^bEstimated value owing to unavailability of 2019 record.

Table 5. Total, industrial, and municipal withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, for discrete time periods from 1965 to 2019.

Period	Total withdrawals	Industrial withdrawals	Municipal withdrawals	Percent industrial	Percent municipal
	(acre-feet)				
1965–2019	273,740	155,420	118,320	57	43
1965–2005	218,300	138,100	80,200	63	37
2006–2019	55,440	17,320	38,120	31	69
2019	3,070	670	2,400	22	78

**Figure 6.** Plot of annual withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, 1965–2019.




Figure 7. Map of well systems monitored for annual withdrawals from the N aquifer, Black Mesa area, northeastern Arizona, calendar year 2019, and confined/unconfined zones of the N aquifer.

Table 6. Water-level changes in monitoring program wells completed in the N aquifer, Black Mesa area, northeastern Arizona, prestress period (prior to 1965) to calendar year 2019.

[ft, feet; ---, no data]

Common name or location	Bureau of Indian Affairs site number	Water level (ft below land surface), 2019	Prestress period water level		Change in water level from prestress period to 2019 (ft)
			(ft below land surface)	(date)	
Unconfined areas					
BM observation well 1 ^a	8T-537	374.2	374.0	(^a)	−0.2
BM observation well 4 ^a	2T-514	216.7	216.0	(^a)	−0.7
Goldtooth	3A-28	232.7	230.0	10–29–53	−2.7
Long House Valley	8T-510	139.4	99.4	08–22–67	−40.0
Northeast Rough Rock	8A-180	44.5	46.9	11–13–53	2.4
Rough Rock	9Y-95	111.4	119.5	08–03–49	8.1
Rough Rock	9Y-92	166.5	168.8	12–13–52	2.3
Shonto	2K-300	172.4	176.5	06–13–50	4.1
Shonto Southeast	2K-301	291.1	283.9	12–10–52	−7.2
Shonto Southeast	2T-502	418.3	405.8	08–22–67	−12.5
Tuba City	3T-333	28.2	23.0	12–02–55	−5.2
Tuba City	3K-325	205.1	208.0	06–30–55	2.9
Tuba City Rare Metals 2	---	48.8	57.0	09–24–55	8.2
Tuba City NTUA 1R	---	53.2	^b 29.0	02–12–69	−24.2
Tuba City NTUA 3R	---	57.2	^b 34.2	11–08–71	−23.0
Tuba City NTUA 4	3T-546	57.6	33.7	08–06–71	−23.9
Confined area					
BM observation well 2 ^a	8T-538	207.0	125.0	(^a)	−82.0
BM observation well 3 ^a	8T-500	161.3	55.0	04–29–63	−106.3
BM observation well 5 ^a	4T-519	420.2	324.0	(^a)	−96.2
BM observation well 6 ^a	---	833.3	697.0	(^a)	−136.3
Forest Lake NTUA 1	4T-523	1,155.6	^c 1,096	05–21–82	−59.6
Howell Mesa	3K-311	450.2	463.0	11–03–53	12.8
Kayenta West	8T-541	284.0	230.0	03–17–76	−54.0
Keams Canyon PM2	---	477.5	292.5	06–10–70	−185.0
Kits’iili NTUA 2	---	1,336.8	^d 1,297.9	01–14–99	−38.9
Kykotsmovi PM1	---	207.1	220.0	05–20–67	12.9
Kykotsmovi PM3	---	248.8	210.0	08–28–68	−38.8
Marsh Pass	8T-522	131.9	125.5	02–07–72	−6.4
Piñon PM6	---	907.8	743.6	05–28–70	−164.2
Rough Rock	10R-119	260.9	256.6	12–02–53	−4.3
Rough Rock	10T-258	309.9	301.0	04–14–60	−8.9
Rough Rock	10R-111	191.8	170.0	08–04–54	−21.8
Sweetwater Mesa	8K-443	548.1	529.4	09–26–67	−18.7
White Mesa Arch	1K-214	218.8	188.0	06–04–53	−30.8

^aContinuous recorder. Prestress water levels estimated from a groundwater model, except at BM observation well 3 (Brown and Eychaner, 1988).^bWater level is from a previous well drilled to a shallower depth at the same location.^cReported from driller's log.^dWater level is the first water level measured after completion of well.




Figure 8. Map showing water-level changes in wells completed in the N aquifer, Black Mesa area, northeastern Arizona, prestress period (prior to 1965) to 2019.

Table 7. Well-construction characteristics, depth to top of N aquifer, and 2019 static water level for wells used in annual water-level measurements and for continuous-record observation wells, Black Mesa area, northeastern Arizona, 2018–19.

[ft, feet]

Bureau of Indian Affairs site number and (or) common name	Date well was completed	Land-surface elevation (ft)	Well depth (ft) below land surface)	Screened/open interval(s) (ft below land surface)	Depth to top of N aquifer (ft below land surface) ^a	2019 static water level (ft below land surface)
8T-537 (BM observation well 1)	02-01-1972	5,864	851	300–360; 400–420; 500–520; 600–620; 730–780	290	374.2
8T-538 (BM observation well 2)	01-29-1972	5,656	1,338	470–1,338	452	207.0
8T-500 (BM observation well 3)	07-29-1959	5,724	868	712–868	155	161.3
2T-514 (BM observation well 4)	02-15-1972	6,320	400	250–400	160	216.7
4T-519 (BM observation well 5)	02-25-1972	5,869	1,683	1,521–1,683	1,520	420.2
BM observation well 6	01-31-1977	6,332	2,507	1,954–2,506	1,950	833.3
1K-214	05-26-1950	5,771	356	168–356	250	218.8
2K-300	06-00-1950 ^b	6,264	300	260–300	0	172.4
2K-301	06-12-1950	6,435	500	318–328; 378–500	≈30	291.1
2T-502	08-10-1959	6,670	523	12–523	25	418.3
3A-28	04-19-1935	5,381	358	(^d)	60	232.7
3K-311	11-00-1934 ^b	5,855	745	380–395; 605–745	615	450.2
3K-325	06-01-1955	5,250	450	75–450	≈30	205.1
3T-333	12-02-1955	4,940	229	63–229	24	28.2
3T-546 (Tuba City NTUA 4)	08-00-1971 ^b	5,206	612	256–556	0	57.6
4T-523 (Forest Lake NTUA 1)	10-01-1980	6,654	2,674	1,870–1,910; 2,070–2,210; 2,250–2,674	(^e)	1,155.6
8A-180	01-20-1939	5,200	107	60–107	≈40	44.5
8K-443	08-15-1957	6,024	720	619–720	590	548.1
8T-510	02-11-1963	6,262	314	130–314	≈125	139.4
8T-522	07-00-1963 ^b	6,040	933	180–933	480	131.9
8T-541	03-17-1976	5,885	890	740–890	700	284.0
9Y-92	01-02-1039	5,615	300	154–300	≈50	166.5
9Y-95	11-05-1937	5,633	300	145–300	≈68	111.4
10R-111	04-11-1935	5,757	360	267–360	210	191.8
10R-119	01-09-1935	5,775	360	(^d)	310	260.9
10T-258	04-12-1960	5,903	670	465–670	460	309.9
Keams Canyon PM2	05-00-1070 ^b	5,809	1,106	906–1,106	900	477.5
Kits'iili NTUA 2	10-30-1993	6,780	2,549	2,217–2,223; 2,240–2,256; 2,314–2,324; 2,344–2,394; 2,472–2,527	2,205	1,336.8
Kykotsmovi PM1	02-20-1967	5,657	995	655–675; 890–990	880	207.1
Kykotsmovi PM3	08-07-1968	5,618	1,220	850–1,220	840	248.8

Table 7.—Continued

Bureau of Indian Affairs site number and (or) common name	Date well was completed	Land-surface elevation (ft)	Well depth (ft) below land surface)	Screened/open interval(s) (ft below land surface)	Depth to top of N aquifer (ft below land surface) ^a	2019 static water level (ft below land surface)
Piñon PM6	02-00-1970 ^b	6,397	2,248	1,895-2,243	1,870	907.8
Tuba City NTUA 1R ^f	11-02-2018	5,123	650	95-450	0	53.2
Tuba City NTUA 3R ^g	11-02-2018	5,184	650	164-520	34	57.2
Tuba City Rare Metals 2	09-00-1955 ^b	5,108	705	100-705	255	48.8

^aDepth to top of N aquifer from Eychaner (1983) and Brown and Eychaner (1988).
^b00, indicates day is unknown.
^cAll material between land surface and top of the N aquifer is unconsolidated—soil, alluvium, or dune sand.
^dScreened and (or) open intervals are unknown.
^eDepth to top of N aquifer was not estimated.
^fWell is a replacement for Tuba City NTUA 1.
^gWell is a replacement for Tuba City NTUA 3.

Table 8. Median changes in water levels in monitoring-program wells of the N aquifer, Black Mesa area, northeastern Arizona, prestress period (prior to 1965) to 2019.

Years	Aquifer conditions	Number of wells	Median change in water level (feet)
Prestress–2019	Unconfined	16	–1.7
	Confined	18	–38.8

Water levels measured in 2019, and changes in water levels from the prestress period to 2019, are shown in table 6. The 34 wells in table 6 are grouped by location in the unconfined or confined area of the aquifer. From the prestress period to 2019, water levels in 16 wells in the unconfined part of the aquifer had a median change of –1.7 ft (table 8), and water-level changes ranged from –40.0 ft at Long House Valley (8T-510) to +8.2 ft at Tuba City Rare Metals (fig. 8; table 6). Water levels in 18 wells in the confined part of the aquifer had a median change of –38.8 ft (table 8), and water-level changes ranged from –185.0 ft at Keams Canyon PM2 to +12.9 ft at Kytotsmovi PM1 (fig. 8; table 6).

Hydrographs of groundwater levels in the network of wells observed annually show the temporal changes from the 1950s to present (fig. 9). In most of the unconfined areas, water levels have changed only slightly (generally less than 10 ft). Near Shonto, however, the water level in well 8T-510 (Longhouse Valley) has declined 40.0 ft (fig. 8; table 6). Water levels have declined in most of the confined area, but the magnitudes of declines are varied. Larger declines have occurred near the municipal pumping centers

(wells Piñon PM6 and Keams Canyon PM2) and near the well field for the PWCC (BM6). Smaller declines occurred away from the pumping centers in or near towns in the study area (wells 10T-258, 10R-119, 8T-522; figs. 8, 9).

Hydrographs for the Black Mesa continuous-record observation wells (fig. 10) show water levels since the early 1970s. The two wells in the unconfined areas (BM1 and BM4) have shown small seasonal or year-to-year variation since 1972 but no apparent long-term decline. In the confined area, water levels (not corrected for barometric pressure effects or seasonal effects) in wells BM2, BM3, BM5, and BM6 consistently declined from the 1970s to the mid-2000s (fig. 10). After the mid-2000s, water levels in BM2, BM5, and BM6 began to level off and then to rise. The water-level recoveries in BM2, BM5, and BM6 since the mid-2000s have been 11.5 ft, 7.9 ft, and 28.3 ft, respectively. Water levels in BM3 are more variable because of nearby municipal pumping. After the mid-2000s, water levels in BM3 continued to vary, but the overall trend flattened out (fig. 10).




Figure 9 (pages 23–26). Plots of observed water levels in annual observation well network of the N aquifer, Black Mesa area, northeastern Arizona, 1950–2019.




Figure 9 (pages 23–26).—Continued



Figure 9 (pages 23–26).—Continued




Figure 9 (pages 23–26).—Continued

Spring Discharge from the N Aquifer

Groundwater in the N aquifer discharges from many springs around the margins of Black Mesa, and changes to the discharge from those springs could indicate effects of withdrawals from the N aquifer. Moenkopi School Spring (360632111131101), Burro Spring (354156110413701), Pasture Canyon Spring (36102111115901), and Unnamed Spring near Dennehotso (364656109425400) have been measured intermittently since the late 1980s and all four springs were measured for discharge in 2019. Additionally, trend analyses were performed on the flow data from the four springs.

Moenkopi School Spring, also called Susunova Spring by the Hopi Tribe, is in the western part of the Black Mesa area (fig. 11). Discharge from Moenkopi School Spring was measured in April 2019 using the volumetric method and compared to discharge data from previous years to determine changes over time (fig. 12A). The trend for discharge measurements at this spring is not corrected for seasonal variability, but discharge measurements are generally made annually at or close to the same time of year. In 2019, the measured discharge from Moenkopi School Spring was 7.5 gallons per minute (gal/min) (table 9). A Kendall's tau analysis, using the Theil-Sen slope estimator, indicated a decreasing trend ($p < 0.05$) of about 0.3 gal/min per year during the period of record (fig. 12A).

Burro Spring is in the southwestern part of the study area and discharges from the Navajo Sandstone and alluvium (fig. 11). Burro Spring discharges from the aquifer through a metal pipe and into a cement trough for livestock. As in previous years, the 2019 discharge measurement and water-quality sampling point was from the end of the metal pipe

before the livestock trough. Discharge at Burro Spring has fluctuated since 1989 between 0.2 and 0.4 gal/min, but there is no significant ($p > 0.05$) trend from a Kendall's tau analysis (fig. 12B). In 2019, the measured discharge was 0.3 gal/min (table 9).

Pasture Canyon Spring is in the western part of the study area and discharges from the Navajo Sandstone and alluvium (fig. 11). Discharge is measured at two locations: where the spring issues from the Navajo Sandstone (also the water-quality sampling point), and farther down the canyon at the USGS gaging station (Pasture Canyon Springs 09401265). The USGS gaging station at Pasture Canyon measures the discharge from Pasture Canyon Spring as well as additional discharge from seeps in Pasture Canyon. As in previous years, discharge was measured at Pasture Canyon Spring at its emergence point in April 2019 using the volumetric method. The measured discharge was 43.0 gal/min (table 9), indicating an increase in discharge of about 1.0 gal/min from the 2018 measurement. The discharge measured in 2019 is one of the highest discharges recorded at the spring emergence point. Discharge at Pasture Canyon Spring has fluctuated since 1995 between 26.5 and 43.0 gal/min, but there is no significant ($p > 0.05$) trend from a Kendall's tau analysis (fig. 12C). The discharge data measured at this spring is not corrected for seasonal variability, but annual discharge measurements are generally made close to the same time of year.

Unnamed Spring near Dennehotso is the only spring located in the northeastern part of the study area (fig. 11), and it discharges from the Navajo Sandstone. As in previous years, measurements at Unnamed Spring near Dennehotso are made using a flume. Discharge has decreased markedly at Unnamed




Figure 10. Plots of observed groundwater levels in continuous-record observation wells BM1–BM6 in the N aquifer, Black Mesa area, northeastern Arizona, 1963–2019.

Spring near Dennehotso since 2005. That year, the discharge at the spring was 21.5 gal/min. The discharge wasn't measured again until 2010 when it was 9.0 gal/min. In 2019 discharge was 12.0 gal/min after several years of being lower (table 9).

For the period of record, which includes a gap in data from 2005 to 2010, a decreasing trend ($p < 0.05$) is evident from a Kendall's tau analysis (fig. 12D). The discharge data measured at this spring is not corrected for seasonal variability.




Figure 11. Map of surface-water and water-chemistry data-collection sites in the N aquifer, Black Mesa area, northeastern Arizona, 2019.




Figure 12. Plots of discharge from *A*, Moenkopi School Spring (360632111131101); *B*, Burro Spring (354156110413701); *C*, Pasture Canyon Spring (361021111115901); and *D*, Unnamed Spring near Dennehotso (364656109425400), N Aquifer, Black Mesa area, northeastern Arizona, 1987–2019. Moenkopi School Spring data from 1952 and Pasture Canyon Spring data from 1988 to 1993 are not shown because they were taken from different measuring locations.

Table 9. Discharge from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso, N Aquifer, Black Mesa area, northeastern Arizona, 1952–2019.

[Measured discharges do not represent the total discharge from the springs. gal/min, gallons per minute]

Bureau of Indian Affairs site number	Rock formation(s)	Date of measurement	Discharge (gal/min)	Bureau of Indian Affairs site number	Rock formation(s)	Date of measurement	Discharge (gal/min)		
Moenkopi School Spring ^a				Burro Spring ^a —Continued					
3GS-77-6	Navajo Sandstone ^b	05–16–1952	40.0	6M-31—Continued	Navajo Sandstone—Continued	06–07–2010	0.3		
		04–22–1987	^c 16.0			06–08–2011	0.4		
		11–29–1988	^c 12.5			06–14–2012	0.3		
		02–21–1991	^c 13.5			07–30–2013	0.3		
		04–07–1993	^c 14.6			09–02–2014	0.3		
		12–07–1994	^c 12.9			06–23–2016	0.2		
		12–04–1995	^c 10.0			07–18–2017	0.3		
		12–16–1996	^c 13.1			06–06–2018	0.3		
		12–17–1997	^c 12.0			04–22–2019	0.3		
		12–08–1998	^c 13.3			Pasture Canyon Spring ^a			
		12–13–1999	^c 13.7			3A-5	Navajo Sandstone, alluvium	11–18–1988	^c 211
		03–12–2001	^c 10.2					03–24–1992	^c 233
		06–19–2002	^c 11.2					10–12–1993	^c 211
		05–01–2003	^c 11.2					12–04–1995	^f 38.0
		03–29–2004	^c 12.2					12–16–1996	^f 38.0
		04–04–2005	^c 11.5					12–17–1997	^f 40.0
		03–13–2006	^c 11.1					12–10–1998	^f 39.0
		05–31–2007	^c 9.0					12–21–1999	^f 39.0
		06–03–2008	^c 8.3					06–12–2001	^f 37.0
		06–03–2009	^c 8.0					04–04–2002	^f 37.0
		06–14–2010	^c 7.4					05–01–2003	^f 30.9
		06–10–2011	^c 9.0					04–26–2004	^f 30.6
		06–07–2012	^c 6.3					04–27–2005	^f 33.3
		07–29–2013	^c 6.4					06–03–2008	^f 29.4
		08–27–2014	^c 6.3					06–03–2009	^f 31.1
		06–21–2016	^c 6.0					06–14–2010	^f 34.3
		07–11–2017	^c 6.8					06–09–2011	^f 31.4
		06–06–2018	^c 6.4					06–07–2012	^f 26.5
		04–22–2019	^c 7.5					07–29–2013	^f 35.7
Burro Spring ^a				08–27–2014	^f 39.3				
6M-31	Navajo Sandstone	12–15–1989	0.4	06–21–2016	^f 37.5				
		12–13–1990	0.4	07–11–2017	^f 43.0				
		03–18–1993	0.3	06–06–2018	^f 42.0				
		12–08–1994	0.2	04–22–2019	^f 43.0				
		12–17–1996	0.4	Unnamed Spring near Dennehotso ^e					
		12–30–1997	0.2	8A-224	Navajo Sandstone	10–06–1954	^g 1		
		12–08–1998	0.3			06–27–1984	^g 2		
		12–07–1999	0.3			11–17–1987	^g 5		
		04–02–2001	0.2			03–26–1992	16.0		
		04–04–2002	0.4			10–22–1993	14.4		
		04–30–2003	0.4			12–05–1995	17.0		
		04–06–2004	^d 0.2			12–19–1996	15.7		
		03–28–2005	0.2			12–30–1997	25.6		
		03–28–2006	0.2			12–14–1998	21.0		
		06–04–2009	0.3			12–15–1999	14.8		

Table 9.—Continued

Bureau of Indian Affairs site number	Rock formation(s)	Date of measurement	Discharge (gal/min)
Unnamed Spring near Dennehotso ^d —Continued			
8A-224—	Navajo	03-14-2001	26.8
Continued	Sandstone—	04-03-2002	5.8
	Continued	07-15-2002	9.0
		05-01-2003	17.1
		04-01-2004	12.6
		04-06-2005	21.5
		06-17-2010	9.0
		06-04-2012	4.5
		08-06-2013	6.7
		09-03-2014	8.1
		10-26-2016	9.0
		07-03-2018	3.0
		04-23-2019	12.0

^aVolumetric discharge measurement.

^bInterfingering with the Kayenta Formation at this site.

^cDischarge measured at water-quality sampling site and at a different point than the measurement in 1952.

^dDischarge is approximate because the container used for the volumetric measurement was not calibrated.

^eDischarge measured in channel below water-quality sampling point.

^fDischarge measured at water-quality sampling point about 20 feet below upper spring on west side of canyon.

^gDischarge measured at a different point than later measurements.

Surface-Water Discharge, Calendar Year 2019

Continuous surface-water discharge data have been collected at selected streams since the monitoring program began in 1971. Surface-water discharge in the study area generally originates as groundwater that discharges to streams or as surface runoff from rainfall or snowmelt. Groundwater discharges to some stream reaches at a fairly constant rate throughout the year; however, the amount of groundwater discharge that results in surface flow is affected by seasonal fluctuations in evapotranspiration (Thomas, 2002a). In contrast, the amount of rainfall or snowmelt runoff varies widely throughout the year. In the winter and spring, the amount and timing of snowmelt runoff are a result of the temporal variation in factors such as snow accumulation, air temperatures, and rate of snowmelt. However, snowmelt usually does not lead to large runoff events on any of the four washes. Rainfall can occur throughout the year, but more typically occurs during the summer months than during other times of the year. The amount and timing

of rainfall depends on the intensity and duration of thunderstorms during the summer and mid-latitude low pressure systems during the fall, winter, and spring.

In 2019, discharge data were collected at four continuous-recording streamflow-gaging stations (fig. 13). Data collection at these stations began in July 1976 (Moenkopi Wash at Moenkopi, 09401260), June 1993 (Dinnebito Wash near Sand Springs, 09401110), April 1994 (Polacca Wash near Second Mesa, 09400568), and August 2004 (Pasture Canyon Springs, 09401265) (table 10). Most of the daily mean discharge values highlighted as estimated in figure 13 were either estimated because the streamflow record was affected by ice during the winter months or because the stage recorder was damaged during high flow conditions. Estimated daily values are based on adjacent good records, records from comparable stations, and discrete discharge measurements. Areas of hydrographs in figure 13 where no line is present are periods when mean daily discharge was zero or near zero. The hydrographs present mean daily discharge using a logarithmic axis, which cannot have a zero value. The geologic and hydrologic settings along with trend analyses of base flow at the four streamflow sites are described briefly below.

Moenkopi Wash

Moenkopi Wash has a drainage area of 1,629 mi² and drains a large portion of the western part of Black Mesa. The streamflow gage is located near the Village of Moenkopi in a portion of the wash that cuts down into interbedded Navajo Sandstone and Kayenta Formation. During the period of streamflow gage operation, there has generally been continuous flow at the gage except for the summer months, when the stream is often dry at the gage (fig. 13). Monsoon rain events occurring between July and September can result in large sediment laden flows in Moenkopi Wash. The maximum instantaneous discharge recorded at the gage was 10,100 cubic feet per second (ft³/s) on September 30, 1983.

There are no observed N-aquifer springs issuing directly from the Navajo Sandstone near the streamflow gage. During base flow conditions, flow seems to initiate from Moenkopi Wash alluvium, but it is assumed this flow is supplied by the N aquifer from below and through the alluvium.

Dinnebito Wash

Dinnebito Wash has a drainage area of 473 mi² and drains part of the middle portion of Black Mesa. The streamflow gage is located in a part of the wash that is cut down into the Navajo Sandstone. Dinnebito Wash is an intermittent stream with small sections that flow year round, though most of the stream is dry much of the year. The streamflow gaging station is in a perennial reach (fig. 13). From July through September monsoon rain events can result in large sediment laden flows in Dinnebito Wash. The maximum instantaneous discharge recorded at the gage was 3,970 ft³/s on September 20, 2004. The minimum daily mean discharge recorded at the gage was 0.05 ft³/s on August 16, 23, and October 1–6, 2002.




Figure 13. Plots of daily mean discharge for *A*, Moenkopi Wash at Moenkopi (09401260); *B*, Dinnebito Wash near Sand Springs (09401110); *C*, Polacca Wash near Second Mesa (09400568); and *D*, Pasture Canyon Springs (09401265), Black Mesa area, northeastern Arizona, calendar year 2019.

Table 10. Streamflow-gaging stations used in the Black Mesa monitoring program, their periods of record, and drainage areas.

[mi², square miles; ---, not determined]

Station name	Station number	Date data collection began	Drainage area (mi ²)
Moenkopi Wash at Moenkopi	09401260	July 1976	1,629
Dinnebito Wash near Sand Springs	09401110	June 1993	473
Polacca Wash near Second Mesa	09400568	April 1994	905
Pasture Canyon Springs	09401265	August 2004	---

There are no observed N-aquifer springs issuing directly from the Navajo Sandstone near the streamflow gage. During base flow conditions, flow seems to initiate from Dinnebito Wash alluvium, but it is assumed this flow is supplied by the N aquifer from below and through the alluvium.

Polacca Wash

Polacca Wash has a drainage area of 905 mi² and drains a large section of the eastern part of Black Mesa. The streamflow gage is in a part of the wash that is cut down into the Kayenta Formation. Much of Polacca Wash is ephemeral, remaining dry except during and after precipitation runoff events. However, the streamflow gage is in a stream reach that often has flow. During the period of

streamflow-gage operation, there has been continuous flow at the gage for most months of the year with the exception of the summer months, when the stream is often dry at the gage (fig. 13). From July through September, monsoon rain events can result in large sediment-laden flows in Polacca Wash. The maximum instantaneous discharge recorded at the gage was 2,140 ft³/s on July 30, 2017. Most of the base flow at the Polacca Wash streamflow gage is likely provided by a spring issuing from the base of the Navajo Sandstone located about 1 mi upstream of the gage.

Pasture Canyon Springs

Pasture Canyon Springs discharges to a small perennial stream that begins near the head of Pasture Canyon, a narrow box canyon carved into the Navajo Sandstone. Base flow begins near the head of the canyon from a piped spring. Discharge from that spring accounts for around 20 percent of the total flow measured at the streamflow gage, which is located approximately 1/4 mi downstream from the spring. The remaining base flow measured at the streamflow gage comes from additional springs issuing through the alluvium between the head of the canyon and the gage. Because the drainage area is small, very little surface runoff from rainstorms or snowmelt occurs above the Pasture Canyon Springs streamflow gage (fig. 13). In addition, most of the alluvium in the wash is composed of reworked dune sand, so precipitation tends to infiltrate rather than run off. During the operational period of record for the gage, the minimum daily mean discharge recorded was 0.14 ft³/s on September 5–8, 2017, and the maximum instantaneous discharge was 4.96 ft³/s on October 3, 2018.

Surface-Water Base Flow

Trends in the groundwater-discharge component of total flow at the four streamflow-gaging stations were evaluated on the basis of the median of 120 consecutive daily mean flows for four winter months (November through February) as a surrogate measure for base flow (fig. 14). Median winter streamflow is reported as the year in which the winter season began, which for this report is 2018. Groundwater discharge was assumed to be constant throughout the year, and the median winter flow was assumed to represent constant annual groundwater discharge. Most flow that occurs during the winter is groundwater discharge; rainfall and snowmelt runoff are infrequent, and evapotranspiration is at a minimum during the winter. Not all groundwater discharge ends up as surface-water flow; during summer months some or all of the groundwater discharge can be taken up by plants or evaporated directly to the atmosphere. Rather than the average flow, the median flow for November, December, January, and February is used to estimate groundwater discharge because the median is less affected by occasional winter runoff. Nonetheless, the median flow for November, December, January, and February is an estimate of groundwater discharge rather than a calculation of base-flow groundwater discharge. A more rigorous and accurate calculation of base-flow would involve detailed evaluations of streamflow hydrographs, flows into and out of bank storage, gain and loss of

streamflow as it moves down the stream channel, and interaction of groundwater in the N aquifer with groundwater in the shallow alluvial aquifers in the stream valleys. The median winter flow, however, is useful as a consistent, easily measurable index for evaluating possible temporal trends in groundwater discharge.

Median winter flows calculated for winter 2018 were 2.7 ft³/s for Moenkopi Wash at Moenkopi, 0.31 ft³/s for Dinnebito Wash near Sand Springs, 0.07 ft³/s for Polacca Wash near Second Mesa, and 0.43 ft³/s for Pasture Canyon Springs (fig. 14A–D). A significant decreasing trend in median winter flows, calculated using Kendall's tau ($p < 0.05$), is indicated at the Moenkopi Wash and Dinnebito Wash streamflow-gaging stations, but no significant trends are indicated at the Polacca Wash and Pasture Canyon Springs streamflow-gaging stations (fig. 14A–D).

Water Chemistry

Water samples for water-chemistry analyses were collected in April, July, and August 2019 from selected wells and springs as part of the Black Mesa monitoring program. Field measurements were made and water samples were analyzed for major ions, nutrients, and the trace elements arsenic, boron, and iron. Field measurements were made in accordance with standard USGS protocols documented in the USGS National Field Manual for the Collection of Water-Quality Data (U.S. Geological Survey, variously dated). Field measurements include pH, specific conductance, temperature, dissolved oxygen, alkalinity, and discharge rates at springs. Field alkalinities were determined using incremental equivalence. Dissolved constituent samples were filtered through a 0.45-micron pore size filter and preserved according to sampling and analytical protocols. Laboratory analyses for samples were done at the USGS National Water Quality Laboratory according to techniques described in Fishman and Friedman (1989), Fishman (1993), Struzeski and others (1996), and Garbarino and others (2006).

Quality assurance for this study was maintained through the use of standard USGS training of field personnel, use of standard USGS field protocols (U.S. Geological Survey, variously dated), collection of quality control (QC) samples, and thorough review of the analytical results. All USGS scientists involved with this study have participated in the USGS National Field Quality Assurance Program.

A sequential replicate QC sample was collected at Moenkopi School Spring to better understand potential variability associated with field conditions and field and laboratory procedures (table 11). For all constituents except orthophosphate, relative percent differences (difference in sample results divided by the mean result) ranged from 0 to 6.1 percent. Orthophosphate values from the environmental and replicate samples differed by 22.2 percent. The percent difference for orthophosphate is larger than desired, but the magnitude of the difference is small, about 0.001 milligrams per liter (mg/L). The one replicate sample suggests that there is an acceptably low level of variability affecting the data; no QC data was collected to assess potential bias affecting the data.




Figure 14. Plots of median winter discharge for November through February for A, Moenkopi Wash at Moenkopi (09401260); B, Dinnebito Wash near Sand Springs (09401110); C, Polacca Wash near Second Mesa (09400568); and D, Pasture Canyon Springs (09401265), Black Mesa area, northeastern Arizona, winter 1977–2018. Median winter flow is calculated by computing the median flow for 120 consecutive daily mean flows for the winter months of November, December, January, and February.

In past years water-chemistry samples were systematically collected from as many as 12 different wells as part of the Black Mesa monitoring program. The number of wells sampled has been reduced owing to budgetary constraints. Three industrial wells were sampled in 2019, Peabody 3, Peabody 7, and Peabody 9. Since 1989, samples have been collected from the same four springs—Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso and in 2019 all four springs were sampled. Long-term data for specific conductance, dissolved solids, chloride, and sulfate for the wells and springs sampled each year are shown in the annual reports (table 2). These constituents are monitored on an annual basis

because an increase in the concentration of these constituents in the N aquifer could indicate leakage from the overlying D aquifer. On average, the concentration of dissolved solids in water from the D aquifer is about 7 times greater than that of water from the N aquifer; concentration of chloride ions is about 11 times greater, and concentration of sulfate ions is about 30 times greater (Eychaner, 1983). Historical data for other constituents for all the wells and springs in the Black Mesa study area are available from the USGS NWIS Web Interface for water-quality data (<https://waterdata.usgs.gov/nwis/qw>), and they can be found in monitoring reports cited in the “Previous Investigations” section of this report and listed in table 2.

Table 11. Comparison of chemical analyses of environmental and replicate water samples taken from Moenkopi School Spring (360632111131101), Black Mesa area, northeastern Arizona, 2019.

[mg/L, milligrams per liter; µg/L, micrograms per liter; °C, degrees Celsius; <, less than; --- no data]

Physical property or chemical analysis	Water samples		Relative Percent difference
	Environmental	Replicate	
Common well name	Moenkopi School Spring	Moenkopi School Spring	---
U.S. Geological Survey identification number	360632111131101	360632111131101	---
Date of samples	04–22–19	04–22–19	---
Alkalinity, field, dissolved (mg/L as CaCO ₃)	95.0	91.6	3.6
Nitrogen, NO ₂ + NO ₃ , dissolved (mg/L as N)	2.87	2.88	–0.3
Orthophosphate, dissolved (mg/L as P)	0.004	0.005	–22.2
Calcium, dissolved (mg/L as Ca)	43.5	43.4	0.2
Magnesium, dissolved (mg/L as Mg)	8.99	9.00	–0.1
Potassium, dissolved (mg/L as K)	1.50	1.46	2.7
Sodium, dissolved (mg/L as Na)	32.9	32.9	0.0
Chloride, dissolved (mg/L as Cl)	39.3	39.3	0.0
Fluoride, dissolved (mg/L as F)	0.16	0.16	0.0
Silica, dissolved (mg/L as SiO ₂)	13.6	13.6	0.0
Sulfate, dissolved (mg/L as SO ₄)	49.1	49.1	0.0
Arsenic, dissolved (µg/L as As)	2.4	2.4	0.0
Boron, dissolved (µg/L as B)	51	51	0.0
Iron, dissolved (µg/L as Fe)	<10.0	<10.0	---
Dissolved solids, residue at 180 °C (mg/L)	255	271	–6.1

Water-Chemistry Data for Wells Completed in the N Aquifer

Previous monitoring (table 2) has found that the primary types of water in the N aquifer in the Black Mesa study area are calcium bicarbonate water and sodium bicarbonate water. Calcium bicarbonate water is mostly found in the recharge and unconfined areas of the northern and northwestern parts of the Black Mesa study area. Sodium bicarbonate water is mostly found in the area that is confined and downgradient to the south and east (Lopes and Hoffmann, 1997). Water-chemistry results from well samples in 2019 are similar to results from previous years and are presented in figures 15 and 16 and in tables 12 and 13. The three wells are located in the confined part of the N aquifer, and all had sodium bicarbonate water in 2019.

Chemical constituents analyzed from the three wells were compared to the U.S. Environmental Protection Agency (EPA) primary and secondary drinking water standards (U.S. Environmental Protection Agency, 2003). Maximum contaminant levels (MCLs), which are the primary regulations, are legally enforceable standards that apply to public water systems. They protect drinking-water quality by limiting the levels of specific contaminants that can adversely affect public health. Secondary maximum contaminant levels (SMCLs) provide guidelines for the control of contaminants that may cause cosmetic effects

(such as skin or tooth discoloration) or aesthetic effects (such as taste, odor, or color) in drinking water. The EPA recommends compliance with SMCLs for public water systems; however, compliance is not enforced.

In 2019, all of the analyzed constituents from the three wells were below the EPA MCLs for drinking water. However, all three wells had pH values that were higher than the SMCL pH range (6.5–8.5) (table 12). The SMCL for pH is a guideline, waters with a pH higher than the SMCL may have a slippery feel, soda taste, and cause water deposits (U.S. Environmental Protection Agency, 2003). All other analyzed constituents were below the SMCL for drinking water (table 12).

Water-Chemistry Data for Springs that Discharge from the N Aquifer

In 2019, water samples were collected from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso (figs. 11, 15). Geologic maps and field observations indicate that these four springs discharge water from the unconfined part of the N aquifer. At Moenkopi School Spring, samples were collected from a horizontal metal pipe built into the hillside to collect water from the spring. At Burro Spring, samples were collected




Figure 15. Map showing water chemistry and distribution of dissolved solids at wells and springs in the N aquifer, Black Mesa area, northeastern Arizona, 2019.




Figure 16. Plot of dissolved-solids concentrations for water samples from selected wells completed in the N aquifer, Black Mesa area, northeastern Arizona: Peabody 3, 1971–2019; Peabody 7, 1972–2019; and Peabody 9, 1986–2019.

Table 12. Physical properties and chemical analyses of water samples from selected industrial wells completed in the N aquifer, Black Mesa area, northeastern Arizona, 2019.

[°C, degrees Celsius; $\mu\text{S}/\text{cm}$, microsiemens per centimeter at 25 °C; mg/L , milligrams per liter; $\mu\text{g}/\text{L}$, micrograms per liter; $<$, less than]

Common well name	U.S. Geological Survey identification number	Date of samples	Temperature, field (°C)	Specific conductance, field ($\mu\text{S}/\text{cm}$)	pH, field (units)	Alkalinity, field, dissolved (mg/L as CaCO_3)	Nitrogen, $\text{NO}_2 + \text{NO}_3$, dissolved (mg/L as N)	Ortho-phosphate, dissolved (mg/L as P)	Calcium, dissolved (mg/L as Ca)	Magnesium, dissolved (mg/L as Mg)	Potassium, dissolved (mg/L as K)
Peabody 3	362625110223701	08–23–19	32.1	218	9.1	79.7	0.950	0.009	4.96	0.042	0.77
Peabody 7	362456110242301	08–09–19	31.3	222	9.3	72.8	0.861	0.010	3.16	0.026	0.67
Peabody 9	362333110250001	07–27–19	34.1	182	9.1	58.6	0.766	0.027	3.85	0.047	0.55

Common well name	U.S. Geological Survey identification number	Date of samples	Sodium, dissolved (mg/L as Na)	Chloride, dissolved (mg/L as Cl)	Fluoride, dissolved (mg/L as F)	Silica, dissolved (mg/L as SiO_2)	Sulfate, dissolved (mg/L as SO_4)	Arsenic, dissolved ($\mu\text{g}/\text{L}$ as As)	Boron, dissolved ($\mu\text{g}/\text{L}$ as B)	Iron, dissolved ($\mu\text{g}/\text{L}$ as Fe)	Dissolved solids, residue at 180°C (mg/L)
Peabody 3	362625110223701	08–23–19	43.7	4.03	0.21	20.4	15.1	3.1	29	<10.0	147
Peabody 7	362456110242301	08–09–19	92	3.10	0.16	21.2	15.1	2.9	23	<10.0	146
Peabody 9	362333110250001	07–27–19	30.2	2.05	0.16	20.5	3.24	2.8	19	10.8	97

Table 13. Specific conductance and concentrations of selected chemical constituents in water samples from selected industrial wells completed in the N aquifer, Black Mesa area, northeastern Arizona, 1968–2019.

[µS/cm, microsiemens per centimeter at 25 °C; °C, degrees Celsius; mg/L, milligram per liter; ---, no data]

Year	Specific conductance, field (µS/cm)	Dissolved solids, residue at 180 °C (mg/L)	Chloride, dissolved (mg/L as Cl)	Sulfate, dissolved (mg/L as SO ₄)	Year	Specific conductance, field (µS/cm)	Dissolved solids, residue at 180 °C (mg/L)	Chloride, dissolved (mg/L as Cl)	Sulfate, dissolved (mg/L as SO ₄)
Peabody 3					Peabody 7—Continued				
1968	236	^a 171	4.0	17.0	1980	210	136	3.7	11.0
1971	247	^a 159	3.5	17.0	1988	240	151	4.4	18.0
1973	---	^a 213	---	---	1993	228	128	3.8	16.0
1974	230	154	3.2	13.0	1994	225	130	3.80	16.0
1975	240	157	2.8	14.0	2019	222	146	3.10	15.1
1976	250	145	3.0	11.0	Peabody 9				
1977	230	^a 159	3.2	12.0	1986	181	^a 134	3.1	4.9
1979	240	^a 171	3.3	17.0	1987	148	102	2.8	4.1
1980	230	151	3.5	14.0	1990	158	106	1.6	3
1987	171	112	3.6	6.1	1991	155	83	2.70	3.10
1993	177	169	4.2	19.0	1993	157	94	1.6	2.90
2019	218	147	4.03	15.1	1995	154	122	1.60	1.60
Peabody 7					1998	109	109	1.68	2.48
1972	222	^a 157	2.5	20.0	2001	88	112	1.83	2.72
1973	---	^a 151	---	---	2002	152	96	1.66	2.54
1974	210	141	3.3	14.0	2006	---	111	2.59	4.43
1975	220	146	3.0	15.0	2013	159	114	2.03	3.42
1976	240	140	3.2	13.0	2019	182	97	2.05	3.24
1979	225	^a 159	3.4	20.0	^a Value determined by sum of constituents.				

from the end of a pipe that fills a trough for cattle. At Pasture Canyon Spring, samples were collected from a pipe at the end of a channel that is approximately 50 ft away from the spring. At Unnamed Spring near Dennehotso, samples were collected from a pool along the bedrock wall from which the spring discharges.

Samples from all springs yielded a calcium bicarbonate-type water, except Burro Spring, which had a sodium-calcium bicarbonate-type water (fig. 15; table 14). Dissolved solid concentrations measured 255 mg/L at Moenkopi School Spring, 332 mg/L at Burro Spring, 148 mg/L at Pasture Canyon Spring, and 105 mg/L at Unnamed Spring near Dennehotso (tables 14, 15). Chloride concentration was highest at Moenkopi School Spring (39.3 mg/L; tables 14, 15). Concentration of sulfate was highest at Burro Spring (63.0 mg/L; tables 14, 15). Concentrations of all the analyzed constituents in samples from all four springs were less

than current EPA MCLs and SMCLs (U.S. Environmental Protection Agency, 2003).

There are significant increasing trends in concentrations of dissolved solids, chloride, and sulfate in water from Moenkopi School Spring ($p < 0.05$) (table 15; fig. 17). Concentrations of the same constituents from Pasture Canyon Spring, and Unnamed Spring near Dennehotso did not show any significant trends (table 15; fig. 17). However, in 2010, 2011, and 2012, Unnamed Spring near Dennehotso showed an increase in dissolved solids concentrations which may be a result of sampling from an alternate sample location. Since then, Unnamed Spring near Dennehotso has been sampled from the same location that was used prior to 2010 and the results for dissolved-solids analysis returned to levels observed prior to 2010 (fig. 17). Concentrations of dissolved solids and chloride from Burro Spring did not show significant trends; however, concentrations of sulfate from Burro Spring now show a decreasing trend.

Table 14. Physical properties and chemical analyses of water samples from four springs that discharge from the N aquifer, Black Mesa area, northeastern Arizona, 2019.

[°C, degree Celsius; µS/cm, microsiemens per centimeter at 25 °C; mg/L, milligrams per liter; µg/L, micrograms per liter; <, less than]

U.S. Geological Survey identification number	Bureau of Indian Affairs site number	Common spring name	Date of samples	Temperature, field (°C)	Specific conductance, field (µS/cm)	pH, field (units)	Alkalinity, field, dissolved (mg/L as CaCO ₃)	Nitrogen, NO ₂ + NO ₃ , dissolved (mg/L as N)	Ortho-phosphate, dissolved (mg/L as P)	Calcium, dissolved (mg/L as Ca)	Magnesium, dissolved (mg/L as Mg)	Potassium, dissolved (mg/L as K)
360632111131101	3GS-77-6	Moenkopi School Spring	04-22-19	17.5	428	7.5	95.0	2.87	0.004	43.5	8.99	1.50
354156110413701	6M-31	Burro Spring	04-22-19	12.5	522	7.7	160	<0.040	<0.004	53.7	3.87	<0.30
361021111115901	3A-5	Pasture Canyon Spring	04-22-19	16.7	224	7.9	68.9	4.58	0.021	30.6	4.53	1.41
364656109425400	8A-224	Unnamed Spring near Dennehotso	04-23-19	12.7	156	7.6	68.9	1.46	0.019	26.4	3.99	1.08

U.S. Geological Survey identification number	Bureau of Indian Affairs site number	Common spring name	Date of samples	Sodium, dissolved (mg/L as Na)	Chloride, dissolved (mg/L as Cl)	Fluoride, dissolved (mg/L as F)	Silica, dissolved (mg/L as SiO ₂)	Sulfate, dissolved (mg/L as SO ₄)	Arsenic, dissolved (µg/L as As)	Boron, dissolved (µg/L as B)	Iron, dissolved (µg/L as Fe)	Dissolved solids, residue at 180°C (mg/L)
360632111131101	3GS-77-6	Moenkopi School Spring	04-22-19	32.9	39.3	0.16	13.6	49.1	2.4	51	<10.0	255
354156110413701	6M-31	Burro Spring	04-22-19	58.6	23.7	0.35	14.3	63.0	0.7	75	<10.0	332
361021111115901	3A-5	Pasture Canyon Spring	04-22-19	12.4	5.02	0.16	10.5	16.8	1.8	36	<10.0	148
364656109425400	8A-224	Unnamed Spring near Dennehotso	04-23-19	4.75	2.39	0.14	12.7	6.02	2.8	17	<10.0	105

Table 15. Specific conductance and concentrations of selected chemical constituents in water samples from four springs that discharge from the N aquifer, Black Mesa area, northeastern Arizona, 1948–2019.

[μS/cm, microsiemens per centimeter at 25 °C; °C, degrees Celsius; mg/L, milligrams per liter; ---, no data]

Year	Specific conductance, field (μS/cm)	Dissolved solids, residue at 180 °C (mg/L)	Chloride, dissolved (mg/L as Cl)	Sulfate, dissolved (mg/L as SO ₄)	Year	Specific conductance, field (μS/cm)	Dissolved solids, residue at 180 °C (mg/L)	Chloride, dissolved (mg/L as Cl)	Sulfate, dissolved (mg/L as SO ₄)
Moenkopi School Spring					Burro Spring—Continued				
1952	222	---	6	---	1999	545	346	24.8	69.2
1987	270	161	12.0	19.0	2001	480	348	23.6	67.8
1988	270	155	12.0	19.0	2002	591	374	30.6	77.0
1991	297	157	14.0	20.0	2003	612	374	30.5	81.1
1993	313	204	17.0	27.0	2004	558	337	24.9	63.6
1994	305	182	17.0	23.0	2005	558	357	25.8	68.9
1995	314	206	18.0	22.0	2006	576	359	25.0	68.2
1996	332	196	19.0	26.0	2009	577	372	25.7	72.5
1997	^a 305	185	17.8	23.8	2010	583	355	25.9	71.5
1998	296	188	17.6	23.7	2011	560	353	25.7	69.5
1999	305	192	18.7	25.6	2012	553	330	23.1	64.7
2001	313	194	18.3	25.5	2013	560	350	24.4	67.7
2002	316	191	18.3	23.1	2014	549	360	22.8	64.6
2003	344	197	18.6	23.4	2016	544	318	22.2	61.7
2004	349	196	19.1	21.3	2017	536	329	22.0	59.5
2005	349	212	23.3	29.6	2018	532	330	21.6	58.7
2006	387	232	27.2	34.2	2019	522	332	23.7	63.0
2007	405	^b 253	30.6	39.9	Pasture Canyon Spring				
2008	390	230	28.3	37.6	1948	^a 227	(^c)	6.0	13
2009	381	240	27.0	35.6	1982	240	---	5.1	18.0
2010	480	217	26.2	33.4	1986	257	---	5.4	19.0
2011	374	216	28.5	36.2	1988	232	146	5.3	18.0
2012	382	218	27.5	33.3	1992	235	168	7.10	17.0
2013	370	220	27.2	33.3	1993	242	134	5.3	17.0
2014	382	226	28.5	34.6	1995	235	152	4.80	14.0
2016	427	244	34.6	43.8	1996	238	130	4.70	15.0
2017	424	266	35.5	44.7	1997	232	143	5.27	16.9
2018	434	260	36.9	45.9	1998	232	147	5.12	16.2
2019	428	255	39.3	49.1	1999	235	142	5.06	14.2
Burro Spring					2001	236	140	5.06	17.0
1989	485	308	22.0	59	2002	^d 243	143	5.14	16.5
1990	^a 545	347	23.0	65.0	2003	236	151	5.09	16.1
1993	595	368	30.0	85.0	2004	248	150	5.50	16.4
1994	^a 597	368	26.0	80.0	2005	250	149	5.07	16.3
1996	525	324	23.0	62.0	2008	240	149	5.01	18.3
1997	^a 511	332	26.0	75.0	2009	241	160	5.10	18.6
1998	504	346	24.6	70.4	2010	314	157	5.25	17.9

Table 15.—Continued

Year	Specific conductance, field ($\mu\text{S}/\text{cm}$)	Dissolved solids, residue at 180 °C (mg/L)	Chloride, dissolved (mg/L as Cl)	Sulfate, dissolved (mg/L as SO_4)
Pasture Canyon Spring—Continued				
2011	236	146	5.47	18.5
2012	248	142	5.20	17.5
2013	245	145	5.16	17.7
2014	249	149	5.03	17.2
2016	252	155	5.09	17.2
2017	236	151	5.07	17.1
2018	238	145	5.04	16.9
2019	224	148	5.02	16.8
Unnamed Spring near Dennehotso				
1984	195	112	2.8	7.1
1987	178	^c 109	3.4	7.5
1992	178	108	3.60	7.30
1993	184	100	3.2	8.00
1995	184	124	2.60	5.70
1996	189	112	2.80	8.20
1997	^a 170	98	2.40	6.10
1998	179	116	2.43	5.36
1999	184	110	2.76	6.30
2001	176	116	2.61	5.96
2002	183	104	2.67	7.38
2003	180	118	2.95	7.16
2004	170	117	2.72	5.05
2005	194	114	2.65	8.67
2010	259	155	9.38	15.5
2011	292	172	14.5	24.1
2012	298	179	13.5	21.9
2013	196	127	3.06	8.24
2014	160	122	2.68	7.40
2016	197	116	3.21	9.46
2018	157	108	2.33	5.39
2019	156	105	2.39	6.02

^aValue is different in Black Mesa monitoring reports before 2000. Earlier reports showed values determined by laboratory analysis.

^bValue is different in Black Mesa monitoring reports prior to this report. Earlier reports showed values determined by the sum of constituents.

^cValue is different in Black Mesa monitoring reports before 2000. Earlier reports showed values determined by the sum of constituents.

^dValue was measured in the laboratory.

Summary

The N aquifer is extensive and the primary source of groundwater for industrial and municipal users in the Black Mesa area of northeastern Arizona. Availability of quality water is an important issue in the Black Mesa area because of past industrial use and continued municipal use, a growing population, and limited precipitation. This report presents results of groundwater and water-chemistry monitoring in the Black Mesa area from January 2019 to December 2019, and additionally uses streamflow statistics from November 2018 to December 2019. These monitoring data are compared to historical data from the 1950s through December 2018.

In 2019, total groundwater withdrawals were about 3,070 acre-ft; industrial withdrawals were about 670 acre-ft, and municipal withdrawals were about 2,400 acre-ft. From the prestress period (before 1965) to 2019, water levels in the unconfined areas of the N aquifer had a median change of -1.7 ft, and the changes ranged from -40.0 ft to $+8.2$ ft. Water levels in the confined area of the N aquifer had a median change of -38.8 ft, and the changes ranged from -185.0 ft to $+12.9$ ft.

Discharge has been measured annually at Moenkopi School Spring and Pasture Canyon Spring and intermittently at Burro Spring and Unnamed Spring near Dennehotso. For the period of record, discharge at Moenkopi School Spring and Unnamed Spring near Dennehotso has fluctuated, and the data indicate a decreasing trend in discharge for both springs; however, no trend is apparent for either Burro Spring or Pasture Canyon Spring.

Streamflow was measured continuously at four streamflow-gaging stations—Moenkopi Wash, Dinnebito Wash, Pasture Canyon Springs, and Polacca Wash—and varied during the periods of record. Median flows for November through February of each winter are used as an indicator of groundwater discharge to those streams. For the period of record at Moenkopi Wash and Dinnebito Wash, winter flows indicate a decreasing trend in discharge. Winter flows at Polacca Wash and Pasture Canyon Springs have fluctuated but show neither a significant increase nor decrease.

In 2019, water samples were collected from three wells and four springs and analyzed for selected chemical constituents. A replicate quality assurance sample suggests that there is an acceptably low level of variability affecting the data. Dissolved-solids concentrations in water samples from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso were 255 mg/L, 332 mg/L, 148 mg/L, and 105 mg/L, respectively. From the mid-1980s to 2019, long-term data from Moenkopi School Spring indicate increasing concentrations of dissolved solids, chloride, and sulfate. Concentrations of dissolved solids, chloride, and sulfate from Pasture Canyon Spring and Unnamed Spring near Dennehotso do not indicate a trend for the period of record. Concentrations of dissolved solids and chloride at Burro Spring also have varied for the period of record, but there is no statistical trend in the data; however, concentrations of sulfate from Burro Spring now show a trend towards lower concentrations.




Figure 17. Plots of concentrations of dissolved solids, chloride, and sulfate for water samples from Moenkopi School Spring, Burro Spring, Pasture Canyon Spring, and Unnamed Spring near Dennehotso, which discharge from the N aquifer, Black Mesa area, northeastern Arizona, 1982–2019. A, Dissolved solids; B, chloride; and C, sulfate.

References Cited

- Aubrey, W.M., 1992, New interpretations of the stratigraphy and sedimentology of uppermost Jurassic to lowermost Cretaceous strata in the San Juan Basin of northwestern New Mexico: U.S. Geological Survey Bulletin, 1808–J, 17 p.
- Beitler, B., Parry, W.T., and Chan, M.A., 2003, Bleaching of Jurassic Navajo Sandstone on Colorado Plateau Laramide highs—Evidence of exhumed hydrocarbon supergiants?: *Geology*, v. 31, p. 1041–1044.
- Beitler, B., Parry, W.T., and Chan, M.A., 2005, Fingerprints of fluid flow—Chemical diagenetic history of the Jurassic Navajo Sandstone, southern Utah, U.S.A.: *Journal of Sedimentary Research*, v. 75, p. 547–561.
- Blakey, R.C., 2008, Pennsylvanian–Jurassic sedimentary basins of the Colorado Plateau and Southern Rocky Mountains, in Miall, A.D., ed., *The sedimentary basins of the United States and Canada*: Amsterdam, Elsevier, p. 245–296.
- Blakey, R.C., Peterson, F., Caputo, M.V., Geesaman, R.C., and Voorhees, B.J., 1983, Paleogeography of Middle Jurassic continental, shoreline, and shallow marine sedimentation, southern Utah, in Reynolds, M.W., and Dolly, E.D., eds., *Mesozoic paleogeography of west-central United States*: Society of Economic Paleontologists and Mineralogists, Rocky Mountain Section, Symposium, v. 2, p. 77–100.
- Blakey, R.C., and Ranney, W., 2008, Ancient landscapes of the Colorado Plateau: Grand Canyon Arizona, Grand Canyon Association, 156 p.
- Billingsley, G.H., Block, D., and Hiza Redsteer, M., 2013, Geologic map of the Winslow 30' x 60' quadrangle, Coconino and Navajo Counties, northern Arizona: U.S. Geological Survey Scientific Investigations Map 3247, pamphlet 25 p., 3 sheets, scale 1:50,000, accessed June 2018, at <http://pubs.usgs.gov/sim/3247/>.
- Billingsley, G.H., Stoffer, P.W., and Priest, S.S., 2012, Geologic map of the Tuba City 30' x 60' quadrangle, Coconino County, northern Arizona: U.S. Geological Survey Scientific Investigations Map 3227, pamphlet 31 p., 3 sheets, scale 1:50,000, accessed November 2017, at <https://pubs.usgs.gov/sim/3227/>.
- Boner, F.C., Davis, R.G., and Duet, N.R., 1992, Water-resources data, Arizona, water year 1991: U.S. Geological Survey Water-Data Report AZ–91–1, 411 p.
- Boner, F.C., Garrett, W.B., and Konieczki, A.D., 1989, Water-resources data, Arizona, water year 1988: U.S. Geological Survey Water-Data Report AZ–88–1, 391 p.
- Boner, F.C., Konieczki, A.D., and Davis, R.G., 1991, Water-resources data, Arizona, water year 1990: U.S. Geological Survey Water-Data Report AZ–90–1, 381 p.
- Boner, F.C., Smith, C.F., Garrett, W.B., and Konieczki, A.D., 1990, Water-resources data, Arizona, water year 1989: U.S. Geological Survey Water-Data Report AZ–89–1, 383 p.
- Brown, J.G., and Eychaner, J.H., 1988, Simulation of five ground-water withdrawal projections for the Black Mesa area, Navajo and Hopi Indian Reservations, Arizona: U.S. Geological Survey Water-Resources Investigations Report 88–4000, 51 p.
- Cooley, M.E., Harshbarger, J.W., Akers, J.P., and Hardt, W.F., 1969, Regional hydrogeology of the Navajo and Hopi Indian Reservations, Arizona, New Mexico, and Utah: U.S. Geological Survey Professional Paper 521–A, 61 p.
- Daly, C., Neilson, R.P., and Phillips, D.L., 1994, A statistical-topographic model for mapping climatological precipitation over mountainous terrain: *Journal of Applied Meteorology*, v. 33, p. 140–158.
- Davis, G.E., Hardt, W.F., Thompson, L.K., and Cooley, M.E., 1963, Records of ground-water supplies, part 1 of geohydrologic data in the Navajo and Hopi Indian Reservations, Arizona, New Mexico, and Utah: Arizona State Land Department Water-Resources Report 12–A, 159 p.
- Dubiel, R.F., 1989, Sedimentology and revised nomenclature for the upper part of the Upper Triassic Chinle Formation and the Lower Jurassic Wingate Sandstone, northwestern New Mexico and northeastern Arizona, in Anderson, O.J., Lucas, S.G., Love, D.W., and Cather, S.M., eds., *Southeastern Colorado Plateau: New Mexico Geological Society 40th Annual Fall Field Conference Guidebook*, p. 213–223.
- Eychaner, J.H., 1983, Geohydrology and effects of water use in the Black Mesa area, Navajo and Hopi Indian Reservations, Arizona: U.S. Geological Survey Water-Supply Paper 2201, 26 p.
- Fishman, M.J., ed., 1993, Methods of analysis by the U.S. Geological Survey National Water Quality Laboratory—Determination of inorganic and organic constituents in water and fluvial sediments: U.S. Geological Survey Open-File Report 93–125, 217 p.
- Fishman, M.J., and Friedman, L.C., 1989, Methods for determination of inorganic substances in water and fluvial sediments: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A1, 545 p.
- Fisk, G.G., Duet, N.R., Evans, D.W., Angereth, C.E., Castillo, N.K., and Longworth, S.A., 2004, Water-resources data, Arizona, water year 2003: U.S. Geological Survey Water-Data Report AZ–03–1, 326 p.
- Fisk, G.G., Duet, N.R., McGuire, E.H., Angereth, C.E., Castillo, N.K., and Smith, C.F., 2005, Water-resources data, Arizona, water year 2004: U.S. Geological Survey Water-Data Report AZ–04–1, 415 p.

- Fisk, G.G., Duet, N.R., McGuire, E.H., Roberts, W.P., Castillo, N.K., and Smith, C.F., 2006, Water-resources data, Arizona, water year 2005: U.S. Geological Survey Water-Data Report AZ-05-1, 347 p.
- Flowers, R.M., 2010, The enigmatic rise of the Colorado Plateau: *Geology*, v. 38, no. 7, p. 671–672.
- Franczyk, K.J., 1988, Stratigraphic revision and depositional environments of the Upper Cretaceous Toreva Formation in the northern Black Mesa area, Navajo and Apache counties, Arizona: U.S. Geological Survey Bulletin, 1685, 32 p.
- Garbarino, J.R., Kanagy, L.K., and Cree, M.E., 2006, Determination of elements in natural-water, biota, sediment, and soil samples using collision/reaction cell inductively coupled plasma-mass spectrometry: U.S. Geological Survey Techniques and Methods, book 5, sec. B, chap. 1, 88 p.
- GeoTrans, Inc., 1987, A two-dimensional finite-difference flow model simulating the effects of withdrawals to the N aquifer, Black Mesa area, Arizona: Boulder, Colo., GeoTrans, Inc., report prepared for Peabody Western Coal Company.
- Harshbarger, J.W., Lewis, D.D., Skibitzke, H.E., Heckler, W.L., and Kister, L.R., 1966, Arizona water: U.S. Geological Survey Water-Supply Paper 1648, 85 p.
- Harshbarger, J.W., Repenning, C.A., and Irwin, J.H., 1957, Stratigraphy of the uppermost Triassic and the Jurassic rocks of the Navajo country: U.S. Geological Survey Professional Paper 291, 74 p.
- Harshbarger, J.W., Repenning, C.A., and Jackson, R.L., 1951, Jurassic stratigraphy of the Navajo country: New Mexico Geological Society Guidebook, 2nd Field Conference, p. 95–99.
- Hart, R.J., and Sottolare, J.P., 1988, Progress report on the ground-water, surface-water, and quality-of-water monitoring program, Black Mesa area, northeastern Arizona—1987–88: U.S. Geological Survey Open-File Report 88–467, 27 p.
- Hart, R.J., and Sottolare, J.P., 1989, Progress report on the ground-water, surface-water, and quality-of-water monitoring program, Black Mesa area, northeastern Arizona—1988–89: U.S. Geological Survey Open-File Report 89–383, 33 p.
- Helsel, D.R., Hirsch, R.M., Ryberg, K.R., Archfield, S.A., and Gilroy, E.J., 2020, Statistical methods in water resources: U.S. Geological Survey Techniques and Methods, book 4, chap. A3, 458 p., <https://doi.org/10.3133/tm4a3>.
- Hendricks, D.M., 1985, Arizona soils: Tucson, Ariz., University of Arizona, College of Agriculture, 244 p.
- Hill, G.W., 1985, Progress report on Black Mesa monitoring program—1984: U.S. Geological Survey Open-File Report 85–483, 24 p.
- Hill, G.W., and Sottolare, J.P., 1987, Progress report on the ground-water, surface-water, and quality-of-water monitoring program, Black Mesa area, northeastern Arizona—1987: U.S. Geological Survey Open-File Report 87–458, 29 p.
- Hill, G.W., and Whetten, M.I., 1986, Progress report on Black Mesa monitoring program—1985–86: U.S. Geological Survey Open-File Report 86–414, 23 p.
- HSIGeoTrans, Inc., 1993, Investigation of the N- and D-aquifer geochemistry and flow characteristics using major ion and isotopic chemistry, petrography, rock stress analyses, and dendrochronology in the Black Mesa area, Arizona: Boulder, Colo., HSIGeoTrans, Inc., report prepared for Peabody Coal Company, 400 p.
- HSIGeoTrans, Inc., and Waterstone Environmental Hydrology and Engineering, Inc., 1999, A three-dimensional flow model of the D and N aquifers, Black Mesa Basin, Arizona: Boulder, Colo., HSIGeoTrans, Inc., and Waterstone Environmental Hydrology and Engineering, Inc., report prepared for Peabody Western Coal Company, 75 p.
- Imlay, R.W., 1980, Jurassic paleobiogeography of the conterminous United States in its continental setting: U.S. Geological Survey Professional Paper 1062, 134 p.
- Kirkland, J.I., 1991, Lithostratigraphic and biostratigraphic framework for the Mancos Shale (late Cenomanian to middle Turonian) at Black Mesa, northeastern Arizona: Geological Society of America, Special Paper 260, p. 85–111.
- Kister, L.R., and Hatchett, J.L., 1963, Selected chemical analyses of the ground water, part 2 of Geohydrologic data in the Navajo and Hopi Indian Reservations, Arizona, New Mexico, and Utah: Arizona State Land Department Water-Resources Report 12–B, 58 p.
- Lazear, G., Karlstrom, K., Aslan, A., and Kelly, S., 2013, Denudation and flexural isostatic response of the Colorado Plateau and southern Rocky Mountains region since 10 Ma: *Geosphere*, v. 9, p. 792–814.
- Lee, L., 2020, NADA—Nondetects and Data Analysis for Environmental Data, Package (ver. 1.6-1.1): The Comprehensive R Archive Network web page, accessed December 16, 2020, at <https://cran.r-project.org/package=NADA>.
- Littin, G.R., 1992, Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1990–91: U.S. Geological Survey Water-Resources Investigations Report 92–4045, 32 p.
- Littin, G.R., 1993, Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1991–92: U.S. Geological Survey Water-Resources Investigations Report 93–4111, 23 p.

- Littin, G.R., Baum, B.M., and Truini, M., 1999, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1997: U.S. Geological Survey Open-File Report 98–653, 27 p.
- Littin, G.R., and Monroe, S.A., 1995a, Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1992–93: U.S. Geological Survey Water-Resources Investigations Report 95–4156, 37 p.
- Littin, G.R., and Monroe, S.A., 1995b, Results of ground-water, surface-water, and water-chemistry monitoring, Black Mesa area, northeastern Arizona—1994: U.S. Geological Survey Water-Resources Investigations Report 95–4238, 25 p.
- Littin, G.R., and Monroe, S.A., 1996, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1995: U.S. Geological Survey Open-File Report 96–616, 22 p.
- Littin, G.R., and Monroe, S.A., 1997, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1996: U.S. Geological Survey Open-File Report 97–566, 27 p.
- Lopes, T.J., and Hoffmann, J.P., 1997, Geochemical analyses of ground-water ages, recharge rates, and hydraulic conductivity of the N Aquifer, Black Mesa area, Arizona: U.S. Geological Survey Water-Resources Investigations Report 96–4190, 42 p.
- Macy, J.P., 2009, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2007–2008: U.S. Geological Survey Open-File Report 2009–1148, 43 p.
- Macy, J.P., 2010, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2008–2009: U.S. Geological Survey Open-File Report 2010–1038, 43 p.
- Macy, J.P., and Brown, C.R., 2011, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2009–2010: U.S. Geological Survey Open-File Report 2011–1198, 41 p.
- Macy, J.P., Brown, C.R., and Anderson, J.R., 2012, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2010–2011: U.S. Geological Survey Open-File Report 2012–1102, 41 p.
- Macy, J.P., and Mason, J.P., 2017, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2013–2015: U.S. Geological Survey Open-File Report 2017–1127, 49 p., <https://doi.org/10.3133/ofr20171127>.
- Macy, J.P., and Truini, M., 2016, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2012–2013: U.S. Geological Survey Open-File Report 2015–1221, 43 p., <https://doi.org/10.3133/ofr20151221>.
- Macy, J.P., and Unema, J.A., 2014, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2011–2012: U.S. Geological Survey Open-File Report 2014–1304, 42 p., <https://doi.org/10.3133/ofr20131304>.
- Mason, J.P., and Macy, J.P., 2018, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2015–2016: U.S. Geological Survey Open-File Report 2018–1193, 60 p., <https://doi.org/10.3133/ofr20181193>.
- Mason, J.P., 2021, Groundwater, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2016–2018: U.S. Geological Survey Open-File Report 2021–1124, 50 p., <https://doi.org/10.3133/ofr20211124>.
- McCormack, H.F., Fisk, G.G., Duet, N.R., Evans, D.W., and Castillo, N.K., 2002, Water-resources data, Arizona, water year 2001: U.S. Geological Survey Water-Data Report AZ–01–1, 399 p.
- McCormack, H.F., Fisk, G.G., Duet, N.R., Evans, D.W., Roberts, W.P., and Castillo, N.K., 2003, Water-resources data, Arizona, water year 2002: U.S. Geological Survey Water-Data Report AZ–02–1, 337 p.
- McKee, E.D., and MacLachlan, M.E., 1959, Stratigraphic problems of the Wingate sandstone, Moenave formation, and related units, *in* McKee, E.D., Oriel, S.S., Ketner, K.B., MacLachlan, M.E., Goldsmith, J.W., MacLachlan, J.C., and Mudge, M.R., Paleotectonic maps of the Triassic System: U.S. Geological Survey Miscellaneous Geologic Investigations Map I-300, p. 22, pl. 8, figs. 4–6.
- Molenaar, C.M., 1983, Major depositional cycles and regional correlations of Upper Cretaceous rocks, southern Colorado Plateau and adjacent areas, *in* Reynolds, M.W., and Dolly, E.D., eds., Mesozoic paleogeography of west-central United States: Society for Sedimentary Geology Rocky Mountain Section, p. 201–224.
- O’Sullivan, R.B., Repenning, C.A., Beaumont, E.C., and Page, H.G., 1972, Stratigraphy of the Cretaceous rocks and the Tertiary Ojo Alamo Sandstone, Navajo and Hopi Indian reservations, Arizona, New Mexico, and Utah: U.S. Geological Survey Professional Paper 521–E, 65 p.
- Page, H.G., and Repenning, C.A., 1958, Late Cretaceous stratigraphy of Black Mesa, Navajo and Hopi Indian reservations, Arizona, *in* New Mexico Geological Society, Black Mesa Basin, 9th Field Conference: New Mexico Geological Society, p. 115–122.
- Peterson, F., 1988, Stratigraphic nomenclature of Middle Jurassic and Upper Jurassic rocks, western Colorado Plateau, Utah and Arizona, *in* Revisions to stratigraphic nomenclature of Jurassic and Cretaceous rocks of the Colorado Plateau: U.S. Geological Survey Bulletin, 1633–B, p. 13–56.

- Peterson, F., and Pippingos, G.N., 1979, Stratigraphic relations of the Navajo Sandstone to middle Jurassic formations, southern Utah and northern Arizona: U.S. Geological Survey Professional Paper 1035-B, 43 p.
- PRISM Climate Group, 2018, Norm81m: PRISM Climate Group, Oregon State University, accessed March 1, 2018, at <http://prism.oregonstate.edu>.
- R Development Core Team, 2020, R—A language and environment for statistical computing: Vienna, Austria, R Foundation for Statistical Computing, accessed May 2020, at <https://www.r-project.org/>.
- Raisz, E., 1972, Physiographic provinces in the Rocky Mountain region and landforms of western United States, in *Geologic atlas of the Rocky Mountain region*: Denver Colo., Rocky Mountain Association of Geologists, 331 p.
- Repenning, C.A., and Irwin, J.H., 1954, Bidahochi Formation of Arizona and New Mexico, in *Geological notes: American Association of Petroleum Geologist Bulletin*, v. 38, no. 8, p. 1821–1826.
- Sellers, W.D., and Hill, R.H., 1974, Arizona climate 1931–1972: University of Arizona, Tucson, 616 p.
- Smith, C.F., Anning, D.W., Duet, N.R., Fisk, G.G., McCormack, H.F., Pope, G.L., Rigas, P.D., and Wallace, B.L., 1995, Water-resources data, Arizona, water year 1994: U.S. Geological Survey Water-Data Report AZ-94-1, 320 p.
- Smith, C.F., Boner, F.C., Davis, R.G., Duet, N.R., and Rigas, P.D., 1993, Water-resources data, Arizona, water year 1992: U.S. Geological Survey Water-Data Report AZ-92-1, 360 p.
- Smith, C.F., Duet, N.R., Fisk, G.G., McCormack, H.F., Partin, C.K., Pope, G.L., and Rigas, P.D., 1997, Water-resources data, Arizona, water year 1996: U.S. Geological Survey Water-Data Report AZ-96-1, 328 p.
- Smith, C.F., Duet, N.R., Fisk, G.G., McCormack, H.F., Partin, C.K., Pope, G.L., Rigas, P.D., and Tadayon, S., 1996, Water-resources data, Arizona, water year 1995: U.S. Geological Survey Water-Data Report AZ-95-1, 306 p.
- Smith, C.F., Rigas, P.D., Ham, L.K., Duet, N.R., and Anning, D.W., 1994, Water-resources data, Arizona, water year 1993: U.S. Geological Survey Water-Data Report AZ-93-1, 360 p.
- Sottolare, J.P., 1992, Results of ground-water, surface-water, and water-quality monitoring, Black Mesa area, northeastern Arizona—1989–90: U.S. Geological Survey Water-Resources Investigations Report 92-4008, 38 p.
- Struzeski, T.M., DeGiacomo, W.J., and Zayhowski, E.J., 1996, Methods of analysis by the U.S. Geological Survey National Water Quality Laboratory—Determination of dissolved aluminum and boron in water by inductively coupled plasma-atomic emission spectrometry: U.S. Geological Survey Open-File Report 96-149, 17 p.
- Tadayon, S., Duet, N.R., Fisk, G.G., McCormack, H.F., Partin, C.K., Pope, G.L., and Rigas, P.D., 1999, Water-resources data, Arizona, water year 1998: U.S. Geological Survey Water-Data Report AZ-98-1, 454 p.
- Tadayon, S., Duet, N.R., Fisk, G.G., McCormack, H.F., Partin, C.K., Pope, G.L., and Rigas, P.D., 2000, Water-resources data, Arizona, water year 1999: U.S. Geological Survey Water-Data Report AZ-99-1, 389 p.
- Tadayon, S., Duet, N.R., Fisk, G.G., McCormack, H.F., Partin, C.K., Pope, G.L., and Rigas, P.D., 2001, Water-resources data, Arizona, water year 2000: U.S. Geological Survey Water-Data Report AZ-00-1, 390 p.
- Tadayon, S., Duet, N.R., Fisk, G.G., McCormack, H.F., Pope, G.L., and Rigas, P.D., 1998, Water-resources data, Arizona, water year 1997: U.S. Geological Survey Water-Data Report AZ-97-1, 416 p.
- Thomas, B.E., 2002a, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2000–2001, and performance and sensitivity of the 1988 USGS numerical model of the N aquifer: U.S. Geological Survey Water-Resources Investigations Report 02-4211, 75 p.
- Thomas, B.E., 2002b, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2001–02: U.S. Geological Survey Open-File Report 02-485, 43 p.
- Thomas, B.E., and Truini, M., 2000, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1999: U.S. Geological Survey Open-File Report 00-453, 42 p.
- Truini, M., Baum, B.M., Littin, G.R., and Shingoitewa-Honanie, G., 2000, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—1998: U.S. Geological Survey Open-File Report 00-66, 37 p.
- Truini, M., and Longworth, S.A., 2003, Hydrogeology of the D aquifer and movement and ages of ground water determined from geochemical and isotopic analyses, Black Mesa area, northeastern Arizona: U.S. Geological Survey Water-Resources Investigations Report 03-4189, 38 p.
- Truini, M., and Macy, J.P., 2005, Lithology and thickness of the Carmel Formation as related to leakage between the D and N aquifer, Black Mesa, Arizona: U.S. Geological Survey Scientific Investigations Report 2005-5187, 7 p.
- Truini, M., and Macy, J.P., 2006, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2004–05: U.S. Geological Survey Open-File Report 2006-1058, 42 p.
- Truini, M., and Macy, J.P., 2007, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2005–06: U.S. Geological Survey Open-File Report 2007-1041, 42 p.

- Truini, M., and Macy, J.P., 2008, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2006–07: U.S. Geological Survey Open-File Report 2008–1324, 33 p.
- Truini, M.t, Macy, J.P., and Porter T.J., 2005, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2003–04: U.S. Geological Survey Open-File Report 2005–1080, 44 p.
- Truini, M., and Thomas, B.E., 2004, Ground-water, surface-water, and water-chemistry data, Black Mesa area, northeastern Arizona—2002–03: U.S. Geological Survey Open-File Report 03–503, 43 p.
- U.S. Environmental Protection Agency, 2003, Current drinking water standards, national primary and secondary drinking water regulations: Washington, D.C., U.S. Environmental Protection Agency, accessed October 5, 2010, at <http://www.epa.gov/safewater/mcl.html>.
- U.S. Geological Survey, 1963–64a, Surface-water records of Arizona: 91 p.
- U.S. Geological Survey, 1963–64b, Ground-water records of Arizona: 80 p.
- U.S. Geological Survey, 1965–74a, Water-resources data for Arizona—Part 1, surface-water records: 212 p.
- U.S. Geological Survey, 1965–74b, Water-resources data for Arizona—Part 2, ground-water records: 89 p.
- U.S. Geological Survey, 1976–83, Water-resources data for Arizona, water years 1975–81: U.S. Geological Survey Water-Data Reports AZ–75–1 to AZ–81–1, [pagination varies].
- U.S. Geological Survey, 1978, Progress report on Black Mesa monitoring program—1977: U.S. Geological Survey Open-File Report 78–459, 38 p.
- U.S. Geological Survey, variously dated, National field manual for the collection of water-quality data: U.S. Geological Survey Techniques of Water-Resources Investigations, book 9, chaps. A1–A10, accessed January 2018, at <http://pubs.water.usgs.gov/twri9A>.
- White, N.D., and Garrett, W.B., 1984, Water resources data, Arizona, water year 1982: U.S. Geological Survey Water-Data Report AZ–82–1, 440 p.
- White, N.D., and Garrett, W.B., 1986, Water resources data, Arizona, water year 1983: U.S. Geological Survey Water-Data Report AZ–83–1, 387 p.
- White, N.D., and Garrett, W.B., 1987, Water resources data, Arizona, water year 1984: U.S. Geological Survey Water-Data Report AZ–84–1, 381 p.
- White, N.D., and Garrett, W.B., 1988, Water resources data, Arizona, water year 1985: U.S. Geological Survey Water-Data Report AZ–85–1, 343 p.
- Wilson, R.F., 1965, Triassic and Jurassic strata of southwestern Utah, in *Geology and resources of south-central Utah*: Utah Geological Society, Guidebook to Geology of Utah, no. 19, p. 31–46.
- Wilson, R.P., and Garrett, W.B., 1988, Water resources data, Arizona, water year 1986: U.S. Geological Survey Water-Data Report AZ–86–1, 341 p.
- Wilson, R.P., and Garrett, W.B., 1989, Water-resources data, Arizona, water year 1987: U.S. Geological Survey Water-Data Report AZ–87–1, 385 p.
- Zhu, C., 2000, Estimate of recharge from radiocarbon dating of groundwater and numerical flow and transport modeling: *Water Resources Research*, v. 36, no. 9, p. 2607–2620.
- Zhu, C., and Kipfer, R., 2010, Noble gas signatures of high recharge pulses and migrating jet stream in the late Pleistocene over Black Mesa, Arizona, United States: *Geology*, v. 38, no. 1, p. 83–86.
- Zhu, C., Waddell, R.K., Jr., Star, I., and Ostrander, M., 1998, Responses of ground water in the Black Mesa basin, northeastern Arizona, to paleoclimatic changes during the late Pleistocene and Holocene: *Geology*, v. 26, no. 2, p. 127–130.

Moffett Field Publishing Service Center
Manuscript approved September 21, 2022
Edited by Regan Austin
Illustration support by JoJo Mangano
Layout by Cory Hurd

