

TABLE 1.—DISTRIBUTION OF CONODONTS IN THE CHAPPEL LIMESTONE

[Faunal zone: B. C. = *Bactrognathus communis* zone; G. P. = *Gnathodus punctatus* zone; S. C. = *Siphonodella cooperi* zone]

PLEASE REPLACE IN POCKET IN BACK OF BOARD VOLUME (200) 5B no. 254 U.S. GEOLOGICAL SURVEY PROFESSIONAL PAPER 294-J LIBRARY

Table with columns for Plate, Figure, Locality C-1 (sections 1 and 2), Locality C-8, C-10, C-11, C-12, C-13, C-14, C-15, C-16, C-18 (sections 1 and 2), C-20, C-21, C-22, C-23, C-24, C-25, C-26, C-27, C-28, and Number of specimens of each species.

Species not found in pre-Chappel rocks of Llano region

Table listing species not found in pre-Chappel rocks of Llano region, including *Stauronathus anchoraria*, *Neopriodontus lanceolatus*, *Scalognathus anchoralis*, etc.

Species found in pre-Chappel rocks of Llano region

Table listing species found in pre-Chappel rocks of Llano region, including Mississippian species like *Polygnathus communis* and Devonian species like *Panderodella glabra*.

Summary table with columns for Locality C-1 through C-28 and a final column for Total number of specimens in collection.