

Albian stage	English standard zones (After Spath, 1924, 1930, 1941)	Killik-Chandler- Itkillik valleys, mostly northern foothills, Alaska	Recorded ranges of fossils in northern Alaska		Recorded ranges of fossils in Alberta and British Columbia		Lower Smoky River and Lower Peace River, Alberta (After McLearn and Kindler, 1950; Workman, 1954)				
			Ammonites	Other fossils	Ammonites and Inocerami	Other fossils common to northern Alaska and Canada					
Upper	<i>Stoliczkaia dispar</i>	Killik tongue of Chandler formation	?	Starfish and brittle star <i>Ditrypa cornu</i> <i>Nucula cf. N. dowlingsi</i> <i>Yoldia kissoumi</i> <i>Dicranodonta dowlingsi</i> <i>Astarte ignekensis</i> <i>Astarte portana</i> <i>Tancredia stelcki</i> <i>Tancredia kurupana</i> <i>Veniella</i> sp. <i>Flaventia? kukpovrukensis</i> <i>Solecurtus? chapmani</i> <i>Cultellus kokolikensis</i> <i>Pleuromya sikanni</i> <i>Panope? kissoumi</i> <i>Panope? elongatissima</i> <i>Goniomya matonabbei</i> <i>Thracia stelcki</i> <i>Oxytoma camseli</i> <i>Inoceramus anglicus</i> <i>Inoceramus cf. I. anglicus</i> <i>Inoceramus cf. I. cadottensis</i> <i>Camptonectes dettermanni</i> <i>Entolium utukokense</i> <i>Modiolus archisikanni</i> <i>Xenohelix? borings</i>	<i>Neogastropiles cornutus</i> and <i>Inoceramus nahwisi</i>	?	<i>Nucula dowlingsi</i> , <i>Tancredia stelcki</i> , <i>Solecurtus? chapmani</i> <i>Thracia stelcki</i> <i>Pleuromya sikanni</i> <i>Panope? kissoumi</i> <i>Modiolus archisikanni</i> (Not arranged stratigraphically)	Shaftesbury formation			
	<i>Pervinqueria inflata</i>								?	<i>Paragastropiles hardensis</i>	
Middle	<i>Euhoplites lautus</i>	Tuktu formation	?	Crinoid remains <i>Gastropiles kingi</i> and <i>Paragastropiles speikeri</i> (Ranges of species not known) <i>Cleoniceras (Grycia) sabiei</i> <i>Cleoniceras (Neosaynella)? whittingtoni</i>	?	<i>Gastropiles kingi</i> , <i>G. allani</i> , <i>G. stantoni</i> , <i>G. canadensis</i> , <i>Paragastropiles speikeri</i> (Ranges of species not known)	<i>Inoceramus cadottensis</i> and <i>I. altifluminis</i>	Peace River formation			
	<i>Hoplites dentatus</i>								?	<i>Dicranodonta dowlingsi starfish</i>	Paddy sandstone member
	<i>Douvilleiceras mammillatum</i>								?		Cadotte sandstone member
Lower	<i>Leymeriella tardefurcata</i>	Torok formation	?	<i>Cleoniceras tailleuri</i> <i>Subarthropiles belli</i> <i>Puzosia? sp. juv.</i>	?	<i>Lemuroceras? mcconnelli</i> <i>L.? irense</i> <i>Subarthropiles belli</i> <i>Inoceramus dowlingsi</i> <i>Beudanticeras affine and B. glabrum</i> <i>Placunopsis muka</i>	<i>Yoldia kissoumi</i> <i>Astarte portana</i> <i>Oxytoma camseli</i> <i>Thracia kissoumi</i> <i>Panope? elongatissima</i> <i>Goniomya matonabbei</i> (Not arranged stratigraphically)	Loon River formation			
	<i>Acanthohoplites nodosocostatum</i>								?	<i>Aucellina dowlingsi</i> (Suggested stratigraphic arrangements in Mellon and Wall, 1956)	Notikewin sandstone member
									?	<i>Falher member</i>	Spirit River formation
								Harmon shale member			
								Wilrich member			
								Blue Sky formation			

COMPARATIVE STRATIGRAPHIC DISTRIBUTION OF ALBIAN MEGAFOSSILS IN NORTHERN ALASKA, ALBERTA, AND BRITISH COLUMBIA