

Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length						
	Total in split	Perfect in split	Total in sample	Form measured	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
													Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
C-15; 435 meters	83	76	332	Living	31	O.R. 0.32-0.58	0.42						Proloculus diameter: $\cdot = 0.017$ mm $\bullet = 0.05-0.10$ mm				38°	
				Micro-spheric	30	O.R. 0.22-0.58	0.42	0.08	0.10	23.1								
				Megalo-spheric	46	O.R. 0.15-0.60	0.41	0.09	0.11	27.4								
				Total (micro-spheric plus megalospheric)	76	S.R. 0.10-0.73 O.R. 0.15-0.60	0.41 ± 0.01	0.08	0.09 ± 0.01	25.7 ± 2.1								
C-8; 450 meters	90	57	2880	Total	57	S.R. 0.13-0.70 O.R. 0.26-0.62	0.42 ± 0.01	0.08	0.09 ± 0.01	22.5 ± 2.1		Proloculus diameter: $\cdot = < 0.017$ mm $\bullet = 0.05-0.067$ mm				39°		
C-14; 800 meters	62	33	3968	Micro-spheric	38	O.R. 0.25-0.77	0.57	0.10	0.13	22.6		Proloculus diameter: $\cdot = < 0.017$ mm $\bullet = 0.067-0.10$ mm				31°		
				Megalo-spheric	143	O.R. 0.16-0.95	0.57	0.13	0.17	29.8								
				Total plus living forms	188	S.R. 0.08-1.05 O.R. 0.16-0.95	0.57 ± 0.01	0.13	0.16 ± 0.01	28.1 ± 1.5								
C-9; 855 meters	84	45	2688	Total	45	S.R. 0-1.10 O.R. 0.22-0.82	0.52 ± 0.03	0.18	0.19 ± 0.01	36.8 ± 3.9		Proloculus diameter: $\cdot = 0.017$ mm $\bullet = 0.05-0.10$ mm				27°		
C-10 1700 meters	15	3	15	Total	3	O.R. 0.33-0.67	0.52					Proloculus diameter: $\bullet = 0.050-0.084$ mm						

BOLIVINA (BOLIVINA) SEMINUDA CUSHMAN, QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens