

Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
												Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
C-15; 435 meters	43	22	172	57	S.R. 0.15-0.48 O.R. 0.16-0.42	0.31 ± 0.01	0.04	0.06 ± 0.01	17.5 ± 1.6	10 5 2	0.25 0.50 	10 5 2	0.25 0.50 Proloculus diameter: • = <0.017 mm • = 0.05 mm	0.25 0.50 	2 1 0	0.25 0.50 43°	
C-8; 450 meters	123	110	3936	110	S.R. 0.12-0.55 O.R. 0.18-0.50	0.33 ± 0.01	0.06	0.07 ± 0.004	21.3 ± 1.4	10 5 2	0.25 0.50 	15 10 5 2	0.25 0.50 Proloculus diameter: • = 0.017 mm • = 0.042-0.05 mm	0.50 0.25 	3 2 1 0	0.25 0.50 48°	
C-14; 800 meters	7	6	448	32	S.R. 0.10-0.42 O.R. 0.15-0.38	0.26 ± 0.01	0.04	0.05 ± 0.01	20.2 ± 2.5	20 10 5 2	0.25 0.50 	10 5 2	0.25 0.50 Proloculus diameter: • = <0.017 mm • = 0.042-0.059 mm	0.25 	2 1 0	0.25 0.50 47°	
C-10 1700 meters	33	25	33	25	S.R. 0.08-0.48 O.R. 0.18-0.45	0.29 ± 0.01	0.05	0.08 ± 0.01	23.5 ± 3.3	10 5 2	0.25 0.50 	10 5 2	0.25 0.50 Proloculus diameter: • = 0.017 mm • = 0.05-0.059 mm	0.25 	2 1 0	0.25 0.50 38°	

BOLIVINA (BOLIVINA) HUMILIS CUSHMAN AND MC CULLOCH, QUANTATIVE DATA

For explanation of column headings see text
X indicates living specimens