Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length							
												III. Number of chambers		IV. Width		V. Flare			
	Total in split	Perfect in split	Total in sample	*N*	Range	M	*MD*	σ	*V*	Number of specimens	Length (mm)	Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)		
C-6 80 meters	55	31	14,080	216	S.R. 0.16-0.48 O.R. 0.16-0.47	0.33 ±0.004	0.04	0.05 ±0.003	16.8 ±0.81				Proloculus diameter: •=0.017 mm						
C-17 82 meters	19	9	2432	106	S.R. 0.13-0.50 O.R. 0.16-0.48	0.31 ±0.01	0.05	0.06 ±0.004	19.8 ±1.4				Proloculus diameter: •=0.017 mm						
C-16 144 meters	5	5	5	5	O.R. 0.23-0.32	0.28	0.03						Proloculus diameter: •=0.017 mm						
C-8 450 meters	25	11	800	50	S.R. 0.15-0.45 O.R. 0.22-0.45	0.30 ±0.01	0.04	0.05 ±0.01	16.1 ±1.6				Proloculus diameter: •=0.017 mm •=0.025-0.05 mm						
C-14 800 meters	7	5	448	14	S.R. 0.12-0.40 O.R. 0.18-0.35	0.26 ±0.01	0.04	0.05 ±0.01	18.2 ±3.4				Proloculus diameter: •=0.017 mm •=0.025-0.042 mm						
C-13 1600 meters	8	7	8	7	O.R. 0.15-0.32	0.25							Proloculus diameter: •=0.017 mm •=0.033 mm						
C-10 1700 meters	2	2	2	2	O.R. 0.28-0.32	0.30							Proloculus diameter: •=0.017 mm						

644516 O - 63 (In pocket)

BOLIVINA (BOLIVINA) PACIFICA CUSHMAN, QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens