

Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
												Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
C-6 80 meters	8	5	2048	126	S.R. 0.08-1.03 O.R. 0.28-0.98	0.56 ± 0.01	0.13	0.16 ± 0.01	28.3 ± 1.78	20	0.25 0.50 0.75 1.00	15 10 5	Proloculus diameter: $\bullet = 0.017$ mm $\circ = 0.05-0.084$ mm	0.50 0.25	0.25 0.50 0.75 1.00	3 2 1	24°
C-17 82 meters	13	11	1664	237	S.R. 0-1.13 O.R. 0.17-1.04	0.52 ± 0.01	0.17	0.20 ± 0.01	38.7 ± 1.8	30	0.25 0.50 0.75 1.00	15 10 5	Proloculus diameter: $\bullet = < 0.017$ mm $\circ = 0.05-0.084$ mm x	0.50 0.25	0.25 0.50 0.75 1.00	3 2 1	23°
C-16 144 meters	8	6	8	6	O.R. 0.33-0.78	0.50				10	0.25 0.50 0.75 1.00	10 5	Proloculus diameter: $\bullet = 0.05-0.084$ mm	0.25	0.25 0.50 0.75 1.00	2 1	21°
C-8 450 meters	9	8	288	111	S.R. 0-1.15 O.R. 0.18-1.03	0.54 ± 0.02	0.17	0.21 ± 0.01	37.8 ± 2.5	20	0 0.25 0.50 0.75 1.00	15 10 5	Proloculus diameter: $\bullet = 0.017$ mm $\circ = 0.067-0.117$ mm	0.50 0.25	0 0.25 0.50 0.75 1.00	3 2 1	26°

BOLIVINA (BOLIVINA) INTERJUNCTA BICOSTATA CUSHMAN, QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens