

Sample and depth	I. Abundance			IIa. Length statistics (mm)							IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	Form measured	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
													Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
C-6 80 meters	206	164	52,736	Living	45	O.R. 0.16-0.40	0.33	0.05	0.05	17.3	6		15	Proloculus diameter: $\cdot = 0.017-0.042$ mm	0.25		3	
				Dead	164	S.R. 0.20-0.53 O.R. 0.16-0.50	0.37 ± 0.01	0.04	0.06 ± 0.003	15.6 ± 0.9	20							
C-17 82 meters	230	204	29,440	Living	53	O.R. 0.18-0.45	0.32	0.06	0.07	20.3	6		20	Proloculus diameter: $\cdot = 0.017-0.05$ mm	0.25		4	
				Dead	204	S.R. 0.23-0.55 O.R. 0.20-0.48	0.39 ± 0.002	0.04	0.04 ± 0.003	13.4 ± 0.66	30							

BOLIVINA (BOLIVINA) STRIATULA CUSHMAN, QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens