

Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
												Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
C-6 80 meters	15	13	3840	238	S.R. 0.22-0.47 O.R. 0.20-0.43	0.34 ± 0.003	0.03	0.04 ± 0.002	12.0 ± 0.55			Proloculus diameter: $\bar{x} = 0.025 - 0.033$ mm	0.25				
C-17 82 meters	17	17	2162	87	S.R. 0.22-0.46 O.R. 0.23-0.42	0.34 ± 0.004	0.03	0.04 ± 0.003	11.9 ± 0.9			Proloculus diameter: $\bar{x} = 0.017 - 0.033$ mm	0.25				
C-16 144 meters	11	8	11	8	S.R. 0.27-0.40 O.R. 0.30-0.37	0.33 ± 0.007	0.02	0.02 ± 0.005	6.1 ± 1.52			Proloculus diameter: $\bar{x} = 0.017$ mm	0.25				
C-8 450 meters	3	1	96	17	S.R. 0.15-0.45 O.R. 0.18-0.38	0.30 ± 0.01	0.04	0.05 ± 0.01	16.6 ± 2.8			Proloculus diameter: $\bar{x} = 0.017$ mm	0.25				

BOLIVINA (BOLIVINA) TONGI FILACOSTATA CUSHMAN AND MC CULLOCH, QUANTITATIVE DATA

For explanation of column headings see text

X indicates living specimens