

Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
												Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
C-6 80 meters	5	5	1280	72	S.R. 0.03-0.55 O.R. 0.15-0.55	0.29 ± 0.01	0.07	0.09 ± 0.01	29.2 ± 2.4		0.25 0.50	15 10 5 2	0.25 0.50	0.25 0.50	3 2 1	0.25 0.50	39°
C-17 82 meters	6	4	768	83	S.R. 0.04-0.55 O.R. 0.15-0.50	0.29 ± 0.01	0.07	0.08 ± 0.01	28.5 ± 2.2		0.25 0.50	15 10 5 2	0.25 0.50	0.25 0.50	3 2 1	0.25 0.50	31°
C-16 144 meters	28	24	28	24	S.R. 0.08-0.58 O.R. 0.16-0.50	0.33 ± 0.02	0.07	0.08 ± 0.01	25.6 ± 3.7		0.25 0.50	15 10 5 2	0.25 0.50	0.25 0.50	2 1	0.25 0.50	37°
C-8 450 meters	1	0	32	12	S.R. 0.10-0.45 O.R. 0.16-0.38	0.29 ± 0.02	0.05	0.06 ± 0.01	21.2 ± 4.3		0.25 0.50	15 10 5 2	0.25 0.50	0.25 0.50	2 1	0.25 0.50	47°

BOLIVINA (BOLIVINA) ACUMINATA NATLAND, QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens