

Sample and depth	I. Abundance			IIa. Length statistics (mm)						IIb. Length frequency M = mean length		Growth characteristics: variations with length				
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	III. Number of chambers		IV. Width		V. Flare	
											Length (mm)	Number of pairs of chambers	Width (mm)	Length (mm)	Growth index	Length (mm)
C-8 450 meters	4	4	128	48	S.R. 0-1.03 O.R. 0.16-0.98	0.49 ± 0.03	0.14	0.18 ± 0.02	36.9 ± 3.8							
C-14 800 meters	14	9	896	56	S.R. 0-1.15 O.R. 0.16-0.98	0.50 ± 0.03	0.20	0.22 ± 0.02	45.3 ± 4.1							
C-9 885 meters	20	20	640	20	S.R. 0-1.30 O.R. 0.13-1.18	0.46 ± 0.06	0.22	0.28 ± 0.04	57.8 ± 9.2							
C-13 1600 meters	6	5	6	5	O.R. 0.22-0.47	0.35										

BOLIVINA (BOLIVINA) PSEUDOBEURICHI CUSHMAN, QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens