

Species	I. Abundance			IIa. Length statistics (mm)							IIb. Length frequency M = mean length		Growth characteristics: variations with length						
	Total in split	Perfect in split	Total in sample	Forms measured	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	Number of pairs of chambers	III. Number of chambers		IV. Width		V. Flare	
														Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)	
<i>Bolivina (Bolivina) pacifica</i>	55	31	14,080	Total	216	S.R. 0.16-0.48 O.R. 0.16-0.47	0.33 ± 0.004	0.04	0.05 ± 0.003	16.8 ± 0.81	30	0.25 0.50 0.75 1.00 1.25	30	Proloculus diameter: $\cdot = 0.017$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	52°
<i>Bolivina (Bolivina) interjuncta bicostata</i>	8	5	2048	Total	126	S.R. 0.08-1.03 O.R. 0.28-0.98	0.56 ± 0.01	0.13	0.16 ± 0.01	28.3 ± 1.78	10	0.25 0.50 0.75 1.00 1.25	15	Proloculus diameter: $\cdot = 0.017$ mm $\cdot = 0.05-0.084$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	24°
<i>Bolivina (Bolivina) striatula</i>	206	164	52,736	Living forms	45	O.R. 0.16-0.40	0.33	0.05	0.05	17.3	4	0.25 0.50 0.75 1.00 1.25	15	Proloculus diameter: $\cdot = 0.017-0.042$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	39°
				Dead forms	164	S.R. 0.20-0.53 O.R. 0.16-0.50	0.37 ± 0.01	0.04	0.06 ± 0.003	15.6 ± 0.9	20	0.25 0.50 0.75 1.00 1.25	10	0.25 0.50 0.75 1.00	5	0.25 0.50 0.75 1.00	1	0.25 0.50 0.75 1.00	50°
<i>Bolivina (Bolivina) tongi filicostata</i>	15	13	3840	Total	238	S.R. 0.22-0.47 O.R. 0.20-0.43	0.34 ± 0.003	0.03	0.04 ± 0.002	12.0 ± 0.55	40	0.25 0.50 0.75 1.00 1.25	10	Proloculus diameter: $\cdot = 0.025-0.033$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	50°
<i>Bolivina (Bolivina) acuminata</i>	5	5	1280	Total	72	S.R. 0.03-0.55 O.R. 0.15-0.55	0.29 ± 0.01	0.07	0.09 ± 0.01	29.2 ± 2.4	10	0.25 0.50 0.75 1.00 1.25	15	Proloculus diameter: $\cdot = 0.017$ mm $\cdot = 0.033-0.067$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	39°
<i>Bolivina (Bolivina) saughani</i>	10	10	2560	Total	29	S.R. 0.12-0.22 O.R. 0.13-0.20	0.17 ± 0.003	0.01	0.02 ± 0.002	10.0 ± 1.3	10	0.25 0.50 0.75 1.00 1.25	10	Proloculus diameter: $\cdot = < 0.017$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	2	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) inflata</i>	1	1	256	Total	16	S.R. 0.12-0.28 O.R. 0.15-0.25	0.20 ± 0.01	0.03	0.03 ± 0.01	14.6 ± 2.6	10	0.25 0.50 0.75 1.00 1.25	10	Proloculus diameter: $\cdot = 0.017$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	2	0.25 0.50 0.75 1.00	
<i>Bolivina (Bolivina) bradyi</i>	1	1	256	Total	16	S.R. 0.23-0.87 O.R. 0.30-0.70	0.55 ± 0.03	0.08	0.11 ± 0.02	19.6 ± 3.4	10	0.25 0.50 0.75 1.00 1.25	10	Proloculus diameter: $\cdot = 0.017-0.067$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	38°
<i>Suggrundia eckisi</i>	11	8	2816	Total	115	S.R. 0.12-0.38 O.R. 0.15-0.40	0.25 ± 0.01	0.03	0.05 ± 0.003	18.5 ± 1.2	20	0.25 0.50 0.75 1.00 1.25	10	Proloculus diameter: $\cdot = 0.017-0.033$ mm	0.25 0.50 0.75 1.00	0.25 0.50 0.75 1.00	3	0.25 0.50 0.75 1.00	37°

SAMPLE C-6 (80 METERS), QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens