

Species	I. Abundance			IIa. Length statistics (mm)					IIb. Length frequency M = mean length		Growth characteristics: variations with length						
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare	
												Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)
<i>Bolivina (Bolivina) pacifica</i>	5	5	5	5	O.R. 0.23-0.32	0.28	0.03					Proloculus diameter: $\cdot = 0.017$ mm					58°
<i>Bolivina (Bolivina) interjuncta bicostata</i>	8	6	8	6	O.R. 0.33-0.78	0.50						Proloculus diameter: $\cdot = 0.05-0.084$ mm					21°
<i>Bolivina (Bolivina) tongi filacostata</i>	11	8	11	8	S.R. 0.27-0.40 O.R. 0.30-0.37	0.33 ± 0.007	0.02	0.02 ± 0.005	6.10 ± 1.52			Proloculus diameter: $\cdot = 0.017$ mm					52°
<i>Bolivina (Bolivina) acuminata</i>	28	24	28	24	S.R. 0.08-0.58 O.R. 0.16-0.50	0.33 ± 0.02	0.07	0.08 ± 0.01	25.6 ± 3.70			Proloculus diameter: $\cdot = 0.017$ mm $\bullet = 0.033-0.05$ mm					37°
<i>Suggrunda eckisi</i>	7	6	7	6	O.R. 0.20-0.28	0.24						Proloculus diameter: $\cdot = 0.017$ mm					

SAMPLE C-16 (144 METERS), QUANTITATIVE DATA

For explanation of column headings see text

X indicates living specimens