Species	I. Abundance: Total in split	I. Abundance: Perfect in split	I. Abundance: Total in sample	IIa. Length statistics (mm): N	IIa. Length statistics (mm): Range	IIa. Length statistics (mm): M	IIa. Length statistics (mm): MD	IIa. Length statistics (mm): σ	IIa. Length statistics (mm): V	IIb. Length frequency, M = mean length: Number of specimens / Length (mm)	III. Number of chambers: Number of pairs of chambers / Length (mm)	IV. Width: Width (mm) / Length (mm)	V. Flare: Growth index / Length (mm)
Bolivina (Bolivina) seminuda	90	57	2880	57	S.R. 0.13-0.70 O.R. 0.26-0.62	0.42 ±0.01	0.08	0.09 ±0.01	22.5 ±2.1		Proloculus diameter: · =<0.017 mm; • =0.05–0.067 mm		39°
Bolivina (Bolivina) humilis	123	110	3936	110	S.R. 0.12-0.55 O.R. 0.18-0.50	0.33 ±0.01	0.06	0.07 ±0.004	21.3 ±1.4		Proloculus diameter: · =0.017 mm; • =0.042–0.05 mm		49°
Bolivina (Bolivina) pacifica	25	11	800	50	S.R. 0.15-0.45 O.R. 0.22-0.45	0.30 ±0.01	0.04	0.05 ±0.01	16.1 ±1.6		Proloculus diameter: · =0.017 mm; • =0.025–0.05 mm		52°
Bolivina (Bolivina) interjuncta bicostata	9	8	288	111	S.R. 0-1.15 O.R. 0.18-1.03	0.54 ±0.02	0.17	0.21 ±0.01	37.8 ±2.5		Proloculus diameter: · =0.017 mm; • =0.067–0.117 mm		26°
Bolivina (Bolivina) subadvena	91	81	2912	81	S.R. 0-0.98 O.R. 0.18-0.88	0.43 ±0.02	0.15	0.19 ±0.01	42.3 ±3.3		Proloculus diameter: · =<0.017 mm; • =0.084–0.117 mm		38°
Bolivina (Bolivina) tongi filacostata	3	1	96	17	S.R. 0.15-0.45 O.R. 0.18-0.38	0.30 ±0.01	0.04	0.05 ±0.01	16.6 ±2.8		Proloculus diameter: · =0.017 mm		52°
Bolivina (Bolivina) acuminata	1	0	32	12	S.R. 0.10-0.45 O.R. 0.16-0.38	0.29 ±0.02	0.05	0.06 ±0.01	21.2 ±4.3		Proloculus diameter: · =0.017 mm; • =0.033–0.05 mm		47°
Bolivina (Bolivina) inflata	1	1	32	8	O.R. 0.20-0.28	0.23					Proloculus diameter: · =0.017 mm		
Bolivina (Bolivina) pseudobeyrichi	4	4	128	48	S.R. 0-1.03 O.R. 0.16-0.98	0.49 ±0.03	0.14	0.18 ±0.02	36.9 ±3.8		Proloculus diameter: · =0.017 mm; • =0.05 mm		21°
Bolivina Loxostomum) boltovskoyi	13	8	416	130	S.R. 0.08-1.02 O.R. 0.20-1.05	0.55 ±0.01	0.12	0.16 ±0.01	28.8 ±1.8		Proloculus diameter: · =0.017 mm; • =0.05–0.084 mm		24°
Bolivina Loxostomum) salvadorensis	4	4	128	52	S.R. 0.23-0.98 O.R. 0.26-0.83	0.61 ±0.02	0.10	0.13 ±0.01	20.7 ±2.0		Proloculus diameter: · =0.033–0.067 mm		45°
Saggrunda eckisi	6	5	192	26	S.R. 0.10-0.35 O.R. 0.16-0.35	0.23 ±0.01	0.03	0.04 ±0.01	19.0 ±2.6		Proloculus diameter: · =0.017 mm		33°

SAMPLE C–8 (450 METERS), QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens

644516 O - 63 (In pocket)