

Species	I. Abundance			IIa. Length statistics (mm)							IIb. Length frequency M = mean length		Growth characteristics: variations with length							
	Total in split	Perfect in split	Total in sample	Forms measured	N	Range	M	MD	σ	V	Number of specimens	Length (mm)	III. Number of chambers		IV. Width		V. Flare			
													Number of pairs of chambers	Length (mm)	Width (mm)	Length (mm)	Growth index	Length (mm)		
<i>Bolivina (Bolivina) seminuda</i>	84	45	2688	Total	45	S.R. 0-1.10 O.R. 0.22-0.82	0.52 ± 0.03	0.18	0.19 ± 0.01	36.8 ± 3.9										
<i>Bolivina (Bolivina) argentea</i>	45	33	1440	Total	33	S.R. 0-1.00 O.R. 0.13-1.07	0.35 ± 0.04	0.16	0.21 ± 0.03	60.1 ± 7.4										
<i>Bolivina (Bolivina) semiperforata</i>	66	49	2112	Micro-spheric	16	O.R. 0.20-0.67	0.49	0.09	0.11	22.2										
				Megalospheric	34	O.R. 0.20-0.62	0.46	0.10	0.12	25.2										
				Total (micro-spheric plus megalospheric)	50	S.R. 0.13-0.82 O.R. 0.20-0.67	0.47 ± 0.02	0.10	0.11 ± 0.01	24.3 ± 2.5										
<i>Bolivina (Bolivina) minuta</i>	52	44	1664	Total	44	S.R. 0.10-0.35 O.R. 0.15-0.32	0.23 ± 0.01	0.03	0.04 ± 0.004	17.7 ± 1.9										
<i>Bolivina (Bolivina) subadvena</i>	261	186	8352	Micro-spheric	89	O.R. 0.13-1.32	0.53	0.28	0.33	62.5										
				Megalospheric	97	O.R. 0.12-0.98	0.58	0.17	0.20	35.0										
				Total (micro-spheric plus megalospheric)	186	S.R. 0-1.37 O.R. 0.12-1.32	0.56 ± 0.02	0.23	0.27 ± 0.01	48.1 ± 2.5										
<i>Bolivina (Bolivina) pseudobeyrichi</i>	20	20	640	Total	20	S.R. 0-1.30 O.R. 0.13-1.18	0.46 ± 0.06	0.22	0.28 ± 0.04	57.8 ± 9.2										
<i>Suggrunda eekisi</i>	11	10	352	Total	10	S.R. 0.10-0.30 O.R. 0.16-0.25	0.20 ± 0.01	0.02	0.03 ± 0.01	16.6 ± 3.7										

SAMPLE C-9 (885 METERS), QUANTITATIVE DATA

For explanation of column headings see text
X indicates living specimens