

	I. Abundance			IIa. Length statistics (mm)					IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	σ	V	Number of specimens	III. Number of chambers		IV. Width		V. Flare	
											Length (mm)	Number of pairs of chambers	Width (mm)	Length (mm)	Growth index	Length (mm)
<i>Bolivina (Bolivina) pacifica</i>	8	7	8	7	O.R. 0.15-0.32	0.25				10	0.25 0.50 0.75	10	0.25 0.50 0.75	3	0.25 0.50 0.75	57°
<i>Bolivina (Bolivina) semiperforata</i>	17	12	17	12	S.R. 0-1.03 O.R. 0.22-0.77	0.45 ±0.06	0.18	0.20 ±0.04	43.3 ±8.8	10	0.25 0.50 0.75	15	0.25 0.50 0.75	3	0.25 0.50 0.75	23°
<i>Bolivina (Bolivina) subadvena</i>	5	5	5	5	O.R. 0.18-0.30	0.22				10	0.25 0.50 0.75	10	0.25 0.50 0.75	2	0.25 0.50 0.75	32°
<i>Bolivina (Bolivina) pseudobeyrichi</i>	6	5	6	5	O.R. 0.22-0.47	0.35				10	0.25 0.50 0.75	5	0.25 0.50 0.75	2	0.25 0.50 0.75	21°

SAMPLE C-13 (1,600 METERS), QUANTITATIVE DATA

For explanation of column headings see text

X indicates living specimens