

Species	I. Abundance			IIa. Length statistics (mm)					IIb. Length frequency M = mean length		Growth characteristics: variations with length					
	Total in split	Perfect in split	Total in sample	N	Range	M	MD	$\sigma$	V	Number of specimens	III. Number of chambers		IV. Width		V. Flare	
											Length (mm)	Number of pairs of chambers	Width (mm)	Length (mm)	Growth index	Length (mm)
<i>Bolivina (Bolivina) seminuda</i>	15	3	15	3	O.R. 0.33-0.67	0.52										
<i>Bolivina (Bolivina) humilis</i>	33	25	33	25	S.R. 0.08-0.48 O.R. 0.18-0.45	0.29 $\pm 0.01$	0.05	0.08 $\pm 0.01$	23.5 $\pm 3.3$							
<i>Bolivina (Bolivina) pacifica</i>	2	2	2	2	O.R. 0.28-0.32	0.30										
<i>Bolivina (Bolivina) semiperforata</i>	38	23	38	23	S.R. 0.18-1.05 O.R. 0.28-0.83	0.62 $\pm 0.03$	0.11	0.15 $\pm 0.02$	23.5 $\pm 3.4$							
<i>Bolivina (Bolivina) subadvena</i>	4	4	4	4	O.R. 0.20-0.45	0.33										
<i>Suggrunda eckisi</i>	7	6	7	6	O.R. 0.15-0.30	0.22										

SAMPLE C-10 (1,700 METERS), QUANTITATIVE DATA

For explanation of column headings see text

X indicates living specimens