

Early Jurassic Ammonites from Alaska

GEOLOGICAL SURVEY PROFESSIONAL PAPER 1148



Early Jurassic Ammonites from Alaska

By RALPH W. IMLAY

GEOLOGICAL SURVEY PROFESSIONAL PAPER 1148

Studies of Early Jurassic ammonites from Alaska provide close correlations with Lower Jurassic ammonite zones in Europe and with ammonite successions elsewhere in the world



UNITED STATES DEPARTMENT OF THE INTERIOR

JAMES G. WATT, *Secretary*

GEOLOGICAL SURVEY

Doyle G. Frederick, *Acting Director*

Library of Congress Cataloging in Publication Data

Imlay, Ralph Willard, 1908-
Early Jurassic ammonites from Alaska.

(Geological Survey professional paper ; 1148)

Bibliography: p.

Supt. of Docs. no.: I 19.16:1148

Includes index.

1. Ammonoidea.	2. Paleontology--Jurassic.	3. Paleontology--Alaska.	I. Title.	II. Series.
QE807.A51585	564'.53	81-607901		
	AACR2			

For sale by the Superintendent of Documents, U.S. Government Printing Office
Washington, D.C. 20402

CONTENTS

	Page
Abstract	1
Introduction	2
Biologic analysis	2
Biostratigraphic summary	4
Northern Alaska	4
East-central Alaska	6
Southern Alaska	7
Wrangell Mountains	7
Talkeetna Mountains	7
Chulitna River area	10
Shellabarger and Rainy Pass areas	10
Kenai Peninsula near Seldovia	10
Puale Bay-Alinchak Bay area	11
Ages and correlations	11
Hettangian ammonites	11
Sinemurian ammonites	12
Pliensbachian ammonites	17
Toarcian ammonites	18
Ammonite faunal setting	18
Geographic distribution	18
Systematic descriptions	28
References cited	43
Index	47

ILLUSTRATIONS

[Plates 1-12 follow index]

- PLATE 1. *Psiloceras*, *P.* (*Franziceras*), and *Discamphiceras*.
2. *Waehneroceras*, *Schlotheimia*, and *Badouxia*.
3. *Charmasseiceras*, *Paltechioceras*, and *Laqueoceras*.
4. *Coroniceras*, *C.* (*Paracorniceras*), *Arietites*, and *Paltechioceras*.
5. *Arctoasteroceras*, *Arnioceras*, and *Cruciloboceras*.
6. *Paracaloceras*.
7. *Cruciloboceras* and *Fanninoceras*.
8. *Tropidoceras*, *Cruciloboceras*, *Uptonia*?, and *Apoderoceras*.
9. *Uptonia*.
10. *Arietoceras*, *Prodactylioceras*, *Arietoceras*, and *Amaltheus*.
11. *Pseudolioceras*, *Dactylioceras*, *D.* (*Orthodactylites*), *Grammoceras*, *Eleganticeras*, *Harpoceras*, *Leptaleoceras*, *Protogrammoceras*, and *Fontanelliceras*.
12. *Haugia*, *Catacoeloceras*?, *Brodieia*, *Protogrammoceras*, *Pseudolioceras*, and *Phymatoceras*?

FIGURE 1. Generalized index map of Lower Jurassic ammonite localities in Alaska	Page
2-6. Detailed index maps of:	3
2. Lower Jurassic ammonite localities in the Wrangell Mountains, southern Alaska	4
3. Lower Jurassic ammonite localities in the Nelchina area of the Talkeetna Mountains and in the upper part of the Matanuska Valley, southern Alaska	5
4. Lower Jurassic ammonite localities in the Healy (A-6) quadrangle, Chulitna River area, southern Alaska	6
5. Lower Jurassic ammonite localities in the Seldovia area on the Kenai Peninsula, southern Alaska	7
6. Lower Jurassic ammonite localities in the Puale Bay-Alinchak Bay area, Alaska Peninsula	8

	Page
FIGURE 7. Diagram of European ranges of Early Jurassic ammonite genera present in Alaska	9
8. Diagram correlating Early Jurassic ammonite faunas in northern and east-central Alaska	10
9. Diagram correlating Early Jurassic ammonite faunas in southern Alaska	11
10-12. Diagrams showing occurrences and ages of:	
10. Ammonites present in the Lower Jurassic beds exposed in the Wrangell Mountains in southern Alaska	12
11. Early Jurassic ammonites present in the Nelchina area of the Talkeetna Mountains and in the upper part of the Matanuska River Valley, southern Alaska	13
12. Ammonites present in unnamed Lower Jurassic beds on northeast side of Puale Bay on the Alaska Peninsula	14
13. Diagram correlating Lower Jurassic rocks in northern and east-central Alaska	15
14. Diagram correlating Lower Jurassic rocks in southern Alaska	16

TABLES

	Page
TABLE 1. Ammonite genera and subgenera of Early Jurassic age in Alaska	2
2. Early Jurassic ammonites from well cores in northern Alaska	17
3-6. Geographic distribution of:	
3. Early Jurassic ammonites from outcrops in northern and east-central Alaska	19
4. Early Jurassic ammonites in the Yakutat area and in the Wrangell Mountains in southern Alaska	20
5. Early Jurassic ammonites in the Talkeetna Mountains in southern Alaska	21
6. Early Jurassic ammonites in the Chulitna River area of the Alaska Range, in the Seldovia area of the Kenai Peninsula, and in the Puale Bay-Alinchak Bay and Wide Bay areas of the Alaska Peninsula	22
7. Descriptions of Lower Jurassic ammonite localities in Alaska	24

EARLY JURASSIC AMMONITES FROM ALASKA

By RALPH W. IMLAY

ABSTRACT

Early Jurassic ammonites of early Hettangian to late Toarcian Age are present in the lower part of the Kingak Shale and equivalent beds in northern Alaska. Well cores from the Arctic Coastal Plain contain ammonites of early Hettangian to Sinemurian Age and of late Pliensbachian to early Toarcian Age. Surface exposures have furnished ammonites of early Pliensbachian to late Toarcian Age. In addition the presence of the pelecypod *Otapiria tailleuri* Imlay near the base of the Lower Jurassic exposures throughout much of northern Alaska is good evidence of a Sinemurian Age because this pelecypod occurs in the subsurface with ammonites of early Sinemurian Age and it occurs on the surface near, but not on the same slabs as, *Uptonia* of earliest Pliensbachian Age.

Ammonites of early Hettangian and late Pliensbachian Age are present in part of the Glenn Shale exposed near Old Rampart in east-central Alaska.

In the Wrangell Mountains (lat 62° N., long 141°–143° W.), the upper member of the McCarthy Formation has furnished ammonites of early Hettangian to early Sinemurian Age and of latest Sinemurian to fairly early Pliensbachian Ages. The overlying Lubbe Creek Formation contains ammonites of middle to late Pliensbachian Age and possibly also of early Toarcian Age.

In the Talkeetna Mountains, northeast of Anchorage, the exposed upper part of the Talkeetna Formation has furnished ammonites of latest Sinemurian to earliest Pliensbachian Age and of late Pliensbachian to late Toarcian Age.

In the Chulitna River area, only ammonites of early and latest Sinemurian Age have been found. In the Seldovia area, only ammonites of earliest Hettangian and early Sinemurian Age have been found. In the Puale Bay area, the Hettangian and lower Sinemurian are well represented by ammonites in an unnamed formation. Sharply above lies the Kialagvik Formation, which is mostly of Bajocian Age, but which contains the ammonite *Haugia* of middle to early late Toarcian Age about 30 ft (9 m) above its base.

These data show that parts of all these Lower Jurassic sequences are not represented by ammonites or are poorly represented. Thus ammonites of late Hettangian and middle Pliensbachian Age are represented only by single occurrences, and the Sinemurian zones of *Caenisites turneri* to *Oxynoticeras oxynotum* are not represented by any ammonite occurrences.

The Hettangian ammonite succession in Alaska is similar to that in western Europe and probably includes species in common. Its lower part, equivalent to the *Psiloceras planorbis* zone, is represented in many areas by *P.* cf. *P. planorbis* (J. de C. Sowerby), or by *P. (Franziceras)* cf. *P. (F.) ruidum* (Buckman), or by both. Its middle part, probably equivalent to the *Alsatites lasiacus* zone, is represented in the Wrangell Mountains by *Discamphiceras* cf. *D. toxophorum* (Gümbel), at Wide Bay (lat 57°45' N., long 156° W.) by *Waehneroceras* cf. *W. portlocki* (Wright), and at the Puale Bay-Alinchak Bay area by these taxa plus *W.* cf. *W. tenerum* (Neumayr) and *Laqueoceras* cf. *L. sublaqueus* (Waehner). The upper Hettangian is represented by a single specimen of *Schlotheimia* in the Puale Bay area.

The Sinemurian ammonite succession in Alaska is likewise similar to that in western Europe but does not include genera that are definitive for nearly half of the stage (*Caenisites turneri*-*Oxynoticeras oxynotum* zones). Its lowest part, equivalent to the *Arietites bucklandi* zone of Europe, is represented in the subsurface of the Arctic Coastal Plain by *Arietites* cf. *A. bucklandi* (Sowerby), by *Charmasseiceras*, and by species of *Coroniceras* both below and above *Charmasseiceras*. Its lowest part in southern Alaska is represented in the Chulitna River, Seldovia, and Puale Bay areas by *Paracaloceras rursicostatum* Frebold, which in the Chulitna area is associated with *Badouxia canadensis* (Frebold).

The overlying beds in most of these areas are characterized by *Arnioceras*, whose presence and stratigraphic position are evidence for correlation with the late early Sinemurian *Arnioceras semicostatum* zone of Europe. Such an age for *Arnioceras* in the Arctic Coastal Plain is confirmed by its position below that of *Coroniceras*, a genus not known above the *A. semicostatum* zone. Such an age for *Coroniceras* (*Paracaroniceras*) in the Seldovia area is shown by the known range of the subgenus in Europe.

The highest part of the Sinemurian, representing the *Echioceras raricostatum* zone, has been identified by the occurrence of *Paltechioceras* in the Wrangell and Talkeetna Mountains and in the Chulitna River area. That genus in the Talkeetna Mountains is associated with *Cruciloboceras*, which ranges upward into the lower Pliensbachian. In the Chulitna River area *Paltechioceras* is associated with *Arctoasteroceras*, a genus that in arctic Canada occurs with *Oxynoticeras oxynotum* (Quenstedt).

The Pliensbachian ammonite succession in northern Alaska is incomplete. At its base, the *Uptonia jamesoni* zone is represented by *Uptonia* cf. *U. jamesoni* (J. de C. Sowerby) obtained from outcrops near the Iknavik and Etivluk Rivers (tributaries of the Colville River). The next two higher zones of western Europe have not been recognized. The still higher zone of *Amaltheus margaritatus* is present throughout northern Alaska as well as farther southeast in the Old Rampart area of east-central Alaska. The highest zone of *Pleuroceras spinatum* is represented by *Amaltheus (Pseudoamaltheus) engelhardti* in the subsurface of the Arctic Coastal Plain. These occurrences of *Amaltheus* in abundance at many places furnish exact correlations with the upper Pliensbachian beds of northern Eurasia.

Pliensbachian ammonite successions in southern Alaska have been identified only in the Wrangell and Talkeetna Mountains and questionably in the Yakutat district. The *Uptonia jamesoni* zone is represented in the Wrangell Mountains by *Uptonia* cf. *U. dayiceratoides* Mouterde, in the Talkeetna Mountains by *Apoderoceras*, and probably in the Yakutat district by a fragment that resembles the outer whorl of *Uptonia* rather than *Cruciloboceras*. The next higher zone of *Tragophylloceras ibex*, is represented by *Tropidoceras actaeon* (d'Orbigny) in association with *Cruciloboceras*. The *Prodactylioceras davoei* zone is possibly represented in the Wrangell Mountains by some float specimens of *Prodactylioceras* that were collected along with *Uptonia*. The highest two zones of the Pliensbachian are represented by the ammonites *Protogrammoceras*, *Leptaleoceras*, *Arietoceras*, and *Fontannelluceras*, which are identical

or nearly identical with species in the Mediterranean area. With these taxa at one locality in the Talkeetna Mountains were found two specimens of *Amaltheus*.

This association with *Amaltheus* furnishes a correlation with the upper Pliensbachian of northern Alaska. Otherwise the late Pliensbachian ammonites in southern Alaska have nothing in common generically with ammonites of that age in the Arctic region. The same genera do occur farther south, however, in western British Columbia and in eastern Oregon, as well as in the Mediterranean area.

The Toarcian ammonite succession in Alaska is similar to that in western Europe and probably includes species in common. Characteristic taxa from the base upward include (1) *Dactylioceras* and *Catacoeloceras* (?), (2) *Harpoceras* cf. *H. exaratum* (Young and Bird) and *Eleganticeras*, (3) *Haugia*, *Brodieia*, and *Pseudolioceras*, and (4) *Pseudolioceras* and *Grammoceras*. The most complete Toarcian ammonite sequence is in the Talkeetna Mountains.

INTRODUCTION

Early Jurassic ammonites obtained from eight areas in northern, eastern, and southern Alaska (figs. 1-6) are described herein in order to present all available evidence concerning the stratigraphic and geographic distribution of the taxa (figs. 7-9), to evaluate their faunal setting in relation to other parts of the world, to make regional and continental correlations, and to date the formations in Alaska as precisely as possible in terms of the standard Jurassic zones of western Europe (figs. 10-14).

This study is based mainly on fossil collections and stratigraphic data furnished by many U.S. Geological Survey geologists, as listed in Imlay and Detterman (1973, p. 8, 9). It is also based on excellent fossil collections and data obtained by British Petroleum (Alaska), Inc., in 1964 in northern Alaska and by the Richfield Oil Co. (now Atlantic Richfield Oil Co.) in 1962 on the east shore of Puale Bay in the Alaska Peninsula. Biostratigraphic data for northern Alaska were furnished by W. P. Brosgé, R. L. Detterman, and H. N. Reiser; for east-central Alaska, by E. M. MacKevett, Jr.; for the Talkeetna Mountains, by Arthur Grantz; and for the Chulitna Valley, by D. L. Jones.

The described specimens are deposited in the type collections of the U.S. National Museum and are labeled USNM.

BIOLOGIC ANALYSIS

Alaska ammonites of Early Jurassic (Hettangian to Toarcian) age described or mentioned herein number about 500 specimens whose distribution by family, subfamily, genus, and subgenus is shown on table 1. This table shows that the families present include the Arietitidae (15 percent), Polymorphitidae (20 percent), Schlotheimiidae (16 percent), Hildoceratidae (15 percent), Psiloceratidae (9 percent), Hammoceratidae (6 percent), and Eoderoceratidae (6 percent). The remaining families combined represent only 13 percent of the total number of specimens.

The classification shown in table 1 is essentially that proposed by Donovan and Forsey (1973 p. 2-4), except for *Fanninoceras* and *Arctoasteroceras*, which those writers consider to be synonyms of *Radstockiceras* and *Aegasteroceras*, respectively. Nonetheless, *Fanninoceras* (McLearn, 1930, p. 4, 5; 1932, p. 76-80) differs morphologically from *Radstockiceras* by having a thinner whorl section and an undercut umbilical wall (Frebold, 1967a, p. 1146). It differs timewise by occurring in beds of late Pliensbachian Age in eastern Oregon (Imlay, 1968, p. C10, C11) and in beds of earliest to latest Pliensbachian Age in the Queen Charlotte Islands (Hans Frebold, written commun., 1975), whereas *Radstockiceras* in northwest Europe is characteristic of

TABLE 1.—Ammonite genera and subgenera of Early Jurassic age in Alaska

Family	Subfamily	Genus and subgenus	Number of specimens
Lytoceratidae		<i>Lytoceras</i>	1
Psiloceratidae		<i>Psiloceras</i>	10
		(<i>Franziceras</i>)	14
		<i>Discamphiceras</i>	19
		<i>Laqueoceras</i>	1
Schlotheimiidae		<i>Schlotheimia</i>	4
		<i>Waehneroceras</i>	63
		<i>Badouxia</i>	4
		<i>Charmasseiceras</i>	8
Arietitidae	Arietitinae	<i>Arietites</i>	1
		<i>Coroniceras</i>	17
		(<i>Paracoriceras</i>)	3
		<i>Arnioceras</i>	45
		<i>Arnioceras?</i>	1
		<i>Paracaloceras</i>	10
	Asteroceratinae	<i>Arctoasteroceras</i>	2
Echioceratidae		<i>Paltechioceras</i>	10
		(<i>Orthechioceras</i>)	1
Eoderoceratidae	Xiphoceratinae	<i>Cruciloboceras</i>	26
	Coeloceratinae	<i>Apoderoceras</i>	2
		<i>Apoderoceras?</i>	1
Oxynoticeratidae		<i>Fanninoceras</i>	16
Polymorphitidae		<i>Uptonia</i>	81
		<i>Uptonia?</i>	9
		<i>Tropidoceras</i>	4
Amaltheidae		<i>Amaltheus</i>	18
		(<i>Pseudo-amaltheus</i>)	1
Dactylioceratidae		<i>Prodactylioceras</i>	4
		<i>Dactylioceras</i>	6
		(<i>Orthodactylites</i>)	7
		<i>Catacoeloceras?</i>	1
Hildoceratidae	Arieticeratinae	<i>Arietoceras</i>	42
		<i>Leptaleoceras</i>	1
		<i>Fontanelliceras</i>	5
	Harpoceratinae	<i>Harpoceras</i>	5
		<i>Protogrammoceras</i>	5
		<i>Elegantuliceras</i>	12
		<i>Pseudoloboceras</i>	5
	Grammocerotinae	<i>Grammoceras?</i>	1
Hammatoceratidae	Phymatoceratinae	<i>Phymatoceras?</i>	3
		<i>Brodieia</i>	1
		<i>Haugia</i>	28



FIGURE 1.—Generalized index map of Lower Jurassic ammonite areas in Alaska, as listed in tables 3–7.

the earliest Pliensbachian (Dean and others, 1961, p. 463, fig. 7).

The genus *Arctoasteroceras* of Frebold (1960, p. 13, 14, pl. 2 figs. 1–5, pl. 3, figs. 1–3) differs morphologically from *Aegasteroceras* of Spath (1925, p. 265, 267, fig. 6a) by having weak ribs on the upper parts of the flanks and on the venter and commonly by having an asymmetrical suture line. These differences were not accepted as valid by Hallam (1965, p. 1495) or by Donovan and Forsey (1973, p. 3). Nonetheless, some European specimens of *Aegasteroceras*, including the holotype of the genotype species, *A. simile* Spath (Guerin-Franiatte, 1966, p. 310–313, pls. 189–192), bear strong ribs on the upper

part of the flanks and on the margins of the venter, and some ribs are weakly connected across the venter. This strong ribbing contrasts markedly with the weak ribs on *Arctoasteroceras jeletzkyi* Frebold from Canada.

These genera likewise differ slightly in age. Thus, *Aegasteroceras* in Europe occurs in the zone of *Asteroceras obtusum* of earliest late Sinemurian Age (Dain and others, 1961, p. 454). *Arctoasteroceras jeletzkyi* in Canada is dated as middle late Sinemurian because of its occurrence with *Oxynotoceras oxynotum* and below echioceratid ammonites (Frebold, 1960, p. 14, 26). The Alaskan specimens of *Arctoasteroceras* described herein are dated as probably latest Sinemurian

EARLY JURASSIC AMMONITES FROM ALASKA




FIGURE 2.—Detailed index map of Lower Jurassic ammonite localities in the Wrangell Mountains, southern Alaska.

because they were collected with *Paltechioceras* (*Orthechioceras*?), which in Europe occurs only in the *Echioceras raricostatum* zone (Getty, 1973, p. 6).

The characteristics of the Early Jurassic ammonites described herein agree very well with those of various genera and subgenera defined in the "Treatise on Invertebrate Paleontology" (Arkell and others, 1957). In addition, the taxonomy of the Psiloceratidae has been discussed by Donovan (1952, p. 634, 641) and Frebold (1967b, p. 17); the Schlotheimiidae by Lange (1951, p. 23, 25), Donovan (1952, p. 644-655), Frebold (1960, p. 13), and Guex and Taylor (1976, p. 525); the Arietidae by Donovan (1952, p. 717, 725, 739, 746), Donovan and Forsey (1973, p. 6, 7), and Guerin-Franiatte (1966, p. 106-118, 150, 252, 281, 283, 309, 310, 313); the Echioceratidae by Getty (1973, p. 7, 8, 17-21, 23, 24); the Amaltheidae by Howarth (1958, pt. 1, p. 1, 21; pt. 2); the Polymorphitidae by Bremer (1965, p. 177); the Dactylioceratidae by Pinna and Levi-Setti (1971) and

Schmidt-Effing (1972); the Polymorphitidae, Dactylioceratidae, and Hildoceratidae by Geczy (1976); and the Dactylioceratidae by Fischer (1966, p. 20-48). In addition, the taxonomy of the Early Jurassic ammonite families of southern Germany has been briefly described by Schlegelmilch (1976, p. 32-95).

BIOSTRATIGRAPHIC SUMMARY

NORTHERN ALASKA

Lower Jurassic marine beds cropping out in northern Alaska (figs. 1, 13) have been briefly described or mentioned by Imlay and Detterman (1973, p. 9; p. 12 fig. 11B). Those from the Sagavanirktok River eastward have been described in considerable detail by Detterman and others (1975, p. 18-20). In general the outcrops are poorly exposed and sparsely fossiliferous, but they have furnished fossils representing most of the Pliensbachian and Toarcian Stages and perhaps the Sinemurian (fig. 8). Field studies to 1978 are insufficient to demonstrate




FIGURE 3.—Detailed index map of Lower Jurassic ammonite localities in the Nelchina area of the Talkeetna Mountains and in the upper part of the Matanuska Valley, southern Alaska.

whether the absence of earlier Jurassic is due to lack of deposition, poor outcrops, or insufficient field work in poorly fossiliferous beds.

The westernmost exposures in the DeLong Mountains consist mostly of clay shales that bear limestone concretions, are of unknown thickness, and have furnished ammonites of late Pliensbachian to early Toarcian Age. Next to the east between the Ipnayik and Itkillik Rivers, the Lower Jurassic apparently consists of 60 feet (18 m) or less of siliceous claystone that locally contains some limestone and is dated as Sinemurian to early Pliensbachian. Such an age is shown by the widespread distribution of *Otapiria tailleuri* Imlay, which in the subsurface is found associated only with ammonites of early Sinemurian Age, and by the presence on the surface of the early Pliensbachian ammonite *Uptonia* in slabs that do not contain *Otapiria*, although that genus occurs nearby.

Still farther east, in the area between the

Sagavanirktok and Aichilik Rivers, the Lower Jurassic beds (Detterman and others, 1975, p. 18–20, 44) represent the lower part of the Kingak Shale and range in thickness from 100 feet (30 m) or less, to about 900 feet (274 m). This lower part consists of fissile black papery shale that at the Aichilik River is about 600 feet (180 m) thick and on the Kavik River contains *Otapiria tailleuri* Imlay of Sinemurian Age. The upper part consists of dark-gray clay shale and claystone that in places is as much as 300 feet (92 m) thick and contains ammonites of late Pliensbachian to late Toarcian Age.

The Lower Jurassic sequences in the subsurface of northern Alaska differ from those on the surface mainly by containing minor amounts of glauconitic sandstone, by being somewhat more fossiliferous, and by containing ammonites representing all Lower Jurassic stages (table 2). They have not, however, furnished any ammonites representing middle Sinemurian to middle Pliensbachian time, or late Toarcian time.




FIGURE 4.—Detailed index map of Lower Jurassic ammonite localities in the Healy (A-6) quadrangle, Chulitna River area, southern Alaska.

EAST-CENTRAL ALASKA

In the northern part of east-central Alaska, the Lower Jurassic is poorly exposed but is reported to consist of about 1000 feet (305 m) of sandstone, siltstone, shale, and quartzite that are part of the Glenn Shale (Brosgé

and Reiser 1964, 1969; Imlay and Detterman, 1973, p. 13, 14). The basal 120 feet (37 m) exposed on Spike Mountain about 30 miles (48 km) north of Old Rampart, consists of sandy beds, rests on granite, and contains the Hettangian ammonites *Psiloceras* (*Psiloceras*) and




FIGURE 5.—Detailed index map of Lower Jurassic ammonite localities in the Seldovia area (Seldovia (B-5) quadrangle) on the Kenai Peninsula, southern Alaska.

P. (Franziceras) throughout. The only other Jurassic ammonite from the same general area is a specimen of *Amaltheus stokesi* (J. Sowerby) obtained about 9 miles (14.4 km) east-southeast of Old Rampart.

SOUTHERN ALASKA

WRANGELL MOUNTAINS

The Lower Jurassic sequence in the Wrangell Mountains has been described in detail by MacKevett (1969, 1970, 1971); has been mentioned briefly by Imlay and Detterman (1973, p. 8, 11) regarding its position relative to a Jurassic volcanic island-arc system; and is depicted herein (figs. 10, 14) in regard to fossil occurrences and ages. Most of the sequence is represented by the upper member of the McCarthy Formation which consists of

2,000–2,500 feet (610–760 m) of very thin to medium-bedded, dark-gray laminated chert and spiculite, grades into adjoining lithologic units, and is of Hettangian, Sinemurian, and earliest Pliensbachian Age. At the top, of the sequence is the Lubbe Creek Formation, which consists of 100–300 feet (30–91 m) of medium-gray spiculite and chert, is overlain unconformably by beds of Bajocian Age, and is of middle and late Pliensbachian to possibly early Toarcian Age.

TALKEETNA MOUNTAINS

The Lower Jurassic sequences in the Nelchina area of the Talkeetna Mountains and in the adjoining upper part of the Matanuska River valley (figs. 11, 14) have been studied by Arthur Grantz. Most of the lithologic and




FIGURE 6.—Detailed index map of Lower Jurassic ammonite localities in the Puale Bay-Alinchak Bay area, Alaska Peninsula.

stratigraphic data shown in figure 11 were prepared by Grantz (written commun., 1977), along with these comments:

The Talkeetna Formation comprises a diverse assemblage of volcanic arc deposits. Waterlain pyroclastic and tuffaceous sedimentary rocks are much dominant over lava flows. If all rocks mapped as Talkeetna Formation are correctly assigned, its composition ranges from basaltic to rhyolitic, with andesitic rocks apparently dominant. The Talkeetna volcanic breccias and tuffs were commonly, and perhaps dominantly, deposited in marine waters. Marine fossils are widespread, and locally abundant in the waterlaid tuffs and in associated more-or-less tuffaceous sands and lutites. However, the presence of thick intervals of coarse volcanic breccias, tuffs and some flows that are barren of fossils and some outcrops that are rich in plant fossils suggest that nonmarine facies may also constitute an important part of the formation. Presumably, the fossiliferous marine beds represent inter-volcanic vent or arc-distal facies, and the pyroclastic and plant-rich beds represent submarine and subaerial composite volcanic cones that formed at major volcanic vents.

The Talkeetna Formation in the upper Matanuska Valley and Nelchina area occurs in large but well-separated outcrop areas. The lack of mapping continuity combined with moderate to locally great structural complexity, locally intense plutonism, and some metamorphism makes the recognition of the stratigraphic succession in these rocks difficult. The chief problem is the possibility, indeed probability, that the local lithologic succession in each isolated outcrop area of the formation owes as much, or more, to its geographic position (i.e., facies) as to its biostratigraphic position. However, if the prior assumption is made that the local successions do have regional validity, then [a] * * * gross, and incomplete, succession can be inferred. Even with this assumption, however, the reconstruction given does not represent the only possible explanation of the available facts.

On the basis of these data and inferences, the Talkeetna Formation of the Talkeetna Mountains is 15,000–19,000 feet (4,660–5,790 m) thick, is incompletely exposed basally, is dominated by volcanic rocks, is mostly marine, and contains ammonites of latest




FIGURE 7.—European ranges of Early Jurassic ammonite genera present in Alaska.

Series	Stage	Northwest European Ammonite zones (After Dean, Donovan, and Howarth, 1961)	Arctic Coastal Plain in subsurface of Point Barrow-Cape Simpson area	DeLong Mountains area near Thetis Creek and Epewik River northwestern Alaska	Ignavik-Anaktuvuk Rivers area north-central Alaska	Ignek Valley in Mt. Michelson (C-4) quadrangle to Aichilik River in northeastern Alaska	Old Rampart area in east-central Alaska
Jurassic	Toarcian	<i>Dumortieria levesquei</i>				<i>Psuedolioceras</i>	
		<i>Grammoceras thouarsense</i>					
		<i>Haugia variabilis</i>					
		<i>Hildoceras bifrons</i>		<i>Harpoceras</i> cf. <i>H. exaratum</i> and <i>Elegantoceras</i>		<i>Harpoceras</i> cf. <i>H. exaratum</i>	
		<i>Harpoceras falcifer</i>	<i>Dactylioceras</i> (Orthodactylites) and <i>Catacoeloceras</i> (?)	<i>Dactylioceras</i> (Orthodactylites) cf. <i>D. (O.) directum</i>			
		<i>Dactylioceras tenuicostatum</i>					
	Plensbachian	<i>Pleuroceras spinatum</i>	<i>Amaltheus engelhardti</i>				
		<i>Amaltheus margaritatus</i>	<i>A. margaritatus</i> <i>A. stokesi</i>	<i>Amaltheus margaritatus</i>		<i>Amaltheus stokesi</i>	<i>Amaltheus stokesi</i>
		<i>Prodactylioceras davoei</i>					
		<i>Tragophylloceras ibex</i>					
		<i>Uptonia jamesoni</i>			<i>Uptonia</i> cf. <i>U. jamesoni</i>		
	Sinemurian	<i>Echioceras raricostatum</i>					
		<i>Oxynoticeras oxynotum</i>					
		<i>Asteroceras obtusum</i>					
		<i>Caenisites turneri</i>					
		<i>Arnioceras semicostatum</i>	<i>Arnioceras</i> and <i>Coroniceras</i>				
		<i>Arietites bucklandi</i>	<i>Arietites</i> , <i>Coroniceras</i> , and <i>Charmasseiceras</i>				
	Hettangian	<i>Schlotheimia angulata</i>					
		<i>Alsatites liasicus</i>	<i>Waehneroceras</i> ?				
		<i>Psiloceras planorbis</i>	<i>Psiloceras</i> (<i>Franziceras</i>)				<i>Psiloceras</i> (<i>Franziceras</i>)

FIGURE 8.—Correlation of Early Jurassic ammonite faunas in northern and east-central Alaska.

Sinemurian to late Toarcian Age (fig. 11). Other studies show (Grantz and others, 1963) that the Talkeetna Formation is overlain unconformably by beds of early Bajocian Age, that the unconformity is dated paleontologically as of latest Toarcian and earliest Bajocian Age, and that the formation is intruded by granitic rocks that may be of the same age as the unconformity.

CHULITNA RIVER AREA

The Lower Jurassic sequence exposed in the Healy (A-6) quadrangle in the south-central part of the Alaska Range has not been described. It has furnished ammonites of early Sinemurian Age at six localities and of latest Sinemurian Age at one locality (fig. 4). Most of the ammonites of early Sinemurian Age are preserved in a matrix of gray calcareous sandstone or of silty phosphatic limestone. The ammonites of latest Sinemurian Age (USGS Mesozoic loc. 31261) are preserved in a unit of massive tuffaceous argillite that apparently underlies, or is in fault contact with, siliceous argillite of Late Jurassic age (Jones and others, 1980).

SHELLABARGER AND RAINY PASS AREAS

The Lower Jurassic sequences exposed in the Talkeetna (C-6) and (B-6) quadrangles in the south-central part of the Alaska Range have not been described. In the Rainy Pass area of the Talkeetna (B-6) quadrangle, Early Jurassic fossils were obtained from gray, calcareous, fine-grained sandstone. In the Shellabarger Pass area of the Talkeetna (C-6) quadrangle, such fossils were obtained from brown siltstone and sandstone in T.28 N., R. 19 W.

KENAI PENINSULA NEAR SELDOVIA

Lower Jurassic beds are exposed on the shore of the Kenai Peninsula for several miles southwestward from Seldovia Bay (figs. 5, 14) in the northwestern part of the Seldovia (B-15) quadrangle. They consist of interbedded tuff, agglomerate, sandstone, shale, and limestone that probably are 2,000–3,000 feet (610–915 m) thick, according to Martin (1926, p. 135). The same sequence, based on unpublished field studies by Don Miller and the author for the U.S. Geological Survey in 1948, includes a thick middle unit consisting of unfossiliferous red lava,

Series	Stage	Northwest European Ammonite zones (Modified from Dean, Donovan and Howarth, 1961)	Wrangell Mountains in southern Alaska	Talkeetna Mountains in southern Alaska	Chulitna River area in Healy (A-6) quad. south-central part of Alaska Range	Seldovia area on the Kenai Peninsula	Puale Bay area on the Alaska Peninsula
Lower Jurassic	Toarcian	<i>Dumortieria levesquei</i>					
		<i>Grammoceras thoursense</i>		<i>Grammoceras</i>			
		<i>Haugia variabilis</i>		<i>Pseudolioceras, Haugia, Brodieia and Phymatoceras (?)</i>			<i>Haugia and Pseudolioceras</i>
		<i>Hildoceras bifrons</i>	?	<i>Harpoceras cf. H. exaratum</i>			
		<i>Harpoceras falcifer</i>	<i>Peronoceras ? Weyla dufrenoyi and W. alata</i>	<i>Dactylioceras cf. D. commune and D. (Orthodactylites) kanense</i>			
		<i>Dactylioceras tenuicostatum</i>					
	Plensbachian	<i>Pleuroceras spinatum</i>	<i>Arietoceras cf. A. domarense and Prodactylioceras (?)</i>	<i>Protogrammoceras, Amaltheus, Arietoceras, Fontanelliceras and Leptaeoceras</i>			
		<i>Amaltheus margaritatus</i>					
		<i>Prodactylioceras davoei</i>	<i>Prodactylioceras italicum</i>				
		<i>Tragophylloceras ibex</i>	<i>Tropidoceras</i>				
		<i>Uptonia jamesoni</i>	<i>Uptonia</i>	<i>Uptonia?</i> <i>Apoderoceras</i>			
	Sinemurian	<i>Echioceras raricostatum</i>	<i>Paltechioceras</i>	<i>Paltechioceras and Cruciloboceras</i>	<i>Paltechioceras and Arctoasteroceras</i>		
		<i>Oxyntoceras oxynotum</i>					
		<i>Asteroceras obtusum</i>					
		<i>Caenites turneri</i>					
		<i>Arnioceras semicostatum</i>	<i>Arnioceras</i>		<i>Arnioceras</i>	<i>Coroniceras (Paracoriceras)</i>	<i>Arnioceras</i>
		<i>Arietites bucklandi</i>	<i>Coroniceras? and arietitid ammonites</i>		<i>Paracaloceras and Badouxia</i>	<i>Paracaloceras rursicostatum</i>	<i>Paracaloceras rursicostatum</i>
	Hettangian	<i>Schlotheimia angulata</i>					<i>Schlotheimia</i>
		<i>Alsatites liasicus</i>	<i>Discamphiceras</i>				<i>Waehneroceras, Discamphiceras, and Lagueoceras</i>
		<i>Psiloceras planorbis</i>	<i>Psiloceras cf. P. planorbis</i>			<i>Psiloceras (Franziceras)</i>	<i>Psiloceras cf. P. planorbis</i>

FIGURE 9.—Correlation of Early Jurassic ammonite faunas in southern Alaska.

green lava, tuff, and agglomerate overlain and underlain by units consisting of green sandstone, tuffaceous sandstone, tuff, agglomerate, and gray limestone. Fossils occur in limestone, tuff, and sandstone beds in both the lower and upper units. The ammonites present are of Hettangian and early Sinemurian Age.

This Lower Jurassic sequence, on the basis of recent studies by Kelley (1978, p. 2356), is about 4,000 m thick and consists mainly of pyroclastic breccias and tuff. It is divisible into three mappable lithofacies, from bottom to top: (1) a "graded pyroclastic lithofacies" deposited below wave base; (2) a "pyroclastic debris flow and turbidite lithofacies" deposited below wave base; and (3) a "reworked volcanoclastic lithofacies" marked basally by fairly thick coal beds but deposited partly under tidal influences. Kelley noted that fossils occur in all three units.

PUALE BAY-ALINCHAK BAY AREA

The Lower Jurassic sequence in the Puale Bay area, as described by Imlay and Detterman (1977), is at least 1,850 feet (550 m) thick. At its base is a limestone unit about 780 feet (238 m) thick that in its lower part contains ammonites of Hettangian Age and that rests con-

formably on limestone of latest Triassic age. The overlying 1,040 feet (317 m) of beds consists mostly of massive tuffaceous conglomeratic sandstone but includes some siltstone and limestone. These beds contain ammonites of early Sinemurian Age. This sandstone is overlain sharply by sandy siltstone of the Kialagvik Formation, which is mostly of early to middle Bajocian Age. Its lowest beds, however, are of middle to early late Toarcian Age, as shown by the presence of *Haugia* about 30 feet (9 m) above its base. The Toarcian probably also includes some of the overlying 320–370 feet (98–113 m) of siltstone between this occurrence of *Haugia* and the lowest occurrence of *Pseudolioceras whiteavesi* (White) of late early Bajocian Age (figs. 9, 12). The sharp contact at the base of the Kialagvik Formation probably represents a fault rather than an unconformity.

AGES AND CORRELATIONS

HETTANGIAN AMMONITES

Hettangian ammonites have been found in the subsurface of northern Alaska, in the Old Rampart area of

Lower Jurassic formation	Lithologic characteristics and thicknesses	Fossiliferous areas in McCarthy C4-C6 and B-4 quadrangles	U.S. Geological Survey Mesozoic fossil localities	Ammonite genera in McCarthy C4-C6 and B-4 quadrangles	Stage
Lubbe Creek Formation	100-300 ft (30.5-91.5 m) Mostly medium gray, impure spiculite chert. A few beds of bioclastic coquina near top	? C-5 West of West Fork ? C-5 East of McCarthy Creek	? 28678 28531	? <i>Peronoceras</i> ? <i>Arietoceras</i>	Lower Toarcian Pliensbachian
Upper Member of McCarthy Formation	2000-2500 ft (610-762 m) Very thin-to medium-bedded, dark-gray, finely laminated organic chert and spiculite. Includes small amounts of fissile, calcareous to carbonaceous shale and limestone. Shale becomes more common in upper part	C-5 West of Nizina River C-5 West of McCarthy Creek C-4 Chitistone Mountain C-6 Ridge west of Root Glacier C-6 Ridge west of Root Glacier C-5 Ridge between Lubbe Creek and Diamond Creek ? C-5 East of McCarthy Creek	28688 28534 28671-73, 28675 31175 31174 ? 28690 14030	<i>Arietoceras</i> <i>Tropidoceras</i> and <i>Cruciloboceras</i> <i>Uptonia</i> and <i>Prodactyloceras</i> <i>Paltehioceras</i> and <i>Cruciloboceras</i> <i>Paltehioceras</i> ? <i>Cruciloboceras</i> <i>Cruciloboceras</i>	Upper
	Contacts are gradational	C-5 West of West Fork C-5 Ridge between McCarthy Creek and West Fork C-5 Ridge west of McCarthy Creek	28535 28538 28537 ? 28533	<i>Amioceras</i> <i>Arnioceras</i> Arietid ammonites ? <i>Coroniceras</i> ?	Sinemurian Lower
		? B-4 Chitistone Mountain ? B-4 Chitistone Mountain	? 29890 ? 29891	<i>Discamphiceras</i> and <i>Alsatites</i> ? ? <i>Psiloceras</i>	Hettangian

FIGURE 10.—Occurrences and ages of ammonites present in the Lower Jurassic beds exposed in the Wrangell Mountains, southern Alaska.

northern east-central Alaska, in the Wrangell Mountains of southern Alaska, in the Seldovia area of the Kenai Peninsula and in the Puale Bay-Alinchak Bay and Wide Bay areas of the Alaska Peninsula (figs. 8-10, 12).

The earliest Hettangian is represented in all these areas, except for Wide Bay, either by a fairly smooth species of *Psiloceras* or by the ribbed subgenus *P. (Franziceras)* similar to *P. (F.) ruidum* (Buckman). The middle Hettangian, probably correlative with the *Alsatites liasicus* zone of Europe, is characterized by *Waehneroceras* at Wide Bay, by *Waehneroceras*, *Discamphiceras*, and *Laqueoceras* at Puale Bay, by *Discamphiceras* in the Wrangell Mountains, and questionably by *Waehneroceras* in the Arctic Coastal Plain near Point Barrow (fig. 8). The late Hettangian, correlative with the *Schlotheimia angulata* Zone, is represented by one specimen of *Schlotheimia* in the Puale Bay area. All these genera are represented by species that are closely similar to, and probably in part identical with, species described from western Europe. Evidently the succession of Hettangian ammonite

genera and species in Alaska is the same as in western Europe.

SINEMURIAN AMMONITES

Sinemurian ammonites have been found in northern Alaska only in the subsurface of the Point Barrow-Cape Simpson area (fig. 7). In southern Alaska, such ammonites occur in five areas from the Wrangell Mountains on the east to Puale Bay on the west (figs. 10-12). The lower Sinemurian zones of *Arietites bucklandi* and *Arnioceras semicostatum* are represented in all these areas except in the Talkeetna Mountains, where beds of that age are not exposed. The uppermost Sinemurian (*Echioceras raricostatum* Zone) is represented in the Wrangell and Talkeetna Mountains and in the Chulitna River area of the Alaska Range. The remainder of the Sinemurian (*Caenisites turneri*, *Asteroceras obtusum*, and *Oxynoticeras oxynotum* Zones) has not been identified by ammonites anywhere in Alaska but is probably represented by poorly fossiliferous beds in the Arctic Coastal Plain, in the Wrangell Mountains, and possibly elsewhere.

Talkeetna Formation in the Nelchina area of the Talkeetna Mountains	Areas of Lower Jurassic outcrops in the Nelchina area and in the upper part of Matanuska River valley	U.S. Geological Survey Mesozoic fossil localities	Ammonite genera	Stage	
1500 ft (457 m) Well-bedded marine strata dominated by siliceous lutite, tuffaceous sandstone, and epiclastic volcanic sandstone	Copper River Basin South side of west part	25939	<i>Grammoceras</i>	Toarcian	Upper
	Northern Horn Mountains	24111, 25316-25319, 25342 25359 and 27508	<i>Pseudolioceras</i> , <i>Haugia</i> and <i>Phymatoceras</i> (?)		
2300 ft (701 m) Well-bedded, tuffaceous marine strata and thick beds of tuff and agglomerate	Northern Horn Mountains	24114, 25940	<i>Haugia</i> and <i>Brodieia</i>		Lower
	East end Sheep Mountain	24787	<i>Dactyloceras</i>		
	South side of upper Matanuska River valley	29198	<i>Harpoceras</i> and <i>Orthodactylites</i>		
8000 ft (2438 m) Thick-bedded volcanic breccia, tuff and some lava flows Base not observed.	Southern Horn Mountains and west part of Sheep Mountain	No fossils found		Pliensbachian	Upper
2500 ft (762 m) Well-bedded marine tuff, tuffaceous sandstone and siliceous lutite. Base not observed and possibly not exposed	Northeast end of Sheep Mountain	24107, 24108, 25941, 29449 and 29450	<i>Amaltheus</i> , <i>Fanninoceras</i> , <i>Protogrammoceras</i> , <i>Arietoceras</i> , <i>Fontanelliceras</i> and <i>Leptaleoceras</i> ,		Lower
	South side of upper Matanuska River valley	27586	<i>Arietoceras</i> , <i>Fanninoceras</i> and <i>Fontanelliceras</i>		
	Valley of Doone Creek on west side of Chickaloon River	6697 and M6171	<i>Apoderoceras</i> and <i>Uptonia</i> ?		
1000-5000 ft (305-1524 m) Well-bedded, marine, siliceous, tuffaceous, lutite and sandstone; limestone and marl; tuff; and some lava flows. Base not observed.	Valley of Doone Creek	6706	<i>Cruciloboceras</i>	Sinemurian (part)	Upper
	Between headwaters of Black and Oshetna Rivers	26722, 28661-63	<i>Paltechioceras</i> and <i>Cruciloboceras</i>		

FIGURE 11.—Occurrences and ages of Early Jurassic ammonites present in the Nelchina area of the Talkeetna Mountains and in the upper part of the Matanuska Valley, southern Alaska.

The lowermost Sinemurian (*Arietites bucklandi* Zone) in the Arctic Coastal Plain is represented by *A. cf. A. bucklandi* (Sowerby) in the Avak test well no. 1 at the depth of 1,836 feet (560 m). It is represented by *Charmasseiceras* in the South Barrow test well no. 3 at depths of 2,412 and 2,419.5 feet (735 and 737 m) and in the South Barrow test well no. 12 at depths of 2,056, 2,061, 2,061.5 and 2,068 feet (627-630 m). The Sinemurian Age of *Charmasseiceras* in the South Bar-

row test well no. 3 is attested by the presence of *Coroniceras* sp. C (pl. 4, fig. 14) in the same well at the depth of 2,470 feet (753 m).

The earliest Sinemurian in the Wrangell Mountains is probably represented by fragmentary arietitid ammonites that occur a little below beds containing *Arnioceras* (figs. 9, 10). Beds of earliest Sinemurian Age in the Chulitna River, Seldovia, and Puale Bay areas are represented by *Paracaloceras rursicostatum* Frebold

EARLY JURASSIC AMMONITES FROM ALASKA

Formation	Generalized description	Fossil localities	Characteristic fossils	Stage
Kialagvik Formation (Lower part beneath 30 m of coarse conglomeratic beds)	1300 ft (396 m) Siltstone, sandy, dark gray to black, contains a few beds of hard buff sandstone, becomes sandier toward top, bears many limy concretions in upper two-thirds.	21235	<i>Inoceramus lucifer</i> and <i>Erycites?</i>	Upper lower to lower middle Bajocian
		ROC 1370 ROC 1366 ROC 1351 and 1356	<i>Tmetoceras scissum</i> and <i>Erycitoides T. scissum</i> <i>Pseudolioceras whitevesi</i> and <i>Erycitoides</i>	Upper lower Bajocian (Aalenian)
		19804	<i>Haugia</i> and <i>Pseudolioceras</i>	Upper lower Toarcian
Unnamed Lower Jurassic beds	160 ft (49 m) Sandstone, conglomeratic, massive, tuffaceous	ROC 1303	<i>Coroniceras?</i>	Lower Sinemurian
	130 ft (40 m) Siltstone and sandstone, limy, partly tuffaceous			
	255 ft (78 m) Sandstone, massive, gray, conglomeratic, tuffaceous	ROC 1282 ROC 1283 21237	} <i>Arnioceras</i>	
	265 ft (81 m) Sandstone and shale, limy, well-bedded, some limestone	ROC 1241 ROC 1240 10820 (part) and 12396 (part)	} <i>Paracoloceras rursicostatum</i> and <i>Badouxie canadensis</i>	?
	245 ft (75 m) Tuff, gray to green interbedded with brown limestone and agglomerate			
	780 ft (238 m) Limestone, mostly thin-bedded, dense, dark-gray; some interbedded limy shale and tuffaceous sandstone that become more common toward top. Grades downward into upper Triassic beds.	? 10820 (part) 3110, 19803, 25694, M1738, 12396 (part) and ROC 1185 3109, 31370	? <i>Schlotheimia</i> <i>Waehneroceras</i> , <i>Psiloceras</i> and <i>Discamphiceras</i> <i>Waehneroceras</i>	Hettangian
Upper Triassic beds (highest part)	320 ft (97.5 m) Limestone, thin-bedded, dense, gray. Some thin beds of sandstone and shale	19806 3108	<i>Monotis</i> cf. <i>M. salinaria</i> <i>Metasibirites</i>	Upper Norian

FIGURE 12.—Occurrences and ages of ammonites present in unnamed Lower Jurassic beds on northeast side of Puale Bay, Alaska Peninsula.

Series	Stage	Arctic Coastal Plain in subsurface of Point Barrow- Cape Simpson area	DeLong Mountains area in northwestern Alaska	Ipnavik-Anaktuvuk Rivers area in north-central Alaska	Ignek Valley to Aichilik River area in northeastern Alaska	Old Rampart area in east-central Alaska
Lower Jurassic	Toarcian	Upper	No fossil data		Kingak Shale (part) 300 ft (91 m) Dark gray clay shale and claystone	
		Lower	?	Thickness unknown Clay shale and limestone concretions		
	Pliensbachian	Upper	Kingak Shale (part) 426 ft (130 m) Clay shale, siltstone, and minor glauconitic sandstone			Glenn Shale (part) 1000 ft (305 m) Shale, siltstone sandstone
		Lower	Kingak Shale (part) 214 ft (65 m) in Barrow test well 3			No fossil data
	Sinemurian	Upper	Exact age span unknown because of lack of fossil evidence	60 ft (18 m) or less Siliceous claystone, some chert and limestone	Kingak Shale (part) 600 ft (183 m) Fissile, black, papery shale	
		Lower	?			
	Hettangian		Kingak Shale (part) 195 ft (59 m) Clay shale and claystone			
						Glenn Shale (part) (included with above)

FIGURE 13.—Correlation of Lower Jurassic rocks in northern and east-central Alaska.

(1967b, p. 26, pl. 7, figs. 1, 2; pl. 9, fig. 1). That species in the Chulitna River area occurs with *Psiloceras canadense* Frebold (1951, p. 3, pl. 6, figs. 1-6; pl. 2, fig. 1; pl. 3, fig. 1) which is now assigned to *Badouxia* Guex and Taylor (1976, p. 525). Those species of *Paracoloceras* and *Badouxia* in association with others in north-central British Columbia were tentatively assigned a middle Hettangian Age by Frebold (1967b, p. 31). Later studies by Guex and Taylor (1976) show, however, that *Paracoloceras* and *Charmasseiceras* in many parts of the world are associated with ammonites of early Sinemurian Age as young as the *Arnioceras semicostatum* zone. Such an age for those genera in British Columbia is confirmed by their association with *Vermiceras multicostatum* (Frebold), which genus is characteristic of the earliest Sinemurian.

The lower Sinemurian zone of *Arnioceras semicostatum* is represented in four areas in Alaska by a finely ribbed species of *Arnioceras* similar to *A. densicosta* (Quenstedt). In the Seldovia area, the same zone is probably represented by the subgenus *Coroniceras* (*Paracoroniceras*), which is not known in younger zones.

The next three younger Sinemurian zones, although not identified faunally, could be represented by poorly fossiliferous beds, or in part by the presence of the genus *Arnioceras*, which in Europe ranges upward through most of the *Asteroceras obtusum* Zone.

The uppermost Sinemurian zone of *Echioceras raricostatum* is represented in both the Wrangell and Talkeetna Mountains by *Paltechioceras* cf. *P. harbledownense* (Crickmay) in association with species of *Cruciloboceras* that closely resemble described species

Series	Stage	Wrangell Mountains in southern Alaska	Talkeetna Mountains Nelchina area southern Alaska	Chulitna River area in Healy (A-6) quadrangle, south-central part of Alaska Range	Seldovia area on the Kenai Peninsula	Puale Bay area on the Alaska Peninsula
Lower Jurassic	Toarcian		1500 ft (457 m) Tuffaceous sandstone and siliceous lutite	No fossil data		Kialagvik Formation Lower 200 ft (62 m) Sandy siltstone. Rests sharply on Sinemurian beds
			2300 ft (701 m) Tuffaceous, marine beds; thick-bedded tuff and agglomerate			
		Lubbe Creek Formation 100–300 ft (30.5–91.5 m) Gray spiculite chert. Some bioclastic coquina near top	8000 ft (2438 m) Volcanic rocks			
	Pliensbachian		2500 ft (762 m) Marine tuff tuffaceous sandstone and siliceous lutite		No fossil data	No fossil data
		Upper member of McCarthy Formation 2000–2500 ft (610–762 m) Mostly dark gray, thin- to medium- bedded, finely laminated organic chert and spiculite. Some fissile shale and limestone that is most common in upper part				Absence probably due to faulting rather than to erosion
	Sinemurian		1000–5000 ft (305–1524 m) Marine lutite, sandstone, marl, tuff, limestone, and lava flows	Massive tuffaceous argillite		
				No fossil data		
	Hettangian	All contacts are gradational		Gray calcareous sandstone and silty limestone	2000–3000 ft (610–915 m) Red to green lava, tuff, agglomerate and sand- stone. Some gray limestone	1040 ft (317 m) Massive conglomeratic sandstone and siltstone
			Not exposed	No fossil data		780 ft (238 m) Limestone, dense, thin-bedded; some limy shale. Grades down- ward into Triassic beds.

FIGURE 14.—Correlation of Lower Jurassic rocks in southern Alaska.

from western Europe. *Paltechioceras* occurs also in the Chulitna River area in association with *Arctoasteroceras jeletzkyi* Frebold, whose type specimens in arctic Canada were collected along with *Oxynoticeras oxynotum* (Quenstedt) below beds containing echioceratid ammonites. This shows that *Arctoasteroceras* occurs in the highest two zones of the Sinemurian.

Other mollusks of some age significance during Sinemurian time include the bivalves *Otapiria tailleuri* Imlay and *Entolium? semiplicatum* Hyatt. Of these, *O. tailleuri* in the subsurface of northern Alaska occurs in the South Barrow test well No. 3 at depths of 2,412–2,417 feet (735 m) and in the South Barrow test well No. 12 at depths of 2,055–2,078.5 feet (627–634 m). Comparisons with the ammonite occurrences in those test wells (table 2) show clearly that *O. tailleuri* is

associated with ammonites of early Sinemurian Age, occurs above ammonites of Hettangian Age, and has its highest occurrence at 214 feet (65 m) below ammonites of late Pliensbachian Age.

Otapiria tailleuri Imlay has been found in outcrops in northern Alaska in a fairly thin unit of fissile clay shale that extends from the Ipnayik River eastward at least 200 miles (320 km) to the Kavik River (Imlay, 1967, p. B6, B7; Imlay and Dettnerman, 1973, p. 12; Dettnerman and others, 1975, p. 44). Apparently it does not occur throughout the entire thickness of the shale in all areas. It does occur in the same rock slabs as some crushed, generically undeterminable ammonites (Imlay, 1967, p. B6). It does not occur on the same slabs as some ammonites of early Pliensbachian Age that are described herein as *Uptonia* cf. *U. jamesoni* (J. de C. Sowerby) and

TABLE 2.—Early Jurassic ammonites from well cores in northern Alaska

Genus and species	Depths in test wells, in feet, from which fossils were obtained				Stages
	Simpson test well 1	Avak test well 1	South Barrow test well 3	South Barrow test well 12	
<i>Dactyloceras</i> (<i>Orthodactylites</i>) cf. <i>D. (O.) kanense</i> McLearn.	-----	-----	2,016	-----	Lower Toarcian.
<i>D. sp</i> -----	-----	-----	1,772; 2,017; 2,018	-----	Do.
<i>Catacoeloceras</i> sp. juv -----	-----	-----	2,063	-----	Do.
<i>Amaltheus margaritatus</i> (Montfort) -----	-----	-----	2,099; 2,111	-----	Upper Pliensbachian.
<i>A. cf. A. margaritatus</i> (Montfort) -----	-----	-----	2,107	-----	Do.
<i>A. stokesi</i> (J. Sowerby) -----	-----	-----	2,193; 2,198	-----	Do.
<i>A. cf. A. stokesi</i> (J. Sowerby) -----	-----	-----	2,186	-----	Do.
<i>A. sp</i> -----	5,677; 5,680; 5,691	-----	2,069-2,184	-----	Do.
<i>A. (Pseudoamaltheus) engelhardti</i> (d'Orbigny).	-----	-----	2,090	-----	Do.
<i>Arietites</i> cf. <i>A. bucklandi</i> (Sowerby) -----	-----	1,836	-----	-----	Lower Sinemurian.
<i>Coroniceras</i> sp. B -----	-----	-----	-----	1,987.4-1,987.6	Do.
<i>Coroniceras</i> sp. C -----	-----	-----	2,470	-----	Do.
<i>Arnioceras</i> sp. juv -----	-----	-----	-----	2,056	Do.
<i>Charmasseiceras</i> sp. -----	-----	-----	-----	2,056	Do.
<i>C. cf. C. marmoreum</i> (Oppel) -----	-----	-----	2,412; 2,419.5	2,061.5; 2,068	Do.
<i>Psiloceras</i> (<i>Franziceras</i>) sp -----	-----	-----	-----	2,170.5	Hettangian.
<i>Waehneroceras?</i> sp -----	-----	-----	-----	2,181.5	Do.

Uptonia sp. (pl. 9, figs. 9-11, 17). These occurrences and associations in both surface and subsurface rocks show that the range of *O. tailleuri* is probably only Sinemurian and that it may not be younger than early Sinemurian.

The other pelecypod that is useful in dating Sinemurian rocks is *Entolium? semiplicatum* Hyatt. (See Crickmay, 1933, p. 52, pl. 14, figs. 4-7.) That species to date has been found in Alaska only in the Wrangell Mountains, where it is common in dark-gray to black fissile shale at the top of the McCarthy Formation. In that shale, which is similar lithologically to the fissile shale in which *Otapiria* occurs in northern Alaska, it is associated with the ammonites *Paltechioceras*, (USGS Mesozoic locs. 30139 and 31174) and *Cruciloboceras* cf. *C. crucilobatum* Buckman (USGS Mesozoic loc. 14030) of latest Sinemurian Age. It has been found elsewhere in the Pacific Coast region on the North Fork of the American River in eastern California (Crickmay, 1933, p. 52; Imlay, 1968, p. C7) in association with *Cruciloboceras*. It is closely similar to and possibly within the range of variation of *Entolium balteatum* Crickmay (1928, p. 62, pl. 4e-g) from Parson Bay in British Columbia. That species is associated with *Paltechioceras harbledownense* (Crickmay) (1928, p. 61, pl. 4a-d), which is probably represented at the top of the McCarthy Formation in Alaska, (pl. 4, figs. 15-17, 22).

These data concerning *Entolium? semiplicatum* and *Otapiria tailleuri* show that both are of Sinemurian Age and both occur in similar fissile shale. The fact that they have never been found associated in the same bed suggests, therefore, that they are of slightly different ages, or lived in somewhat different marine environments, or were separated by a land barrier.

PLIENSCHACHIAN AMMONITES

Pliensbachian ammonites have been found in both the subsurface and surface in northern Alaska, in the Old Rampart area in east-central Alaska, and in the Wrangell and Talkeetna Mountains in the eastern part of southern Alaska. They probably occur in the Yakutat District in the northern part of southeastern Alaska (figs. 8-11).

In northern and east-central Alaska, most Pliensbachian ammonite occurrences consist of *Amaltheus* of late Pliensbachian Age. In the South Barrow test well No. 3, *Amaltheus* is represented by the same species in the same succession as in northwestern Europe (table 2). The only other ammonite of Pliensbachian Age found in northern Alaska consists of a single large specimen of *Uptonia* similar to *U. jamesoni* (J. de C. Sowerby) from an outcrop on Lisburne Ridge (USGS Mesozoic loc. 29774) between the Iqnavik and Etivluk Rivers in the Howard Pass quadrangle. This specimen is good evidence that the beds in which it occurs are of earliest Pliensbachian Age. In addition *Uptonia*, rather than *Cruciloboceras*, may be represented in the Iqnavik-Anatuvuk River area by several crushed molds (pl. 9, figs. 9-11) (USGS Mesozoic locs. 29281, 29282, and 29775) that resemble the inner whorls of the large specimen of *Uptonia*.

The Pliensbachian ammonite successions in the Wrangell and Talkeetna Mountains in southern Alaska contrast with those in northern Alaska by being fairly complete, by containing some ammonites that are closely similar to taxa in the Mediterranean region, and by the genus *Amaltheus* being represented by only two specimens from a single locality. In the Wrangell Mountains the *Uptonia jamesoni* zone is represented by *U. cf.*

U. dayiceroides Mouterde, which was collected with *Prodactylioceras italicum* (Fucini) less than 100 feet (30 m) below the upper contact of the upper member of the McCarthy Formation (USGS Mesozoic locs. 28671-28673, 28175). Most of the *Tragophylloceras ibex* zone is represented by *Tropidoceras actaeon* (d'Orbigny), which was collected with *Cruciloboceras* cf. *C. pacificum* Frebold about 1 m below the upper contact of the McCarthy Formation (USGS Mesozoic loc. 28534). The uppermost Pliensbachian is represented by *Arietoceras* at the top of the McCarthy Formation (USGS Mesozoic loc. 28688) and in the overlying Lubbe Creek Formation (USGS Mesozoic loc. 28531).

In the Talkeetna Mountains (figs. 9, 11), the *Uptonia jamesoni* zone is represented by a well-preserved specimen of *Apoderoceras* cf. *A. subtriangulare* (Young and Bird) and a fragmentary specimen that probably represents *Uptonia*. The next two younger zones of western Europe are not represented by fossils, but the uppermost two Pliensbachian zones are represented by *Leptaleoceras pseudoradians* (Reynes), *Fontanelliceras* cf. *F. fontanellense* (Gemmellaro) *Arietoceras* cf. *A. domarense* (Meneghini), *Protogrammoceras* cf. *P. paltum* (Buckman), *P. cf. P. argutum* (Buckman), *Fanninoceras kunae* McLearn, and *Amaltheus* cf. *A. stokesi* (J. Sowerby).

TOARCIAN AMMONITES

The Toarcian is not well represented by ammonites in Alaska except in the Talkeetna Mountains (figs. 8, 9, 11). In the subsurface of the arctic Coastal Plain, only the lowermost Toarcian (characterized by *Dactylioceras*) has been identified. In the DeLong Mountains, that genus is succeeded upward by *Harpoceras* cf. *H. exaratum* (Young and Bird), which succession accounts for about two-fifths of Toarcian time. The same species of *Harpoceras* was obtained in northeastern Alaska on a tributary of the Sadlerochit River (USGS Mesozoic loc. 22081). Near Red Hill on Ignek Creek in the Mt. Michelson (C-4) quadrangle, was obtained *Pseudo-*

lioceras cf. *P. compactile* (Simpson) and *P. cf. P. lythense* (Young and Bird) of late Toarcian Age at USGS Mesozoic loc. 23772 (Imlay, 1955, p. 89 pl. 12, figs. 17, 18, 20, 21).

In the Talkeetna Mountains of southern Alaska, the lower Toarcian is represented by a similar sequence of *Dactylioceras* and *Harpoceras* as in northern Alaska. The lower upper Toarcian is represented by *Haugia*, *Brodieia*, and *Pseudolioceras*, which genera in association are good evidence for correlation with the *Haugia variabilis* Zone of western Europe. Still higher occurs *Grammoceras*, a genus that in western Europe occurs only in the uppermost two zones of the Toarcian.

In the Puale Bay area, the middle to lower upper Toarcian is represented by *Haugia* and *Pseudolioceras*.

AMMONITE FAUNAL SETTING

No distinct ammonite faunal province existed in Alaska during most of Early Jurassic time, as shown by the remarkable resemblance of most of the ammonites present to those elsewhere in the world. During late Pliensbachian time, a close tie with northern Europe is shown by the presence of *Amaltheus* in northern Alaska and in the Talkeetna Mountains of southern Alaska. At the same time, a close tie with the Mediterranean region is shown by the presence of *Arietoceras* in the Wrangell Mountains and of *Arietoceras*, *Leptaleoceras*, *Fontanelliceras*, and *Protogrammoceras* in the Talkeetna Mountains but not farther north. These occurrences of *Arietoceras* and *Leptaleoceras* in southern Alaska are not much farther north than their northernmost known occurrences in the southern Yukon (Frebold, 1970, p. 446).

GEOGRAPHIC DISTRIBUTION

The geographic occurrences of Early Jurassic ammonites from Alaska described herein are shown on figures 1-6 and tables 3-6. Detailed descriptions of the occurrences are given in table 7.

TABLE 3.—*Geographic distribution of Early Jurassic ammonites from outcrops in northern and east-central Alaska*

[Quadrangle occurrences are listed in table 7. Numbers 5–15 are keyed to area numbers in figure 1. Higher numbers are USGS Mesozoic locality numbers]

	Northwestern Alaska					North-central Alaska			Northeastern Alaska				East-central Alaska			
	Delong Mountains												Old Rampart area			
	Clay shale					Claystone			Kingak shale				Glenn shale (in part)			
	5	6	7	8	9	10	11	12	13				14		15	
	29159	29160	29161	29163	29776	M2441	29164	29775	29774	29281	29282	23772	22081	29165	30074	
<i>Psiloceras</i> sp															X	
<i>P.?</i> sp																
<i>P. (Franziceras)</i> sp															X	
<i>P. (F.)</i> cf. <i>P. (F.) ruidum</i> (Buckman)														X	X	
<i>Uptonia</i> cf. <i>U. jamesoni</i> (J. de C. Sowerby)								X								
<i>Uptonia?</i> sp								X	X	X						
<i>Amaltheus margaritatus</i> (Montfort)						X										
<i>A. stokesi</i> (J. Sowerby)													X			X
<i>A.</i> cf. <i>A. stokesi</i> (J. Sowerby)													X			
<i>A.</i> sp							X									
<i>Dactylioceras</i> (<i>Orthodactylites</i>) cf. <i>D. (O.) directum</i> Buckman			X													
<i>Harpoceras</i> cf. <i>H. exaratum</i> (Young and Bird)	X	X	X	X								X				
<i>Eleganticeras</i> sp. juv				X												
<i>Pseudolioceras</i> cf. <i>P. compactile</i> (Simpson)										X						
<i>P.</i> cf. <i>P. lythense</i> (Young and Bird)										X						

[illegible]

TABLE 5.—*Geographic distribution of Early Jurassic ammonites in the Talkeetna Mountains in southern Alaska*

[Quadrangle occurrences are listed in table 7. Numbers 20–23 are keyed to generalized area numbers in figure 1. Higher numbers are USGS Mesozoic locality numbers]

Genus and species	Talkeetna Mountains																									
	Western Copper River basin	Sheep Mountain to Horn Mountain										Chickaloon- Kings Mountain area				Oshetna River headwaters										
	Talkeetna Formation																									
	20	21										22				23										
25939	24107	24108	24111	24114	24787	25316	25317	25318	25319	25342	25359	25940	25941	27508	29449	29450	M6171	6697	6706	27586	29198	26722	28661	28662	28663	
<i>Paltechioceras</i> cf. <i>P. harbledownense</i> (Crickmay) -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	×
<i>C.</i> cf. <i>C. densinodulum</i> Buckman -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---
<i>C.</i> cf. <i>C. muticum</i> (d'Orbigny) -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	×	×
<i>C.</i> cf. <i>C. submuticum</i> (Oppel) -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	×	×	×
<i>C.</i> sp. juv -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---
<i>Apoderoceras</i> cf. <i>A. subtriangulare</i> (Young and Bird) -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	×	---	---	---	---	---	---	---
<i>Fanninoceras kunae</i> McLearn -----	---	×	×	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	×	---	---	---	---	---	---
<i>F.</i> cf. <i>F. carlottense</i> McLearn -----	---	---	×	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---
<i>Uptonia</i> ? sp. B -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---	---	---
<i>Amaltheus</i> cf. <i>A. stokesi</i> (J. Sowerby) -----	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>Dactylioceras</i> cf. <i>D. commune</i> (J. Sowerby) -----	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---
<i>D.</i> (<i>Orthodactylites</i>) <i>kanense</i> McLearn -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---
<i>Arietoceras</i> cf. <i>A. domarensense</i> (Meneghini) -----	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	×	---	---	---	---	---	---
<i>Leptaleoceras pseudoradians</i> (Reynes) -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---
<i>Fontanelliceras</i> cf. <i>F. fontanellense</i> (Gemmellaro) -----	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	×	---	---	---	×	---	---	---	---	---	---
<i>Harpoceras</i> cf. <i>H. exaratum</i> (Young and Bird) -----	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---
<i>Protogrammoceras</i> cf. <i>P. paltum</i> (Buckman) -----	---	---	×	---	---	---	---	---	---	---	---	---	×	---	×	---	---	---	---	---	---	---	---	---	---	---
<i>P.</i> cf. <i>P. argutum</i> (Buckman) -----	---	---	×	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>Pseudolioceras</i> sp -----	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>Grammoceras</i> sp -----	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>Phymatoceras</i> ? sp -----	---	---	---	---	---	×	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>Brodieia</i> cf. <i>B. tenuicostata</i> var. <i>nodosa</i> (Jaworski) -----	---	---	---	---	---	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>Haugia</i> cf. <i>H. grandis</i> Buckman -----	---	---	×	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>H.</i> cf. <i>H. variabilis</i> (d'Orbigny) -----	---	---	---	---	---	---	×	---	---	---	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---
<i>H.</i> cf. <i>H. compressa</i> Buckman -----	---	---	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
<i>H.</i> sp -----	---	---	---	---	---	×	---	---	×	---	×	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

Genus and species	South-central part of Alaska Range									Kenai Peninsula											
	Chulitna River area Healy A-6 quadrangle							Talkeetna B-6 and C-6 quadrangles		Seldovia area											
	Sandstone, siltstone, limestone, and argillite									Volcanic rocks, sandstone, limestone											
	24							25		26											
16229	31260	31261	31262	31263	31264	31265	31266	30907	30908	31270	2978	2979	2981	21242	22664	31128	31634	31637	31640	31648	31650
<i>Lytoceras</i> sp																					
<i>Psiloceras</i> cf. <i>P. planorbis</i> (J. de C. Sowerby)																					
<i>P. (Franziceras)</i> cf. <i>P. (F.) ruidum</i> (Buckman)														x	x					x	
<i>Discamphiceras</i> cf. <i>D. toxophorum</i> (Buckman)																					
<i>Laqueoceras</i> cf. <i>L. sublaqueus</i> (Waehner)																					
<i>Waehneroceras</i> cf. <i>W. tenerum</i> (Neumayr)																					
<i>W.</i> cf. <i>W. portlocki</i> (Wright)																					
<i>Scholtheimia</i> sp																					
<i>Badouxia canadensis</i> (Frebold)							x														
<i>B. columbiae</i> (Frebold)						x															
<i>Charmasseiceras</i> cf. <i>C. marmoreum</i> (Oppel)												x						x			
<i>Coroniceras</i> sp. A									x												
<i>Coroniceras?</i> sp										x							x		x		
<i>C. (Paracoroniceras)</i> sp																x					x
<i>Arnioceras</i> cf. <i>A. densicosta</i> (Quenstedt)	x	x		x	x		x														
<i>Paracoloceras rursicostatum</i> Frebold							x						x								
<i>Arietitid</i> genus undetermined					x			x		x											
<i>Arctoasteroceras jeletzkyi</i> Frebold				x																	
<i>Paltechioceras (Orthechioceras?)</i> sp				x																	
<i>Pseudolioceras</i> sp																					
<i>Haugia</i> cf. <i>H. grandis</i> Buckman																					
<i>H.</i> cf. <i>H. compressa</i> Buckman																					

23

Higher numbers are USGS Mesozoic locality numbers]

Alaska Peninsula																						Wide Bay area
Puale Bay-Alinchak Bay area																						
Limestone, shale, siltstone, sandstone, and massive conglomerate																						
27																						
3109	3110	10820	12075	12394	12396	19803	19804	21257	25694	29267	29268	M1738	31370	31372	ROC1185	ROC1240	ROC1241	ROC1282	ROC1283	ROC1303	ROC3002	ROC Wide Bay Well
--	--	x	x	--	--	--	--	--	--	--	--	x	--	--	x	--	--	--	--	--	x	--
--	--	--	--	--	--	--	--	--	--	--	--	x	--	--	--	--	--	--	--	--	--	--
--	x	--	--	--	--	--	--	--	x	--	--	--	--	--	--	--	--	--	--	--	--	--
--	x	--	x	--	--	x	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	x	--	x	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
--	--	--	--	--	--	--	--	--	--	--	--</											

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska

[Leaders (—), number not assigned]

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
1			Simpson test well 1, lat 70°57'05" N., long 155°21'45" W., Cape Simpson area, Barrow (A-2) quadrangle northern Alaska. Kingak Shale at depths of 5,677-5,691 ft (1,730-1,735 m). Late Pliensbachian.
2			Avak test well, lat 71°15'02" N., long 156°28'06" W., Point Barrow area, Barrow (B-4) quadrangle, northern Alaska. Kingak Shale at depths of 1,836 ft (660 m). Early Sinemurian.
3			South Barrow test well 3, lat 71°15'51" N., long 156°37'55" W., Point Barrow area, Barrow (B-4) quadrangle, northern Alaska. Kingak Shale at depths of 1,772-2,470 ft (540-753 m). Early Sinemurian to early Toarcian.
4			South Barrow test well 12, lat 71°14'18" N., long 156°20'06" W., Point Barrow area, Barrow (A-4) quadrangle, northern Alaska. Kingak Shale at depths of 1,987.4-2,181.5 ft (606-665 m). Hettangian to early Sinemurian.
5	29159	AF1024	British Petroleum (Alaska) Inc., 1964. On north bank of Thetis Creek headwaters, lat 68°41'51" N., long 164°45'05" W., north side of DeLong Mountains, DeLong Mts. (C-5) quadrangle, northern Alaska. Kingak Shale. Early to middle Toarcian.
5	29160	AF1025	British Petroleum (Alaska) Inc., 1964. Near USGS Mesozoic loc. 29159. Kingak Shale. Early to middle Toarcian.
5	29161	AF1027	British Petroleum (Alaska) Inc. 1964. Near USGS Mesozoic loc. 29159. Kingak Shale. Early to middle Toarcian.
5	29163	AF1028	British Petroleum (Alaska) Inc., 1964. Near USGS Mesozoic loc. 29159. Kingak Shale. Early to middle Toarcian.
5	29776	68ATo46F	I. L. Tailleux and H. A. Tourtelot, 1968. Cutbank on large eastern tributary of Thetis Creek (near USGS Mesozoic loc. 29159), lat 68°40.8' N., long 164°45.5' W., Kingak Shale. Early to middle Toarcian.
6	29164	AF26	British Petroleum (Alaska) Inc., 1964. North side of DeLong Mountains, on east bank of Ipewik River, lat 68°40' N., long 164°13' W., DeLong Mts. (C-4) quadrangle, northern Alaska. Unnamed beds. Late Pliensbachian.
6	M2441	64ATr204F	I. L. Tailleux, 1964. On Ipewik River on north side of DeLong Mountains near USGS Mesozoic loc. 29164, lat 68°37' N., long 164°11' W., northern Alaska. Unnamed beds. Late Pliensbachian.
7	29775	68ATo23	I. L. Tailleux and H. A. Tourtelot, 1968. Cutbanks on east side of Ipnavik River, lat 68°40.6' N., long 157°03.5' W., Howard Pass quadrangle, northern Alaska. Unnamed oil shale overlying Shublik Formation (Triassic). Early Pliensbachian or latest Sinemurian.
8	29774	68ATo21d	I. L. Tailleux and H. A. Tourtelot, 1968. Blankenship Creek in Lisburne Ridge area, lat 68°37.5' N., long 156°42.5' W., Howard Pass quadrangle, northern Alaska. Unnamed oil shale overlying Triassic Shublik Formation. Early Pliensbachian.
9	29281	65ATr123B	I. L. Tailleux, 1965. East wall of Tiglukpuk Creek on south flank of Tiglukpuk anticline, lat 68°21'15" N., long 151°52'30" W., Chandler Lake quadrangle, northern Alaska. Unnamed beds overlying Shublik Formation. Early Pliensbachian or latest Sinemurian.
9	29282	65ATr149.4	I. L. Tailleux, 1965. Southwest wall of Firestone Creek, a tributary of Tiglukpuk Creek, just north of mountain front, lat 68°18'15" N., long 151°48'00" W., Chandler Lake quadrangle, northern Alaska. Float from unnamed beds overlying Shublik Formation. Early Pliensbachian or latest Sinemurian.
10	23772	50AGr61	R. W. Imlay, George Gryc, and Allan Kover, 1950. Near Red Hill on Ignek Creek, lat 69°36'

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
			N., long 146°06' W., Mt. Michelson (C-4) quadrangle, northern Alaska. Kingak Shale. Late Toarcian.
11	22081	48ASa146	E. G. Sable, 1948. About 1 mile (1.6 km) northeast of mouth of Fire Creek and 0.4 mi (0.6 km) above the mouth of a south-flowing tributary of the Sadlerochit River, Mt. Michelson (C-2) quadrangle, northern Alaska. Kingak Shale, within middle third and about 1,700 ft (518 m) above its base in black earthy shale. Early to middle Toarcian.
12	29165	AG525	British Petroleum (Alaska) Inc., 1964. Three miles (4.8 km) east of Lake Schrader, lat 69°22' N., long 144°28' W., Mt. Michelson (B-2) quadrangle, northern Alaska. Kingak Shale. Late Pliensbachian.
13	30074	71ADt330c	R. L. Detterman, 1971. Cutbank on west side of Aichilik River about 2 mi (3.2 km) N. 30° E. of VABM ATTE on northern front of Brooks Range, lat 69°33' N., long 143°05' W., Barter Island (C-4) quadrangle, northern Alaska. Kingak Shale, 800 to 1,000 ft (244 to 205 m) above base. Late Pliensbachian.
14	29737	63ABe47B	W. P. Brosge, 1963. Spike Mountain, about 3 miles (4.8 km) north of Old Rampart, lat 63°35.75' N., long 141°40' W., Porcupine River area, Coleen quadrangle, east-central Alaska. Glenn Shale (in part). A thin unit of red conglomeratic sandstone interbedded with black, fine-grained quartzitic sandstone. Early Hettangian.
14	29738	67ARr-337a	W. P. Brosge, 1967. Same location as USGS Mesozoic loc. 29737. Fossils from apparent top of 120 ft (36.6 m) of beds overlying granite. Early Hettangian.
14	29739	67ARr-337d	Same data as Mesozoic loc. 29738 but from 20 ft (6 m) below apparent top. Early Hettangian.
14	29740	67ARr3371-n	Same data as Mesozoic loc. 29738 but from 40 ft (12 m) below apparent top. Early Hettangian.
14	29742	67ARr-337e-h	Same data as Mesozoic loc. 29738 but from 80 ft (24 m) below apparent top. Early Hettangian.
14	29743	67ARr-337q-s	Same data as Mesozoic loc. 29738 but from 100 ft (30.5 m) below apparent top. Early Hettangian.
15	29340	KP0688	British Petroleum (Alaska) Inc. Float from river bluff in the middle fork Salmon Trout River about 9 miles (14.4 km) east southeast of Old Rampart in Porcupine River area, Coleen quadrangle, east-central Alaska. Glenn Shale (in part). Noncalcareous, pyritic, thin-bedded mudstone interbedded with dark-gray papery shale exposed on bluff. Late Pliensbachian.
16	29773	68APs54F	George Plafker, 1968. Float from sec. 34, T. 23 S., R. 35 E., Yakutat (D-4) quadrangle, southeastern Alaska. Yakutat Group. Probably early Pliensbachian rather than late Sinemurian.
17	28671	62AMK-66	E. M. MacKevett, Jr., 1962. On ridge in south-central part of sec. 23, T. 4 S., R. 16 E., McCarthy (C-4) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation. Less than 100 ft (30.5 m) stratigraphically below the Lubbe Creek Formation. Early Pliensbachian.
17	28672	62AMK-66A	E. M. MacKevett, Jr., 1962. Same description as Mesozoic loc. 28671 and same age.
17	28673	62AMK-67	E. M. MacKevett, Jr., 1962. Collected slightly east of Mesozoic loc. 28671 at the same stratigraphic position.
17	28675	62AMK-69	E. M. MacKevett, Jr., 1962. West-central part of NW¼SW¼ sec. 23, T. 4 S., R. 16 E., at altitude of 6,300 ft (1,920 m), McCarthy (C-4) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation, less than 100 ft (30.5 m) stratigraphically below Lubbe Creek Formation. Early Pliensbachian.

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
17	28677	62AMK-88	E. M. MacKevett, Jr., 1962. Near center of NW¼ sec. 24, T. 4 S., R. 16 E., McCarthy (C-4) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation. Float from impure limestone near an intraformational fault. Early Sinemurian.
17	29870	67AWK-112	E. M. MacKevett, Jr., 1967. On north side of Chitistone Mountain in east-central part of NE¼NW¼ sec. 27, T. 4 S., R. 16 E., McCarthy (B-4) quadrangle, Wrangell Mountains, southern Alaska. Upper member(?) of McCarthy Formation. Hettangian.
17	29890	67AWK-113	E. M. MacKevett, Jr., 1967, near Mesozoic loc. 29870 in same quarter of a quarter section and same stratigraphic position. Hettangian.
17	29891	67AMK-233	E. M. MacKevett, Jr., 1967. NE¼NW¼ sec. 26, T. 4 S., R. 16 E., McCarthy (B-4) quadrangle, Wrangell Mountains. Upper member of McCarthy Formation, a few hundred feet (100 m) above base of member. Hettangian.
18	14030	27AM-F5	F. H. Moffit, 1927. McCarthy Creek valley, near creek on east side, 2.5 mi (4 km) from the glacier, NE¼NE¼ sec. 36, T. 3 S., R. 14 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation, probably a few hundred feet (100 m) above base. Late Sinemurian.
18	28531	19946-39	Standard Oil Co. of California. East of McCarthy Creek near center sec. 25, T. 3 S., R. 14 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Lubbe Creek Formation. Late Pliensbachian.
18	28533	F-6	E. M. MacKevett, Jr., 1961. SE¼NE¼NW¼ sec. 2, T. 4 S., R. 14 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation. About 1,000 ft (305 m) stratigraphically above base of member. Early Sinemurian.
18	28534	F10	E. M. MacKevett, Jr., 1961. On ridge west of McCarthy Creek, SW¼NW¼NW¼ sec. 26, T. 3 S., R. 14 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation, a few feet (about 1 m) below upper contact. Early Pliensbachian.
18	28535	F4	E. M. MacKevett, Jr., 1961. South-central part NW¼SE¼ sec. 11, T. 4 S., R. 15 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Float from upper member of McCarthy Formation. Early Sinemurian.
18	28537	F36	E. M. MacKevett, Jr., 1961. NE. cor. sec. 16., T. 4 S., R. 15 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation; probably in middle part of member. Early Sinemurian.
18	28538	F35	E. M. MacKevett, Jr., 1961. A little south of USGS Mesozoic loc. 28537. Early Sinemurian.
18	28540	F45	E. M. MacKevett, Jr., 1961. NW¼SW¼NE¼ sec. 9, T. 4 S., R. 15 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Float collected about 400 ft (122 m) below top of upper member of McCarthy Formation. Early Pliensbachian.
18	28678	62AMK-20	R. W. Imlay, 1962. Southwest of West Fork in NE. cor. NW¼ sec. 3, T. 4 S., R. 15 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Lubbe Creek Formation. Early Toarcian(?)
18	28688	62AMK-143	E. M. MacKevett, Jr., 1962 West of Nizina River in SE¼NW¼SW¼ sec. 16, T. 3 S., R. 16 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Near top of upper member of McCarthy Formation. Late Pliensbachian.
18	28690	62AMK-16	R. W. Imlay, 1962. In saddle on ridge between

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
			Lubbe Creek and Diamond Creek east of McCarthy Creek, in east-central part of NE¼ sec. 7, T. 4 S., R. 15 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation. Probably late Sinemurian.
18	28704	62AMK-62A	E. M. MacKevett, Jr., 1962. West of McCarthy Creek opposite Lubbe Creek on ridge in east-central part of SW¼ sec. 35, T. 3 S., R. 14 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. About 1,200 ft (366 m) above base of upper member of McCarthy Formation. Probably late Sinemurian.
18	28706	62AMK-63	E. M. MacKevett, Jr., 1962. On ridge west of McCarthy Creek near center NW¼NW¼ sec. 35, T. 3 S., R. 14 E., McCarthy (C-5) quadrangle, Wrangell Mountains, southern Alaska. In upper part of upper member of McCarthy Formation. Early Sinemurian.
18	29118	64AMK-82	E. M. MacKevett, Jr., 1964. South-central part of SW¼NW¼ sec. 35, T. 4 S., R. 15 E., McCarthy (B-5) quadrangle, Wrangell Mountains, southern Alaska. Transitional beds at base of upper member of McCarthy Formation. Early Sinemurian.
18	29119	64AMK-84	E. M. MacKevett, Jr., 1964. 300 ft (91.5 m) southeast of USGS Mesozoic loc. 29118 in same quarter of a quarter section. Transitional beds at base of upper member of McCarthy Formation. Early Sinemurian.
18	29121	64AMK-108	E. M. MacKevett, Jr., 1964. In area of small folds 40 ft (12 m) S. 60° W. of summit of Hill 6580. NE¼NE¼ sec. 34, T. 4 S., R. 15 E., McCarthy (B-5) quadrangle, Wrangell Mountains, southern Alaska. Low in upper member of McCarthy Formation. Early Sinemurian.
18	31174	63AWz-150	E. M. MacKevett, Jr., 1963. On ridge between Root Glacier and Kennicott Glacier. NE¼SW¼ sec. 19, T. 3 S., R. 14 E., McCarthy (C-16) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation, 150 ft (46 m) below top. Late Sinemurian.
18	31175	63AWz-152	E. M. MacKevett, Jr., 1963. Same location as USGS Mesozoic loc. 31174 but only 100 ft (30.5 m) below top of upper member of McCarthy Formation. Late Sinemurian.
19	14472	28AM-F9	F. H. Moffit, 1928. Head of first northern tributary of Mill Creek, SE¼ sec. 33, T. 3 S., R. 11 E., McCarthy (C-7) quadrangle, Wrangell Mountains, southern Alaska. Upper member of McCarthy Formation. Late Sinemurian.
19	30137	71AMK-31E	E. M. MacKevett, Jr., 1971. Center of N¼SW¼ sec. 8, T. 4 S., R. 11 E., McCarthy (C-7) quadrangle, Wrangell Mountains, southern Alaska. Lower part of upper member of McCarthy Formation. Early Sinemurian.
19	30138	71AMK-36B	E. M. MacKevett, Jr., 1971. SW¼NE¼SW¼ sec. 17, T. 4 S., R. 12 E., McCarthy (C-7) quadrangle, Wrangell Mountains, southern Alaska. Low in upper member of McCarthy Formation. Early Sinemurian.
19	30139	71AFd-263B	A. B. Ford, 1971. On ridge in south-central part of SE¼NW¼ sec. 17, T. 4 S., R. 12 E., McCarthy (C-7) quadrangle, Wrangell Mountains, southern Alaska. Low in upper member of McCarthy Formation. Late Sinemurian.
19	30140	71AFd264	A. B. Ford, 1971. On ridge in SW. cor. sec. 17, near sec. 18, T. 4 S., R. 12 E., McCarthy (C-7) quadrangle, Wrangell Mountains, southern Alaska. Early Sinemurian.
20	25939	55AGz182	Arthur Grantz, 1955. Lat 61°54'02" N., long 146°52'00" W., Valdez (D-8) quadrangle, south side of western Copper River basin, southern Alaska. Talkeetna Formation. Late Toarcian.

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
21	24107	52AGz63	R. D. Hoare, R. W. Imlay, and Arthur Grantz, 1952. On south tributary of Squaw Creek, 1.98 miles (3.2 km) N. 28½° E. of southeast summit of Gunsight Mountain, Anchorage (D-2) quad. Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Pliensbachian.
21	24108	52AGz66	Arthur Grantz, R. D. Hoare, and R. W. Imlay, 1952. Differs from USGS Mesozoic loc. 24107 by being 1.91 miles (3.1 km) N. 28½° E. of southeast summit of Gunsight Mountain. Late Pliensbachian.
21	24111	52AGz253	Arthur Grantz, R. D. Hoare, and R. W. Imlay, 1952. 5.04 miles (8 kms) N. 33° E. of the mouth of Sheep Creek, Anchorage (D-2) quadrangle, Talkeetna Mountains, southern Alaska. Early Toarcian.
21	24114	52AGz98	Arthur Grantz, R. D. Hoare, and R. W. Imlay, 1952. From north branch of North Creek, 1.61 miles (2.5 km) N. 45° W. of its mouth. Talkeetna Mountains (A-1) quadrangle, southern Alaska. Talkeetna Formation. Late Toarcian.
21	24787	53AGz40	Arthur Grantz and L. F. Fay, 1953. About 0.6 mile (1 km) northwest of point where Camp Creek crosses Glenn Highway, lat 61°50'53" N., long 147°24'43" W., Anchorage (D-2) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Early Toarcian.
21	25316	54AGz48A	Arthur Grantz and L. F. Fay, 1954. Lat 62°00'42" N., long 147°27'49" W., Talkeetna Mountains (A-1) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation, upper part. Late Toarcian.
21	25317	54AGz49A	Arthur Grantz and L. F. Fay, 1954. Lat 62°00'06" N., long 147°31'30" W., Talkeetna Mountains (A-2) quadrangle, Talkeetna Mountains, southern Alaska. Upper part of Talkeetna Formation. Late Toarcian.
21	25318	54AGz50	Arthur Grantz and L. F. Fay, 1954. Lat 62°00'10" N., long 147°32'33" W., Talkeetna Mountains (A-2) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Toarcian.
21	25319	54AGz51	Arthur Grantz and L. F. Fay, 1954. Lat 62°00'00 ½" N., long 147°32'45" W., Talkeetna Mountains (A-2) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Toarcian.
21	25342	54AFy24	L. F. Fay, 1954. Lat 62°02'48" N., long 147°29'34" W., Talkeetna (A-1) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Toarcian.
21	25359	54AFy130	L. F. Fay, 1954. Lat 62°02'40" N., long 147°21'36" W., Talkeetna Mountains (A-1) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Toarcian.
21	25940	55AGz228a	Arthur Grantz, 1955. Lat 62°01'37" N., long 147°17'53" W., Talkeetna Mountains (A-1) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Toarcian.
21	25941	55AGz294	Arthur Grantz, 1955. Lat 61°51'51" N., long 147°21'22" W., at east end of Sheep Mountain. Anchorage (D-1) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation, upper part. Late Pliensbachian.
21	27508	6-10BB	General Petroleum Corp. 1959. Near head of McDougall Creek. Coordinates 15.9-0.37. Talkeetna Mountains (A-2) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Toarcian.
21	29449	66AGz31A	Arthur Grantz, 1966. Same place and stratigraphic position as USGS Mesozoic loc. 24108.
21	29450	66AGz31B	Arthur Grantz, 1966. On south tributary of Squaw Creek, 1.95 miles (3.1 km) southeast of summit of Gunsight Mountain, near USGS Mesozoic loc. 24107. Anchorage (D-2) quadrangle, Talkeetna

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
			Mountains, southern Alaska. Talkeetna Formation. Late Pliensbachian.
22	M6171	73APr85A	George Plafker, 1973. On south-flowing branch of Doone Creek, 2,000 ft (610 m) west of SE. cor. sec. 25, T. 21 N., R. 5 E., Anchorage (D-4) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Early Pliensbachian.
22	6706	Aug. 30, 1910	A. H. Brooks, 1910. Creek entering Chickaloon River from west 1 mile (1.6 km) above Government Bridge at altitude of 2,000 ft (610 m). Coordinates 3.0-9.0. Anchorage D-4 quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Sinemurian.
22	6797	Aug. 30, 1910	G. C. Martin, 1910. Doone Creek at mouth of gulch. Anchorage (D-4) quadrangle, Talkeetna Mountains, southern Alaska. Float from Talkeetna Formation. Early Pliensbachian.
22	27586	13177-2	Standard Oil of California. Carbon Creek, 2.3 miles (3.7 km) S. 7° E. of its mouth near center of SE¼ sec. 12, T. 19 N., R. 5 E., Anchorage (C-4) quadrangle, coordinates 1.4-17.3., Chugach Mountains, southern Alaska. Talkeetna Formation. Late Pliensbachian.
22	29198	SUS62	British Petroleum Expl. (Alaska) Co. From saddle on south side of Kings Mountain 2 ½ miles (4 km) S. 15° E. of VABM 757 on the Glenn Highway. Anchorage (C-5) quadrangle. Chugach Mountains, southern Alaska. Talkeetna Formation. Early Toarcian.
23	26772	57AGz30	Arthur Grantz, 1957. Lat 62°59'58" N., long 147°57'48" W., Coordinates 1.20-17.23. Talkeetna Mountains (A-2) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Sinemurian.
23	28661	62AGz84	Arthur Grantz, 1962, on Nowhere Creek. Lat 62°13'29" N., long 148°01'27" W., Coordinates 15.35-15.63. Talkeetna Mountains (A-3) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Sinemurian.
23	28662	62AGz84A	Arthur Grantz, 1962. On Nowhere Creek, lat 62°13'30" N., long 148°01'18" W., Coordinates 15.43-15.65., Talkeetna Mountains (A-3) quadrangle, Talkeetna Mountains, southern Alaska. Late Sinemurian.
23	28663	62AGz84B	Arthur Grantz, 1962. Talus and stream float, lat 62°13'30" N., long 148°01'12" W., Coordinates 15.5-15.65., Talkeetna Mountains (A-3) quadrangle, Talkeetna Mountains, southern Alaska. Talkeetna Formation. Late Sinemurian.
24	16229	32AT1	Elmer Bedager, 1932. Head of an eastern tributary of Partin Creek that heads against Little Shotgun Creek on west side of Chulitna River valley north of Eldridge Glacier. Probably same place as USGS Mesozoic loc. 31262 in Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Late early to early middle Sinemurian.
24	31260	F6-J-27	N. J. Silberling and D. L. Jones, 1976. On divide between Ohio and Copeland Creeks; 5,800 ft (1,768 m) N. 30° E. of VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Late early to early middle Sinemurian.
24	31261	F6-J-31	N. J. Silberling and D. L. Jones, 1976. On divide between Ohio and Copeland Creeks; 6,800 ft (2,073 m) N. 39° E. from VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Latest Sinemurian.
24	31262	F6-S-322	N. J. Silberling and D. L. Jones, 1976. Divide between Partin and Little Shotgun Creeks in upper part of small drainage into Partin Creek, 39,850 ft (12,147 m) S. 31.5° W. from VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska.

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
			Unnamed calcareous sandstone. Late early to early middle Sinemurian.
24	31263	F6-S-325	N. J. Silberling and D. L. Jones, 1976. Divide between Ohio and Shotgun Creeks 19,800 ft (6,035 m) S. 38° W. from VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Late early to early middle Sinemurian.
24	31264	F6-S-331	N. J. Silberling and D. L. Jones, 1976. Divide between Shotgun and Little Shotgun Creeks, 28,800 ft (8778 m) S. 34° W. from VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Earliest Sinemurian.
24	31265	F6-S-332	N. J. Silberling and D. L. Jones, 1976. Divide between Shotgun and Little Shotgun Creeks, 31,000 ft (9,449 m) S. 33° W. from VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Late early to early middle Sinemurian.
24	31266	F6-S-333	N. J. Silberling and D. L. Jones, 1976. Crest of ridge on east side of upper Long Creek, 32,700 ft (9,967 m) N. 6° E. from VABM Copeland, Healy (A-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Earliest Sinemurian.
25	30907	75AR38A	R. L. Dettnerman and D. L. Jones, 1975. Altitude 5,850 ft (1,768 m), 3.5 mi (5.6 km) west of Shellabarger Pass, sec. 12, T. 28 N., R. 20 W., lat 62°31'49" N., long 152°53'02" W., Talkeetna (C-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds 150 ft (45.7 m) below lowest <i>Buchia</i> -bearing limestone. Sinemurian.
25	30908	75AR39	D. L. Jones, 1975. Altitude 6,100 ft (1,859 m), 5 miles (8 km) southwest of Shellabarger Pass, lat 62°30'20" N., long 152°55'59" W., Talkeetna (C-6) quadrangle, south-central part of Alaska Range, southern Alaska. Unnamed beds. Early Sinemurian.
25	31270	(76 AR17)	Bruce Reed and R. L. Dettnerman, 1976. Float from 100 to 150 ft (30 to 46 m) of siltstone and shale exposed on north side of ridge about 1 mile (1.6 km) northwest of Tatina River in south-central part of SW¼SE¼ sec. 25, T. 28 N., R. 20 W., lat 62°28'51" N., long 152°53'36" W., Talkeetna (B-6) quadrangle, south-central part of Alaska Range, southern Alaska. Early Sinemurian.
26	2978	905	T. W. Stanton and G. C. Martin, 1904. Point Naskowhak, at entrance to Seldovia Bay, Seldovia (B-5) quadrangle, on Kenai Peninsula, southern Alaska. Unnamed beds. Early Sinemurian.
26	2979	906	T. W. Stanton and G. C. Martin, 1904. Three-fourths of a mile (1.2 km) west of entrance to Seldovia Bay, Seldovia (B-5) quadrangle on Kenai Peninsula, southern Alaska. Unnamed beds. Early Sinemurian.
26	2981	907b	T. W. Stanton and G. C. Martin, 1904. Sea cliffs 2 miles (3.2 km) west of Seldovia Bay, eastern part of sec. 34, T. 8 S., R. 15 W., Seldovia (B-5) quadrangle, on Kenai Peninsula, southern Alaska. Talkeetna Formation about 200 ft (61 m) above horizontal conglomerate. Earliest Sinemurian.
26	21242	48AI119	R. W. Imlay and D. J. Miller, 1948. Sea cliffs 2.95 miles (4.7 km) S. 67° W. of Point Naskowhak on south shore of Kachemak Bay, probably NE. cor. sec. 5, T. 9 S., R. 15 W., Seldovia (B-5) quadrangle, on Kenai Peninsula, southern Alaska. Talkeetna Formation in gray limestone and green tuffaceous sandstone. Early Hettangian.

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
26	22664	51AGz21	Arthur Grantz, 1951. Sea cliffs 2.8 miles (4.5 km) west of Point Naskowhak on south shore of Kachemak Bay, near point on west side of sec. 34, T. 8 S., R. 16 W., Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Talkeetna Formation. Early Hettangian.
26	31128	75JK-77B	J. S. Kelley, 1976. Cliff near head of small bay southwest of Point Naskowhak in NE¼NW¼ sec. 36, T. 15 S., R. 14 W., Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Early Sinemurian.
26	31634	76JK-15P	J. S. Kelley, 1976. Sea Cliffs, 1.2 miles (1.9 km) 5° SE. of Pt. Naskowhak, in NW¼SW¼ sec. 31, T. 8 S., R. 14 W., lat 59°26'45" N., long 151°44'25" W., Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Unnamed beds. Early Sinemurian.
26	31637	76JK-16J	J. S. Kelley, 1976. Sea cliffs, 0.6 mile (0.94 km) 6° SE. of Pt. Naskowhak in NW¼NW¼ sec. 31, T. 8 S., R. 14 W., lat 59°26'71" N., long 151°44'26" W., Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Early Sinemurian.
26	31640	76JK-21D	J. S. Kelley, 1976. Sea cliffs, 0.19 mile (0.3 km) 14° SE. of Pt. Naskowhak, NE¼SE¼ sec. 25, T. 8 S., R. 14 W., lat 59°27'22" N., long 151°44'50" W., Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Unnamed beds. Early Sinemurian.
26	31648	76JK-38A	J. S. Kelley, 1976. Sea cliffs, 3 miles (4.8 km) 66° SW. of Pt. Naskowhak in NE¼SW¼ sec. 34, T. 8 S., R. 15 W., lat 59°26'30" N., long 151°48'86" W., Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Unnamed beds. Early Hettangian.
26	31650	H257	D. M. Hopkins, 1962. Sea cliffs just west of buried valley of Miocene age, lat 59°26.6' N., long 151°47.1' W., probably from same places as USGS Mesozoic loc. 2981, Seldovia (B-5) quadrangle, Kenai Peninsula, southern Alaska. Unnamed beds. Early Sinemurian.
27	3109	949	T. W. Stanton, 1904. Northeast shore of Puale Bay, about 1 mile (1.6 km) northwest of Chignik Point, Alaska Peninsula. A short distance southwest of locality 3110 and midway between that locality and Triassic Mesozoic loc. 19806 that contains <i>Monotis</i> . Unnamed beds. Hettangian.
27	3110	950	T. W. Stanton, 1904. Northeast shore of Puale Bay about 1 mile (1.6 km) northwest of Chignik Point and 5,400 ft (1,646 m) N. 18° W. of most easterly point of Puale Bay, Karluk (C4-C5) quadrangle, Alaska Peninsula. Unnamed beds. Hettangian.
27	10820	1-127	S. R. Capps, 1921. Northeast shore of Puale Bay about 1 ½ miles (1.8 km) northwest of Chignik Point. Contains <i>Paracoloceras</i> as at Mesozoic loc. 12396 and ROC loc. 1241 but also contains one specimen of <i>Schlotheimia</i> preserved in a different matrix and some Late Triassic mollusks. Unnamed beds. Early Sinemurian (mostly). Some late Hettangian and Late Triassic.
27	12075	1	W. R. Smith, 1923. Southwest shore Alinchak Bay, Karluk (D-4) quadrangle, Alaska Peninsula. Probably same place as locality ROC 3002 and USGS Mesozoic loc. 29268. Unnamed beds. Early and middle Hettangian.
27	12394	F16	W. R. Smith, 1924. Southwest side of Alinchak Bay at same place as Mesozoic loc. 12075. Unnamed beds, middle Hettangian.
27	12396	F18	W. R. Smith, 1924. From northeast shore of Puale Bay reportedly about 2 miles (3.2 km) northwest of Chignik Point, Karluk (C4-C5) quadrangle, Alaska Peninsula. Apparently obtained from different units as shown by presence

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
			of <i>Paracaloceras</i> as at USGS Mesozoic loc. 10820 and of <i>Waehneroceras</i> as at Mesozoic locality 3110. Unnamed beds. Early Sinemurian and late Hettangian.
27	19803	(44 AKm F75)	L. B. Kellum, S. N. Daviess, and C. M. Swinney, 1944. Northeast shore of Puale Bay 1 mile (1.6 km) northwest of Chignik Point, Alaska Peninsula. From dense limestone interbedded with limy shale, and some tuffaceous sandstone 360 ft (110 m) below base of massive tuff and agglomerate. Unnamed beds. Middle Hettangian.
27	19804	44 AKm F 76	L. B. Kellum, S. N. Daviess, and C. M. Swinney, 1944. On northeast shore of Puale Bay about 2 miles (3.2 km) north-northeast of Chignik Point and 1,000 ft (305 m) north of northernmost large waterfall, Karluk (C-4 and C-5) quadrangles, Alaska Peninsula. In dark-gray shale 30 ft (9 m) above base of Kialagvik Formation. Near middle of Toarcian.
27	21237	48AI 112	R. W. Imlay and D. J. Miller, 1948. Northeast shore of Puale Bay, Alaska Peninsula, 1.63 miles (2.6 km) N. 22° W. of Chignik Point. Float probably from light-gray sandstone exposed in cliffs about 340 ft (103 m) below base of Kialagvik Formation. Unnamed beds. Early to early late Sinemurian.
27	25694	55AKe6	A. S. Keller, 1955. Southwest side of Alinchak Bay in limestone overlying Triassic beds. Probably same place as 12075, 12394, and 25694. Unnamed beds. Middle Hettangian.
27	29267	65AP 175	Marvin Mangus, 1965. Northeast shore of Puale Bay about 1 mile (1.6 km) northwest of Chignik Point, Alaska Peninsula. Probably from same place as Mesozoic locs. 3110 and 19803. Unnamed beds. Middle Hettangian.
27	29268	65AMe55	G. W. Moore, 1965. Southwest shore of Alinchak Bay, lat 57°45.8' N., long 155°18.9' W., Karluk (D-4) quadrangle, Alaska Peninsula. Unnamed beds. Middle Hettangian.
27	M1738	ALp701M	British Petroleum, Inc., 1962. From or near the same place as Mesozoic locs. 3110 and 19803, 3½ miles (5.6 km) west of Cape Kekurnoi, lat 57°43.4' N., long 155°23.5' W., east shore of Puale Bay, Alaska Peninsula. Unnamed beds. Early to middle Hettangian.
27	31370	77AI1	R. W. Imlay, Martha Yount, Carleen Holloway, and Fred Wilson, 1977. Northeast side of Puale Bay. Same place as USGS Mesozoic loc. 3109, about 200 ft (61 m) stratigraphically above top of <i>Monotis</i> -bearing beds, 1 miles (1.6 km) northwest of Chignik Point, Karluk (C4-C5) quadrangle, Alaska Peninsula. Unnamed beds. Hettangian.
27	31372	77AI3	R. W. Imlay, Martha Yount, Carleen Holloway, and Fred Wilson, 1977. Northeast side of Puale Bay. About 4,300 ft (1,311 m) S. 12° E. of VABM Bay 119. About 270 to 280 ft (82-86 m) stratigraphically above base of massive tuffaceous sandstone. Karluk (C4-C5) quadrangle, Alaska Peninsula. Early Sinemurian.
27	ROC 1185		W. T. Rothwell and associates, 1962. From 1,900 ft (1,494 m) N. 30° W. of VABM 96 Hike on northeast shore of Puale Bay, Alaska Peninsula. Float from or near same place as localities 3110 and 19803. Unnamed beds. Early to late Hettangian.
27	ROC 1240		W. T. Rothwell and associates, 1962. From 10 ft (3 m) stratigraphically below locality ROC 1241 and probably 270 to 280 ft (82-86 m) stratigraphically above base of tuffaceous sandstone. Unnamed beds. Early Sinemurian.
27	ROC 1241		W. T. Rothwell and associates, 1962. Northeast shore of Puale Bay. About 4,300 ft (1,311 m) S.

TABLE 7.—Description of Lower Jurassic fossil localities in Alaska—Continued

Locality No. (figs. 2-6)	USGS Mesozoic loc. No.	Collector's field No.	Collector, year of collection, description of locality, and stratigraphic assignment
			12° E. of VABM 119 Bay, Karluk (C4-C5) quadrangle, Alaska Peninsula. About 700 ft (213 m) below base of Kialagvik Formation. Unnamed beds. Early Sinemurian.
27		ROC 1282 N and S.	W. T. Rothwell and associates, 1962. Same general location as Mesozoic loc. 19804. At point about 3,000 ft (914 m) S. 12° E. of VABM 119 Bay (N=bed north of fault, S=bed south of fault). In massive sandstone about 290 ft (88 m) below the base of the Kialagvik Formation. Unnamed beds. Late early to early late Sinemurian.
27		ROC 1283	W. T. Rothwell and associates, 1962. Same description as locality ROC 1282 but 10 ft (3 m) lower in the massive sandstone. Unnamed beds. Late early to early late Sinemurian.
27		ROC 1303	W. T. Rothwell and associates, 1962. Same general location as Mesozoic loc. 19804. At point about 2,700 ft (823 m) S. 12° E. of VABM 119 Bay. From unit of massive sandstone 160 ft. (49 m) thick underlying the Kialagvik Formation. Unnamed beds. Sinemurian.
27		ROC 3002	W. T. Rothwell and associates, 1962. South end of Alinchak Bay about 2.1 mi (3.3 km), N. 35° E. of VABM 1197 Kek, Karluk (D-4) quadrangle, Alaska Peninsula. Unnamed beds. Early and middle Hettangian.
28			Richfield Oil Co., Wide Bay Test Well 1, core 5 at depth of 2,235 to 2,236 ft (681 m), at Wide Bay, Alaska Peninsula. Unnamed beds. Hettangian.

SYSTEMATIC DESCRIPTIONS

Family *PSILO CERATIDAE* Hyatt, 1867
Genus *PSILO CERAS* Hyatt, 1867*Psiloceras* cf. *P. planorbis* (J. de C. Sowerby)

Plate 1, figures 1, 2

Ten compressed molds represent a highly evolute smooth ammonite similar to *P. planorbis* (J. de C. Sowerby) (1824, pl. 448; Dean and others, 1961, pl. 63, fig. 1; Arkell and others, 1957, p. L232, fig. 258-10, a-c) of early Hettangian Age. Their occurrence with *Waehneroceras* of middle Hettangian Age in three collections implies either that some of the specimens herein compared with *P. planorbis* are of that age or that the collections were made from more than one stratigraphic unit. The presence of beds of earliest Hettangian Age in the areas where these collections were made is shown by the upward gradation of marine Upper Triassic beds into marine Lower Jurassic beds.

Figured specimens.—USNM (U.S. National Museum) 247950, 247951.

Occurrences.—Upper member of McCarthy Formation at USGS Mesozoic loc. 29891 in the Wrangell Mountains; unnamed beds on the small peninsula between Puale Bay and Alinchak Bay, Alaska Peninsula, at USGS Mesozoic locs. 12075 and M1738 and ROC (Richfield Oil Co.) localities 1185 and 3002.

Subgenus **FRANZICERAS** Buckman, 1923**Psiloceras** (*Franziceras*) cf. *P. (F.) ruidum* (Buckman)

Plate 1, figures 12-14, 18-24

Twelve external molds of immature ammonites are fairly evolute, have a subovate whorl section that is a little higher than wide, an umbilicus that represents about 50 percent of the diameter, and rather sharp, widely spaced ribs. These ribs begin near the umbilicus, trend radially or slightly adapically on the flanks, and terminate rather abruptly a little above the middle of the flanks. In addition, faint lines that trend radially on the flanks and arch forward on the venter are preserved on shelly material that occurs on a few specimens from the Seldovia area.

The suture line is simple, has fairly narrow saddles, and does not have a strongly retracted suspensive lobe.

This species differs from plicate forms of *Psiloceras* such as *P. (P.) plicatum* (Quenstedt) (1883, pl. 1, figs. 9, 11; Donovan, 1952, pl. 22, figs. 1, 2; Lange, 1941, pl. 2, fig. 20) by having sharp ribs instead of folds. It differs from species of the subgenus *P. (Caloceras)* by being less evolute, by having a smaller umbilicus relative to its diameter, by ribs trending radially instead of projecting adorally, and by having a somewhat different sutural pattern. It appears to have somewhat weaker ribbing than the fragmentary specimens of *Psiloceras* from the northern Yukon that were described by Frebold and Poulton (1977, p. 92, 93, pl. 1 figs. 3-8). Overall it shows most resemblance to *P. (Franziceras) ruidum* Buckman (1923, pl. 423) but differs by having shorter ribs and a less retracted suspensive lobe. These differences may be related to its much smaller size.

It is associated at USGS Mesozoic loc. 29737 with small specimens of *Psiloceras*, of which some are smooth and some bear faint ribs on their innermost whorls.

Figured specimens.—USNM 247952-247955.

Occurrences.—Glenn Shale in Coleen quadrangle of east-central Alaska at USGS Mesozoic locs. 29737-29739 and 29742; unnamed beds in the Seldovia area on the Kenai Peninsula at USGS Mesozoic locs. 21242, 22664, and 31648.

Psiloceras (*Franziceras*) sp. ind.

Plate 1, figures 11, 15-17

This species is represented by two specimens. It differs from the species described herein as *P. (F.)* cf. *P. (F.) ruidum* (Buckman) by having sparser, shorter ribs that are much more prominent on the lower third of the flanks. These ribs are particularly prominent near the umbilical margin on the inner whorls of a specimen from northernmost Alaska as well as on the smaller specimen from east-central Alaska. On the outer whorl of the specimen from northernmost Alaska, the ribs fade considerably near the middle of the flanks but persist

across the venter, where they arch forward and vary from weak to fairly strong. In addition, fine closely spaced radial striae are present on some shelly material that is preserved on the inner whorls of the larger specimen. The suture line is not preserved.

Figured specimens.—USNM 247956, 247957.

Occurrences.—Glenn Shale in the Coleen quadrangle of east-central Alaska at USGS Mesozoic loc. 29742; Kingak Shale in northernmost Alaska in the South Barrow No. 12 well at the depth of 2,170.5 feet (661.4 m).

Genus **DISCAMPHICERAS** Spath, 1923**Discamphiceras** cf. *D. toxophorum* (Waehner)

Plate 1, figures 3, 4, 8-10

This species is represented by 12 specimens, of which 10 are crushed laterally. It is characterized by moderately involute coiling; an umbilicus that is about two-fifths as wide as the shell diameter; a high whorl section; a narrow, fairly sharp venter, and by moderately spaced ribs that trend radially, or incline slightly forward on the flanks, and that arch forward on the ventral area of smaller specimens. These ribs broaden and weaken ventrally and are faint or absent on venters of medium to large specimens. On the largest specimen the adapical half bears widely spaced ribs that fade out ventrally, whereas the adoral half is completely smooth. The suture line is not preserved.

This species closely resembles *D. toxophorum* Waehner (1884, p. 109, pl. 24, figs. 5a,b to 7a-c) in most respects but has somewhat sparser ribbing. The small specimens differ from comparable specimens of *D. kammerkahrense* (Gümbel) (Waehner, 1884, p. 113, pl. 24, figs. 3a-c, 4a-d) and from *D. calcimontanum* (Waehner) (1884, p. 112, pl. 24, figs. 1a-c, 2a,b) by having a wider umbilicus and sharper, sparser ribbing. The ribbing on the outer whorl of the largest Alaskan specimens is somewhat similar to that on the largest specimens of *D. kammerkahrense* (Gümbel) (Waehner, 1884, pl. 25, figs. 1a-c) and of *D. calcimontanum* (Waehner) (1884, pl. 24, figs. 1a,b).

Assignment of these Alaskan specimens to *Discamphiceras* rather than *Waehneroceras* is favored by their moderately involute coiling, by their ribs fading on the venter of small specimens, and by their adult outer whorl being fairly smooth.

Figured specimens.—USNM 247958-247961.

Occurrences.—Unnamed beds on peninsula between Puale Bay and Alinchak Bay on Alaska Peninsula at USGS Mesozoic locs. 3110, 25694, M1738, and ROC loc. 3002.

Discamphiceras sp.

Plate 1, figures 5-7

Seven laterally crushed molds differ from the Alaskan specimens herein compared with *Discamphiceras tox-*

ophorum (Waehner) by having weaker and more closely spaced ribs. In that respect they resemble a small specimen of *D. calcimontanum* (Waehner) (1884, p. 112, pl. 24, figs. 2a, b.) Their ribs are much weaker and do not extend as far ventrally as on the type specimens of *D. toxophorum* (Waehner) (1884, p. 109, pl. 24, figs. 5a-c to 7a-c).

Figured specimens.—USNM 247962.

Occurrence.—Upper member of the McCarthy Formation in the Wrangell Mountains at USGS Mesozoic loc. 29890.

Genus LAQUEOCERAS Lange, 1925

***Laqueoceras* cf. *L. sublaqueus* (Waehner)**

Plate 3, figure 13

This species is represented by one laterally crushed specimen that includes a large internal mold and a small, shell-bearing fragment. The specimen is highly evolute, consists of many whorls, and bears fine radial, rather closely spaced ribs. These ribs are fairly distinct on the inner whorls at diameters as large as 75 mm but gradually become faint at greater diameters and are barely evident on the outermost whorl. The venter is poorly exposed but apparently does not bear a keel.

The specimen is similar to *L. sublaqueus* (Waehner) (1886, p. 142, pl. 15, fig. 2; pl. 16, fig. 10; pl. 30, fig. 4; Donovan, 1952, p. 641, 642) in size and in the strength, closeness, and persistence of its ribbing.

Figured specimen.—USNM 247963

Occurrence.—Unnamed Lower Jurassic beds on south side of Alinchak Bay on Alaska Peninsula at ROC loc. 3002.

Genus WAEHNEROCERAS Hyatt, 1889

***Waehneroceras* cf. *W. tenerum* (Neumayr)**

Plate 2, figures 1-6

cf. *Aegoceras tenerum* Neumayr, 1879, K. K. Geol. Reichsanst., Abh. v. 7, p. 31, pl. 3, figs. 4, 5.

cf. *Waehneroceras tenerum* (Neumayr). Arkell and others, 1957, Treatise on invertebrate paleontology, Part L, Mollusca 4, p. L235, fig. 260-1a,b.

This species is represented by 30 crushed molds. It has highly evolute coiling, a wide umbilicus, and an ovate whorl section. Its ribs are widely spaced, trend nearly radially on the flanks, become stronger ventrally, project forward and become much weaker on the margin of the venter, and are reduced in strength along the midline of the venter.

This species differs from the specimens herein compared with *W. portlocki* (Wright) by having a lower whorl section and much stronger and sparser ribbing. Its features are essentially the same as those of *W. tenerum* (Neumayr) (Arkell and others, 1957, p. L235, fig. 260-1a,b). It is associated at four localities in Alaska with *Waehneroceras* cf. *W. portlocki* (Wright) and at three localities with *Discamphiceras*.

Figured specimens.—USNM 247964-247966.

Occurrences.—Unnamed beds on south side of Alinchak Bay on Alaska Peninsula at USGS Mesozoic locs. 12075, 12394, 25694, 29268 and ROC loc. 3002; unnamed beds on east side of Puale Bay on Alaska Peninsula at USGS Mesozoic locs. 3109, 3110, 19803, 31370, and ROC loc. 1185.

***Waehneroceras* cf. *W. portlocki* (Wright)**

Plate 2, figures 7, 10-15

cf. *Aegoceras portlocki* Wright, 1881, 1882, London, Palaeontographical Soc., l. pl. 48, figs. 4, 5 (1881), p. 372 (1882).

cf. *Aegoceras extracostatum* Waehner, 1882, Beitr. Pal. Geol. Osterr.-Ungarns v. Orient, v. 2, pt. 3, p. 74, pl. 14, fig. 1.

cf. *Macrogrammites grammicus* Buckman, 1928, Type ammonites, v. 7, pl. 761 a,b.

cf. *Schlotheimia* (*Waehneroceras*) *portlocki* (Wright). Donovan, 1952, Annals and Mag. Nat. History, ser. 12, v. 5, p. 646, pl. 22, figs. 5a,b.

cf. *Schlotheimia* (*Waehneroceras*) *portlocki* (Wright). Dean, Donovan, and Howarth, 1961, British Mus. (Nat. History) Bull., v. 4, no. 10, p. 445, pl. 63, figs. 4a,b.

This species is represented by 33 internal and external molds, of which most are laterally crushed. These specimens have highly evolute coiling, a wide umbilicus, and a compressed whorl section. Their ribs trend nearly radially on the flanks, bend forward slightly on the ventral margin, are rather closely spaced on septate whorls and become more widely spaced on the incomplete body chamber. The venter is smooth along its midline as shown on several specimens. The suture line is not exposed.

In coiling and ribbing this species is essentially identical with the specimen of *Waehneroceras portlocki* (Wright) figured by Dean, Donovan, and Howarth (1961, pl. 63, figs. 4a,b) and probably represents the same species. None of the Alaska specimens, however, include the large outer whorl as figured by Wright (1881, pl. 48, figs. 4, 5) and Buckman (1928, pl. 761a,b). *W. curvior-natum* (Waehner) (1882, p. 75, pl. 16, figs. 2-4) appears to have slightly coarser ribbing on its inner whorls and develops a smooth venter on its outer whorls.

Figured specimens.—USNM 247967-247972.

Occurrences.—Unnamed beds on south side of Alinchak Bay on Alaska Peninsula at USGS Mesozoic locs. 12075, 12394, 25694, and 29268, and ROC loc. 3002; unnamed beds on east side of Puale Bay on Alaska Peninsula at Mesozoic loc. 3110, 29267 and 31370; unnamed beds in Richfield Oil Co.-Wide Bay test well No. 1, core no. 5 at depth of 2235 to 36 ft (681 m) on north side of Wide Bay, Alaska Peninsula.

***Waehneroceras*? sp.**

Plate 2, figures 8, 9

Waehneroceras is possibly represented in the subsurface of northern Alaska by one small fragment of a whorl in which only the venter and upper part of the

flank are preserved. The specimen is characterized by simple ribs that project strongly forward on the venter and that are only slightly reduced in strength on the venter. These features are similar to those on immature specimens of *Waehneroceras tenerum* (Neumayr) (Arkell and others, 1957, p. L235, figs. 1a,b).

Figured specimen.—USNM 247973.

Occurrence.—Kingak Shale in South Barrow test well No. 12 at depth of 2181.5 ft. (665 m) in northern Alaska. This occurrence is 11 ft (3.3 m) below that of *Psiloceras* (*Franziceras*).

Family SCHLOTHEIMIIDAE Spath, 1923
Genus SCHLOTHEIMIA Bayle, 1878

***Schlotheimia* sp.**

Plate 2, figures 16, 17

One specimen is moderately evolute. Its whorl section is a little higher than it is wide. The umbilical width is 40 percent of the diameter. Its ribs are simple, sharp, incline forward on the flanks, become a little stronger ventrally, arch forward on the venter, and are only slightly reduced in strength along the midline of the venter.

The ribbing on this specimen resembles that on the inner whorls of *S. montana* (Waehner) (1886, p. 165, pl. 19, fig. 1; pl. 20, fig. 1) and of *S. donar* (Waehner) (1886, p. 173, pl. 21, figs. 2, 5).

Figured specimens.—USNM 247974.

Occurrence.—Unnamed beds in the Puale Bay area on Alaska Peninsula at USGS Mesozoic loc. 10820.

Genus BADOUIA Guex and Taylor, 1976

***Badouxia canadensis* (Frebold)**

Plate 2, figures 18–21, 24–28

Psiloceras canadense Frebold, 1951, Canada Geol. Survey Bull. 18, p. 3, pl. 1, figs. 1–6, pl. 2, fig. 1a–c; pl. 3, fig. 1.

Psiloceras canadense Frebold, 1964a, Canada Geol. Survey Bull. 116, p. 6, pl. 1, figs. 1–5b.

Psiloceras canadense Frebold, 1967b, Canada Geol. Survey Bull. 158, p. 18, pl. 1, figs. 1a,b, 2a,b, 3a–c.

Badouxia canadensis (Frebold). Guex and Taylor, 1976, Eclogae Geol. Helvetiae, v. 69, no. 2, p. 525.

This species is represented in Alaska by 10 fairly small specimens. Most of these (pl. 2, figs. 18, 24, 28) bear rather closely spaced ribbing comparable with that on certain specimens illustrated by Frebold (1951, pl. 1, fig. 5a; 1967b, pl. 1, fig. 2a). A few specimens (pl. 2, figs. 20, 26, 27) bear sparser ribbing, as does another specimen illustrated by Frebold (1951, pl. 1, figs. 1e,d; 1967b, pl. 1 fig. 3a).

The species has ovate whorls that are higher than they are wide, embrace half or more of the preceding whorl, and are rounded ventrally. Its smaller whorls bear straight, fairly sharp, forwardly inclined ribs that extend across the lower two-thirds of the flanks and then either fade out rather abruptly or pass into very weak secondary ribs. Its larger whorls are marked only by

strong forwardly inclined ribs that fade out rather abruptly at the base of the upper third of the flanks. Its assignment to the genus *Psiloceras* is questioned because it is more involute and more strongly ribbed than is typical of that genus. Its ribbing is much stronger on the lower part of the flanks than in *Arc-toasteroceras jeletzkyi* Frebold.

Hypotypes.—USNM 247975–247977 and 248062.

Occurrences.—Unnamed beds in the Healy (A–6) quadrangle in Alaska Range at USGS Mesozoic loc. 31266; unnamed beds at Puale Bay in Alaska Peninsula at ROC loc. 1240, and USGS Mesozoic loc. 31372.

***Badouxia columbiae* (Frebold)**

Plate 2, figures 22, 23

Psiloceras (*Curvicerias*) *columbiae* Frebold, 1967b, Canada Geol.

Survey Bull. 158, p. 20, pl. 1, figs. 10a–c; pl. 2, figs. 1–5;

p. 3, figs. 2a–c.

Schlotheimia? sp. indet. Frebold, 1951, p. 7, pl. 4, figs. 2–4.

One fragmentary ammonite bears simple ribs that incline forward on the flanks as do those on *B. canadensis* (Frebold), but that continue across the venter where they arch strongly forward as in *B. columbiae* (Frebold).

Figured specimen.—USNM 247978.

Occurrence.—Unnamed beds in the Healy (A–6) quadrangle, Alaska Range at USGS Mesozoic loc. 31264.

Genus CHARMASSEICERAS Spath, 1924

***Charmasseiceras* cf. *C. marmoreum* (Oppel)**

Plate 3, figures 1–3

This species is represented by three fragmentary molds of immature specimens. The best preserved internal mold shows that the species is fairly involute, that its ribbing changes during growth from fairly fine to moderately strong, that many ribs fork low on the flanks, and that all ribs curve forward on the upper parts of the flanks. One external mold bears similar forwardly curved ribs that terminate ventrally in shelly material whose smoothness suggest a ventral band.

These specimens are closely similar in involution and ribbing to the small specimen of *Charmasseiceras marmoreum* (Oppel) figured by Waehner (1886, pl. 22, figs. 2–4) and by Frebold (1967b, pl. 3, figs. 1c and 1d; pl. 4, figs. 2a–c).

Figured specimens.—USNM 247979.

Occurrence.—Kingak Shale in northern Alaska in the South Barrow test well No. 12 at depths of 2061.5 and 2068 feet (628 and 630 m). Unnamed beds in Seldovia area in northern Alaska at USGS Mesozoic locs. 2979 and 31637.

***Charmasseiceras* sp.**

Plate 3, figures 4–10

Three small specimens are characterized by a subquadrate whorl section that is much higher than it is

wide; by a flattened venter; by high, sharp, widely spaced ribs that incline forward strongly on the upper parts of the flanks; and by prominent forwardly inclined ventral swellings that on the largest specimen are joined across the venter by low weak chevron-shaped ribs.

These specimens differ from immature specimens of *C. marmoreum* (Oppel) (Waehner, 1886, pl. 22 figs. 2-6) by having slightly coarser ribs of which only a few fork low on the flanks. In whorl shape and ribbing, the specimens show more resemblance to some small specimens that were figured as *C. charmassei* (d'Orbigny) (1844, pl. 91, figs. 1,2). They also resemble some small specimens of *C. posttaurinum* (Waehner) (1886, pl. 23, figs. 17a-c) in ribbing but are much more compressed.

Figured specimens.—USNM 247980, 247981.

Occurrences.—Kingak Shale in northern Alaska in the South Barrow test well No. 12 at depth of 2,056 feet (627 m) and in the South Barrow test well No. 3 at depths of 2,412 and 2,419.5 feet (735 and 737 m).

Family ARIETITIDAE Hyatt, 1874
Subfamily ARIETITINAE Hyatt, 1874
Genus ARIETITES Waagen, 1869

Arietites cf. *A. bucklandi* (J. Sowerby)

Plate 4, figures 11-13

"*Arietites*" cf. "*A. bucklandi*" (J. Sowerby). Imlay, 1955, U.S. Geol. Survey Prof. Paper 274-D, p. 87, pl. 10, figs. 7, 8.

The assignment of this species to *Arietites* was accepted by Arkell (1956, p. 530). No other specimen of the genus has yet been found in Alaska.

Figured specimen.—USNM 108778.

Occurrence.—Kingak Shale in northern Alaska in the Avak test well No. 1 at depth of 1836 ft (560 m).

Genus CORONICERAS Hyatt, 1867

Coroniceras sp. A

Plate 4, figures 1-5

This species, represented only by fragmentary material, is similar in appearance to *Coroniceras multicoatum* (J. de C. Sowerby) (1824, v. 5, p. 75, pl. 454; Reynes, 1879, pl. 25, figs. 1-2; Guérin-Franchette, 1966, p. 141, pls. 29-32). The specimens from Alaska are fairly evolute. Their whorls are higher than they are wide. Their ribs are fairly strong, radial on the flanks, bend forward on the ventral margin, and bear ventrolateral tubercles. Their keel is fairly strong and on internal molds is bordered by shallow furrows.

Figured specimen.—USNM 247982.

Occurrence.—Unnamed Lower Jurassic beds in Talkeetna (C-6) quadrangle, central part of the Alaska Range at USGS Mesozoic loc. 30908.

Coroniceras sp. B

Plate 4, figures 6-10

This species is represented by 12 poorly preserved internal and external molds. Its appearance is similar to that of the specimens herein described as *Coroniceras* sp. A, but differs by having ribs curving forward much more strongly on the upper part of its flanks. In that respect it shows more resemblance to *C. haueri* (Waehner) (1886, p. 38, (127), pl. 19, fig. 1a,b; pl. 20, fig. 2a,b). Its ribbing also resembles some species of *Caenoceras* (J. de C. Sowerby, 1824, pl. 452, fig. 1; Wright, 1879, pl. 12, fig. 1; Arkell, 1956, pl. 31, fig. 1; Dean and others, 1961, pl. 66, figs. 1,2; Guérin-Franchette, 1966, pls. 204, 208). It differs from that genus, however, in that its keel is bordered by very weak instead of deep furrows.

Figured specimens.—USNM 247983, 247984.

Occurrences.—Kingak Shale in northern Alaska in the South Barrow test well No. 12 at depth of 1987.4 to 1987.6 ft (606 m).

Coroniceras sp. C

Plate 4, figure 14

Three small fragmentary molds of ammonites from the subsurface of northern Alaska possibly represent the inner whorls of the species described herein as *Coroniceras* sp. A. They appear, however, to have slightly sparser ribs that bear stronger tubercles on the ventrolateral margin. The keel is fairly low and is not bordered by furrows.

Figured specimens.—USNM 247985.

Occurrence.—Kingak Shale in northern Alaska in the South Barrow test well No. 3 at depth of 2470 ft (753 m).

Subgenus PARACORONICERAS Spath, 1922

Coroniceras (*Paracorniceras*) sp.

Plate 4, figures 23-25

One species, represented by a crushed fragment of an outer whorl and by one internal mold, bears prominent widely spaced ribs that trend radially on the flanks and curve forward on the ventral margin. Its keel is fairly high and is not bordered by furrows. Its whorl section is considerably higher than wide.

This specimen resembles the adoral end of the outermost preserved whorl of the coarsely ribbed holotype of *Coroniceras* (*Paracorniceras*) *charlesi* Donovan (1955, p. 12, 28; Reynes, 1879, pl. 16, figs. 1, 2; Guérin-Franchette, 1966, p. 153, pl. 38).

Figured specimen.—USNM 247986.

Occurrence.—Unnamed beds in the Seldovia area on Kenai Peninsula at USGS Mesozoic locs. 31128 and 31650.

Genus *ARNIOCERAS* Hyatt, 1867*Arnioceras* cf. *A. densicosta* (Quenstedt)

Plate 5, figures 9-11, 16-24

This species is represented by 38 specimens, of which most are molds; but shelly material is preserved on some small specimens obtained from a well core in northern Alaska. It has highly evolute coiling. Its ribs are sharp, fairly closely spaced, trend radially on the flanks, bend forward on the margins of the venter, and then fade out on the venter. Its innermost whorls are smooth to a diameter of 15 to 20 mm. Its venter bears a keel that is bordered by distinct furrows on the internal mold and by weak furrows where shelly material is preserved. These features are essentially identical with those on the lectotype of *Arnioceras densicosta* (Quenstedt) (1858, pl. 7, fig. 7; 1884, p. 100, pl. 13, fig. 7; Guérin-Franiatte, 1966, pl. 142, figs. 1a,b) and on other comparable illustrated specimens (Guérin-Franiatte, 1966, p. 265, pl. 142, figs. 2, 3; Reynes, 1879, pl. 14, figs. 5, 6).

The suture line is similar to that on *A. densicosta* Quenstedt (1884, pl. 13, fig. 7). The first lateral lobe is broad, bifid, and a little deeper than the ventral lobe. The first lateral saddle is bifid.

Figured specimens.—USNM 247987-247989.

Occurrences.—Unnamed beds in the Puale Bay area on Alaska Peninsula at USGS Mesozoic loc. 21237, and ROC locs. 1282N, ROC 1282S, and ROC 1283A; unnamed beds in the Healy (A-6) quadrangle, Chulitna Valley area in the Alaska Range at USGS Mesozoic locs. 16229, 31260, 31262, 31263, 31265. Upper member of McCarthy Formation in the Wrangell Mountains at USGS Mesozoic locs. 28535, 28538, and 30140.

Arnioceras sp. juv.

Plate 5, figures 5, 6, 12-15

Arnioceras is represented in northern Alaska by many small immature specimens obtained from a well core in the Point Barrow area. These specimens are nearly identical in coiling, whorl shape, and ornamentation with the inner whorls of the specimens herein described as *A. cf. A. densicosta* (Quenstedt). They differ by their innermost whorls, being smooth only to diameters of 8 to 15 mm.

Figured specimens.—USNM 247990.

Occurrence.—Kingak Shale in northern Alaska in the South Barrow test well No. 12 at depth of 2056 ft (626.7 m)

Genus *PARACALOCERAS* Spath, 1923*Paracaloceras rursicostatum* Frebold

Plate 6, figures 1-11

Paracaloceras rursicostatum Frebold, 1967b, Canada Geol. Survey Bull. 158, p. 26, pl. 7, figs. 1a-c, 2a-c; pl. 9, fig. 1.

Ten laterally crushed internal molds are characterized

by highly evolute coiling and by strong, rather widely spaced ribs that curve adapically on most flanks, curve adorally and become stronger on the ventral margin, and then terminate abruptly. The keel on the mold is fairly low and is bounded laterally by two furrows that are bounded by ridges. The furrows and the lateral ridges are very weak on the small specimens but are strong on the larger specimens (pl. 6, figs. 1, 5, 11). On the least crushed specimen (pl. 6, figs. 7, 11), the whorl section is nearly as wide as it is high. During growth, the adapical arching of the ribs becomes more pronounced.

All the specimens from Alaska have sparser and somewhat coarser ribbing than those described as *Paracaloceras* cf. *P. coregonense* (Sowerby) by Frebold (1951, p. 7, pl. 5, figs. 1-6; pl. 6, fig. 1; 1967b, p. 24, pl. 7, figs. 3-7). Their ribbing is similar, however, to some specimens illustrated by Waehner (1888, pl. 21, figs. 1-2; pl. 23, fig. 4) as *Arietites coregonensis* Sowerby. They differ mainly by having ribs bending adapically more strongly on their largest whorls, a feature characteristic of *P. rursicostatum* Frebold.

Hypotypes.—USNM 247991-247997 and 248070.

Occurrences.—Unnamed beds in the Puale Bay area on Alaska Peninsula at USGS Mesozoic locs. 10820, 12396, 31372, and ROC loc. 1241; unnamed beds in the Seldovia area on Kenai Peninsula at Mesozoic loc. 2981; unnamed beds in the Healy (A-6) quadrangle in Alaska Range at Mesozoic loc. 31266.

Genus *ARCTOASTEROCERAS* Frebold, 1960*Arctoasteroceras jeletzkyi* Frebold

Plate 5, figures 1-4

Arctoasteroceras jeletzkyi Frebold, 1960, Canada Geol. Survey Bull. 59, p. 14, pl. 2, figs. 1-5; pl. 3, figs. 1-3.

Arctoasteroceras jeletzkyi Frebold, 1964b, Canada Geol. Survey Bull. 63-4, p. 5, pl. 2, figs. 1, 2.

This species is represented in Alaska by two specimens, of which the larger closely resembles the holotype (Frebold, 1960, pl. 2, fig. 1a,b) and the smaller resembles a paratype (Frebold, 1960, pl. 3, fig. 3a,b). Resemblances include a moderately involute shell, an ovate whorl section, gently convex flanks, a low blunt keel, and moderately spaced ribs that incline forward on the lower two-thirds of the flanks and then bend forward and become faint on the upper third and on the venter. The only difference is the presence of faint sulci bordering the keel. The suture line is poorly preserved on the Alaska specimens.

Hypotypes.—USNM 247998, 247999.

Occurrences.—Unnamed beds in the Healy (A-6) quadrangle in Alaska Range at USGS Mesozoic loc. 31261.

Family ECHIOCERATIDAE Buckman, 1913
Genus PALTECHIOCERAS Buckman, 1924

Paltechioceras cf. *P. harbledownense* (Crickmay)

Plate 4, figures, 15–22

cf. *Melanhippites harbledownensis* Crickmay, 1928, California Univ. Pub. Dept. Geol. Sci. Bull., v. 18, no. 2, p. 61, pl. 3, pl. 4a–d.

Ten specimens represent a species characterized by highly evolute coiling, an elliptical whorl section that is much higher than it is wide, a single keel that is not bordered by furrows, and moderately spaced, simple, slightly flexuous ribs. On the smallest specimens up to a diameter of about 27 mm, the ribs trend radially or curve adapically on the flanks, become stronger ventrally, and terminate abruptly on the ventral margin (pl. 4, figs. 18–21). During further growth, as shown on the larger specimens, the ribs gradually become nearly radial, or incline slightly adorally on the flanks, and their ventral ends curve a little adorally.

These specimens closely resemble the paratypes of *Melanhippites harbledownensis* Crickmay (1928, pl. 4, figs. a–d) and likewise are associated with *Entolium? semiplicatum* (Hyatt) (equals *E. balteatum* Crickmay, 1928, pl. 4 figs. e–g). The ribbing on the Alaskan specimens is slightly sparser than on the holotype of *Paltechioceras aplanatum* Hyatt (Buckman, 1924, pl. 482) but nearly identical with that on the same species as figured by Getty (1973, p. 20, pl. 4, fig. 1a,b). The ribbing is likewise similar to that on *Vermiceras bavaricum mexicanum* Erben (1956, p. 207, pl. 36, figs. 507), which is assigned to *Paltechioceras* by Hallam (1965, p. 1493).

Figured specimens.—USNM 248000–248002.

Occurrences.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic locs. 28661 and 28663; upper member of McCarthy Formation in the Wrangell Mountains at USGS Mesozoic locs. 30139 and 31174.

Subgenus ORTHECHIOCERAS Trueman and Williams, 1925

Paltechioceras (*Orthechioceras*?) sp.

Plate 3, figures 11, 12

One small specimen, consisting of five incomplete whorls, is characterized by highly evolute coiling, by a subquadrate whorl section that is slightly wider than high; by a carinate bisulcate venter whose sulci are fairly shallow; and by simple forwardly curved ribs that are rather closely spaced on the innermost whorls but become more widely spaced on the outermost whorl.

The specimen shows considerable resemblance to *P. (O.) radiatum* Trueman and Williams (1925, p. 724, pl. 2, fig. 9a,b; Getty, 1973, p. 23, pl. 5 figs. 1a,b, 2a,b), but its ribbing is denser on its four smallest whorls. It also shows some resemblance to the inner whorls of *P. (P.) elicatum* Buckman (1924, pl. 483) but differs by having shallower sulci and by ribs which become more widely spaced adorally.

The specimen is assigned to the subgenus *Orthechioceras* rather than the subgenus *Paltechioceras* because of its fairly shallow sulci and because its ribs become widely spaced during growth. Its poor preservation, however, does not warrant a positive subgeneric determination.

Figured specimen.—USNM 248003.

Occurrence.—Unnamed beds in the Healy (A–6) quadrangle in Alaska Range at USGS Mesozoic loc. 31261.

Family EODEROCERATIDAE Spath, 1929
Subfamily XIPHEROCERATINAE Spath, 1925
Genus CRUCILOBICERAS Buckman, 1920

Crucilobicer cf. *C. crucilobatum* Buckman

Plate 5, figures 7, 8

One laterally crushed, fairly large external mold represents a highly evolute, fairly large bituberculate ammonite that is similar in appearance to *Crucilobicer* and *Microderoceras*. Its outer two whorls bear rather weak, fairly sparse radial ribs that become stronger ventrally. The tubercles in the outer row occur near the ventral margin, are fairly prominent, and are round to slightly elongate spirally. The tubercles in the inner row occur at about the top of the lower third of the flanks, are fairly weak, are elongate radially, and appear to be a little stronger on the outermost whorl than on the next smaller whorl. The smallest whorls are much corroded.

The same species is probably represented by one small crushed mold that differs from the large specimen mainly by having more pronounced ribs and tubercles.

These specimens are similar in appearance to *C. crucilobatum* Buckman (1920, pl. 178) except for having more closely spaced ribs. Their assignment to *Crucilobicer* rather than *Microderoceras* is based on the weakness of the inner row of tubercles and the rather high ventral position of the outer row of tubercles.

In England *C. crucilobatum* Buckman occurs in the lower part of the *Echioceras raricostatum* zone (Buckman, 1920, v. 3, pl. 178).

Figured specimens.—USNM 248004, 248005.

Occurrence.—Upper member of McCarthy Formation in the Chitina Valley of Wrangell Mountains at USGS Mesozoic loc. 14472 and probably at Mesozoic loc. 14030.

Crucilobicer cf. *C. densinodulum* Buckman

Plate 7, figures 4, 5

cf. *Crucilobicer densinodulum* Buckman, 1923, Yorkshire type ammonites, v. 5, pl. 442.

cf. *C. densinodulum* Buckman. Dean, Donovan, and Howarth, 1961, p. 459, pl. 67, fig. 5.

One worn specimen is characterized by highly evolute coiling; by a quadrate whorl section that is a little wider than high; and by very widely spaced flank ribs that incline slightly adorally, are swollen radially on the lower

part of the flanks, weaken near the middle of the flanks, and terminate ventrally in prominent spirally elongate tubercles. These tubercles are well preserved at only one place.

This specimen differs from *C. densinodulum* Buckman by having weaker ribs near the middle of its flanks. *C. densinodulum* (Quenstedt) as figured by Wright (1880, pl. 39, figs. 6, 7; 1882, pl. 50, figs. 11, 12) has much stronger ribbing and a higher whorl section. The species *C. densinodulum* occurs in the lower part of the *Echioceras raricostatum* zone (Dean and others, 1961, p. 459).

Figured specimen.—USNM 248006

Occurrence.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic loc. 28661.

***Cruciloboceras* cf. *C. muticum* (d'Orbigny)**

Plate 7, figures 6–10, 12–15

This species is represented by 17 laterally compressed specimens, of which 15 are from USGS Mesozoic loc. 28661. It is characterized by highly evolute coiling; a subquadrate whorl section that is a little higher than wide; and ribs that are strong, straight, rather sparse, incline forward on the flanks, become stronger ventrally, and terminate in fairly prominent tubercles on the ventral margin. The venter is gently rounded, smooth on the inner whorls, but marked on some outer whorls by weak swellings that arch gently forward from the tubercles. Most of the specimens have rather widely spaced ribs, as in *C. muticum* (d'Orbigny) (1844, p. 274, pl. 80).

C. muticum (d'Orbigny) in Europe is recorded from the *Uptonia jamesoni* Zone at the base of the Pliensbachian and questionably from the top of the underlying *Echioceras raricostatum* Zone (Bremer, 1965, p. 155; Mouterde and Ruget, 1970, p. 50; Geczy, 1976, p. 57).

The specimens herein compared with *C. muticum* (d'Orbigny) and *C. submuticum* (Oppel) are associated with echioceratid ammonites and hence must be of latest Sinemurian Age.

Figured specimens.—USNM 248007.

Occurrences.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic locs. 28661, 28662, and 28663.

***Cruciloboceras* cf. *C. submuticum* (Oppel)**

Plate 7, figures 1–3

Six specimens differ from those herein compared with *C. muticum* (d'Orbigny) by having finer and more closely spaced ribs as in *C. submuticum* (Oppel) (Quenstedt, 1885, pl. 33, figs. 1, 9, 19, 23). As five of these specimens are associated with the specimens herein compared with *C. muticum* (d'Orbigny), they may all be variants of a single species.

C. submuticum (Oppel) in Europe is recorded from the *Uptonia jamesoni* zone at the base of the Pliensbachian (Arkell, 1956, pl. 128; Bremer, 1965, p. 156). Similar finely ribbed specimens from Hungary are recorded from the *Tragophylloceras ibex* Zone (Geczy, 1976, p. 58).

Figured specimens.—248008, 248009.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 6706 and 28661.

***Cruciloboceras* cf. *C. pacificum* Frebold**

Plate 8, figures 10–12, 15–17

cf. *Cruciloboceras pacificum* Frebold, 1970, Canadian Jour. Earth Sci., v. 7, no. 2, p. 435–437, pl. 1, figs. 4–8; pl. 2, fig. 10.

One specimen, represented by external and internal molds, is nearly identical in whorl shape, ribbing, and tuberculation with the type specimens of *Cruciloboceras pacificum* Frebold. Its cross section is quadrate, slightly higher than wide, and is broadest at the ventrolateral border. Its venter is nearly flat and its flanks are slightly convex. Its ribs on the flanks are widely spaced, fairly strong, incline slightly forward, and terminate on the ventrolateral margin in nodes that are elongate spirally. Between successive strong ribs generally occur single weaker ribs, most of which fade out on the upper parts of the flanks. The surface of the venter is somewhat irregular but is not marked by ribs.

The identification of this specimen with *Cruciloboceras pacificum* Frebold is favored not only by these features but by the fact that it is likewise associated with fairly well preserved specimens of *Tropidoceras actaeon* (d'Orbigny), as in the Queen Charlotte Islands.

Figured specimens.—USNM 248010.

Occurrence.—Near top of upper member of the McCarthy Formation in the Wrangell Mountains at USGS Mesozoic loc. 28534.

Subfamily COELO CERATIDAE Haug, 1910

Genus APODEROCERAS Buckman, 1921

***Apoderoceras* cf. *A. subtriangulare* (Young and Bird)**

Plate 8, figures 14, 18–23

cf. *Deroceras subtriangulare* (Young and Bird). Buckman, 1913, Yorkshire type ammonites, v. 2, pl. 71A, B.

cf. *Platyleuroceras*? sp. indet. Frebold, 1970, Canadian Jour. Earth Sci., v. 7, no. 2, p. 439, pl. 1, fig. 3).

This species is represented by two specimens. It is characterized by evolute coiling; a wide, depressed whorl section that is widest near the venter; a venter that is slightly arched on the smallest preserved whorls, nearly flat on the largest whorls, and smooth or faintly striate; and straight simple ribs that begin low on the umbilical wall, incline slightly adorally on the flanks, and terminate in prominent nodes on the margin of the venter. The ribs exposed in the umbilicus of the innermost preserved whorls are sharp and fairly closely

spaced, but at a diameter of about 25 mm they become much stronger and more widely spaced. The suture line is not preserved.

The largest specimen at a diameter of about 120 mm has a whorl height of about 30 mm, a whorl thickness of 46 mm, and an umbilical width of 68 mm.

The largest preserved whorl of this species resembles that of the holotype of *A. subtriangulare* (Young and Bird) (Buckman, 1913, pl. 71A,B) from England but differs by having a flatter venter and more closely spaced ribs. The smallest whorls resemble those of *Platypleuroceras*? sp. indet. of Frebold (1970, pl. 1, fig. 3) from the McConnell Creek area in western British Columbia, except possibly for an abrupt change in strength of ribbing.

Figured specimens.—USNM 248011, 248012.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 6697 and M6171.

Family OXYNOTICERATIDAE Hyatt, 1875
Genus FANNINOCERAS McLearn, 1930

Fanninoceras kunae McLearn

Plate 7, figure 11

Four immature specimens resemble *F. kunae* McLearn (1930, p. 5, pl. 2, fig. 4; 1932, pl. 9, figs. 1–6; Frebold, 1964b, pl. 9, fig. 4) in having a fairly wide umbilicus, a narrowly rounded venter, and moderately strong ribs that project forward on the flanks and become very weak on the venter.

Hypotype.—USNM 248013.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 24107, 24108, and 27586.

Fanninoceras cf. *F. carlottense* McLearn

Twelve specimens resemble *Fanninoceras carlottense* McLearn (1932, p. 76, pl. 8, figs. 9, 10) in having a tiny umbilicus, a sharp venter, very weak, widely spaced ribs on small and intermediate-sized whorls, and nearly smooth outer whorls.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 24108 and 29449.

Family POLYMORPHITIDAE Haug, 1887
Genus UPTONIA Buckman, 1898

Uptonia cf. *U. dayiceroides* Mouterde

Plate 9, figures 1–4, 8, 12–16

cf. *Uptonia dayiceroides* Mouterde, 1951, Soc. Geol. Portugal Bol., v. 9, no. 3, p. 179, pl. 1, figs. 4–6.

Uptonia cf. *U. dayiceroides* Mouterde. Frebold, 1970, Canadian Journal Earth Sci., v. 7, no. 2, p. 438, pl. 1, figs. 9a,b,c.

Eighty laterally crushed molds from the Wrangell Mountains are identical in coiling and density of ribbing with a specimen from the Queen Charlotte Islands that Frebold (1970, p. 438, pl. 1, figs. 9a,b) compared with *Uptonia dayiceroides* Mouterde.

On the small and intermediate-sized specimens from Alaska, the flank ribs are gently flexuous, become stronger ventrally, bear tubercles on the ventral margin, and then curve adorally. On the largest available specimens the flank ribs differ by not bearing tubercles. None of the specimens shows the middle part of the venter. Several associated specimens (pl. 9, figs. 2, 14) that have slightly coarser and sparser ribbing are herein interpreted as a variant.

Figured specimens.—USNM 248014, 248015.

Occurrences.—Upper member of McCarthy Formation in the Wrangell Mountains at USGS Mesozoic locs. 28671–28673 and 28675.

Uptonia cf. *U. jamesoni* (J. de C. Sowerby)

Plate 9, figure 17

One laterally crushed ammonite from northern Alaska greatly resembles a specimen of *Uptonia jamesoni* (J. de C. Sowerby) from England (Wright, 1882, p. 352, pl. 51, figs. 1–3; Dean and others, 1961, pl. 68, figs. 3a,b) in its evolute coiling and in its ribbing. Its ribs are fairly strong, incline slightly forward on the flanks, incline strongly forward on the margins of the venter, become swollen ventrally, and become much stronger adorally. Sharp ventral tubercles are not evident. Most of the venter is not exposed.

Figured specimen.—USNM 248016.

Occurrence.—Unnamed shale in northern Alaska at USGS Mesozoic loc. 29774.

Uptonia? sp.

Plate 9, figures 9–11

Six small crushed ammonites from northern Alaska are nearly identical in coiling and ornamentation with the inner whorls of the fairly large ammonite described herein (pl. 9, fig. 17) as *Uptonia* cf. *U. jamesoni* (J. de C. Sowerby). The only difference consists of the presence of very small, sharp tubercles on some ribs at the ventral margin. Such close resemblances suggest that the small ammonites are probably immature forms of the same species as the large ammonite. Their assignment to *Uptonia* rather than *Crucilobicerias* is favored by the forward curvature of their ribs on the highest parts of the flanks.

Figured specimens.—USNM 248017.

Occurrences.—Unnamed shale in northern Alaska at USGS Mesozoic locs. 29775, 29281, and 29282.

Uptonia? sp. A

Plate 8, figure 13

One fragmentary internal mold of a fairly large whorl bears ribbing similar to that on large specimens of *Crucilobicerias* and *Uptonia*. Assignment to *Uptonia* instead of *Crucilobicerias* is favored by the forward cur-

vature of the ribs at the ventral margin and by the fact that the ribs are not distinctly tuberculate.

The specimen constitutes the only evidence for the presence of Lower Jurassic beds in the Yakutat district.

Figured specimen.—USNM 248018.

Occurrence.—Float from unnamed beds in the Yakutat D-4 quadrangle, southeastern Alaska, at USGS Mesozoic loc. 29773.

***Uptonia?* sp. B**

Plate 9, figures 5-7

Two small specimens have a compressed, fairly evolute shell and fairly sharp, simple ribs that incline slightly adorally on the flanks, become stronger ventrally, form pronounced chevrons on the venter, and do not bear ventral tubercles.

These features suggest that the specimens are immature forms of *Uptonia*. Such an assignment is suggested by their association with a fairly large, typical specimen of *Apoderoceras*, which genus in Europe is characteristic of the lower part of the *Uptonia jamesoni* zone of earliest Pliensbachian Age (Dean and others, 1961, p. 463).

Figured specimen.—248019.

Occurrence.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic loc. 6697.

Genus TROPIDOCERAS Hyatt, 1867

***Tropidoceras actaeon* (d'Orbigny)**

Plate 8, figures 1-9

(For synonymy see Frebold, 1970, p. 440)

This species is represented in Alaska by four molds that are identical in appearance with the type specimens of *T. actaeon* (d'Orbigny) (1844, p. 232, pl. 61, figs. 1, 2) from France and with specimens of that species from the Queen Charlotte Islands illustrated by Frebold (1970, pl. 2, figs. 13a,b, 14a,b, 15a,b). The Alaskan specimens of the species have fairly evolute coiling. Their ribs are simple, moderately spaced, radial or slightly sigmoidal on the flanks, and they curve forward sharply on the ventral margins, where they fade out abruptly. On internal molds the venter ranges from narrowly rounded to fairly sharp and in places on some specimens bears a low blunt keel.

In Europe, this species is indicative of the lower to middle parts of the *Tragophylloceras ibex* zone of early Pliensbachian Age, according to Frebold (1970, p. 444, 445).

Hypotypes.—USNM 248020-248023.

Occurrence.—Near top of upper member of the McCarthy Formation in Wrangell Mountains at USGS Mesozoic loc. 28534.

Family AMALTHEIDAE Hyatt, 1867
Genus AMALTHEUS Montfort, 1808

***Amaltheus margaritatus* (Montfort)**

Plate 10, figures 25, 26

(For synonymy and description see Howarth, 1958, London, Palaeontographical Soc., p. 13, 14 and Dagis, 1976, p. 7-9.)

This species was provisionally identified by Imlay in 1955 (p. 87, pl. 10, fig. 4) based on a specimen from a test well in the Point Barrow area. The specific identification was subsequently confirmed by Howarth (1958, p. xxvi).

Hypotypes.—USNM 108768, 248024, 248025.

Occurrences.—Kingak Shale in northern Alaska in the South Barrow test well No. 3 at depths of 2,019 feet (639 m), 2,111 feet (634 m), and probably at 2,107 feet (641 m); in the Simpson test well No. 1 at depth of 5,677 feet (1,730 m); and in unnamed beds in the DeLong Mountains at USGS Mesozoic loc. M2441.

***Amaltheus stokesi* (J. Sowerby)**

Plate 10, figures 27, 28

(For most of the synonymy see Howarth, 1958, London, Palaeontographical Soc., pt. 1, p. 3-6.)

Amaltheus cf. *A. nudus* (Quenstedt). Imlay, 1955, USGS Prof. Paper 274-D, p. 87, pl. 10, fig. 5.

Amaltheus stokesi (J. Sowerby). Howarth, 1958, London, Palaeontographical Soc. pt. 2, p. xxvi.

This species has strong, slightly sigmoidal, radial-trending ribs that curve forward rather strongly high on the flanks, where they bifurcate or trifurcate and then merge with chevrons on the venter. It is distinguished from *Amaltheus margaritatus* (Montfort), according to Howarth (1958, p. 6), by having ventral chevrons that are larger, fewer, less forwardly inclined, and distinctly connected by twos or threes with the flank ribs instead of being connected only by striae or separated by a smooth band.

The presence of *A. stokesi* in Alaska was first recognized by Howarth (1958, p. xxvi) on the basis of an illustration published by Imlay (1955, pl. 10, fig. 5). Furthermore, Howarth identified *A. cf. A. stokesi* on the basis of another illustration by Imlay (1955, pl. 10, fig. 1). Other small specimens comparable with *A. stokesi* are illustrated on plate 10, figures 23 and 24.

Hypotypes.—USNM 108769, 248026, 248027.

Occurrences.—Kingak Shale in northern Alaska in the South Barrow test well No. 3 at depths of 2,193 and 2,198 feet (668.4 and 670 m); from unnamed beds cropping out in northern Alaska at USGS Mesozoic loc. 29165; and from Glenn Shale in east-central Alaska at Mesozoic loc. 29340.

Amaltheus cf. *A. stokesi* (J. Sowerby) occurs in the Kingak Shale in northern Alaska in the South Barrow test well No. 3 at the depth of 2,186 feet (666.3 m) and in outcrops at USGS Mesozoic loc. 30074. It occurs in the

Talkeetna Formation in the Talkeetna Mountains in southern Alaska at USGS Mesozoic loc. 25941.

Subgenus PSEUDOAMALTHEUS Frebold, 1922

A. (*Pseudoamaltheus*) engelhardtii (d'Orbigny)

This taxon is represented in northern Alaska by a fragment from the South Barrow test well No. 3 at the depth of 2090 feet (637 m). It was assigned by Imlay (1955, p. 87, pl. 10, fig. 3) to *Amaltheus* sp. and by Howarth (1958, p. xxvi) to *A. (Pseudoamaltheus) engelhardtii* (d'Orbigny). Howarth's identification was based on the presence of spiral ribs on the upper part of the flanks and on the keel, and on the lack of radial ribbing. He noted that the species and subgenus in Europe ranges from the upper part of the *Amaltheus margaritatus* zone to the end of the Pliensbachian, or *Pleuroceras spinatum* zone, and is a descendant of *Amaltheus margaritatus* (Howarth, 1958, p. 21-23).

Hypotype.—USNM 108766.

Family DACTYLIOCERATIDAE Hyatt, 1867

Genus PRODACTYLIOCERAS Spath

***Prodactylioceras italicum italicum* (Fucini)**

Plate 10, figure 3

Coeloceras lorioli Bettoni, 1900, Schweizer. palaeont. Gesell. Abh., v. 27, p. 76, pl. 7, fig. 12.

Coeloceras italicum Meneghini in Fucini, 1900, Palaeontographica Italica, v. 7, p. 72, pl. 13, fig. 4.

Coeloceras italicum Meneghini in Fucini, 1905, Palaeontographica Italica, v. 11, p. 115, pl. 6, figs. 11, 12, 14.

Prodactylioceras italicum italicum (Fucini) in Fischer, 1971, Geologica et Palaeontologica, v. 5, p. 111, pl. 2, fig. 12.

Prodactylioceras (Aveyronicerias) italicum (Meneghini in Fucini). Geczy, 1976, Akad. Kiado, Budapest, p. 145, pl. 25, figs. 8-9; pl. 26, figs. 1-4.

One laterally crushed specimen from Alaska is characterized by highly evolute coiling, by very fine, dense, simple, forwardly inclined ribs that become a little stronger ventrally, and by a few weak tubercles high on the flanks of the innermost whorls. Its appearance is identical with that of the finely ribbed subspecies, *P. italicum italicum* (Fucini) from Italy as described in Fischer (1971, p. 111). It has slightly finer ribbing than a specimen from eastern Oregon described as *P. cf. P. italicum* Meneghini (Imlay, 1968, p. C28, pl. 2, fig. 14). It differs from specimens in northern British Columbia and southern Yukon, described by Frebold (1964a, p. 10, pl. 3, figs. 2; 1970, p. 442, pl. 4, figs. 1, 2), by lacking forked ribs.

The association of the Alaskan specimen of *Prodactylioceras* with *Uptonia* shows that it occurs at or near the base of the Pliensbachian. Comparable finely ribbed specimens from southern Europe as listed herein are recorded from the zone of *Tragophylloceras ibex* near the base of the Pliensbachian (Fischer, 1971, p. 118, 123).

Figured specimen.—USNM 248030.

Occurrence.—Upper member of the McCarthy Formation in upper 100 ft (30.5 m), Wrangell Mountains, at USGS Mesozoic loc. 28671.

***Prodactylioceras cf. P. italicum fucini* R. Fischer**

Plate 10, figures 4, 5

Three crushed molds are characterized by highly evolute coiling; by high, narrow, moderately spaced nearly radial ribs that become slightly stronger ventrally and bifurcate rarely, low on the flanks; and by lacking tubercles, at least on their outer whorls. Their rib spacing resembles that on *P. italicum fucini* R. Fischer (1971, p. 111, pl. 2, figs. 8, 11) and is somewhat denser than in *P. colubriforme* (Bettoni) (1900, pl. 7, fig. 10; Pinna, 1966, pl. 10, fig. 6; Fischer, 1971, pl. 2, fig. 10) from southern Europe. The resemblance of these Alaskan specimens to *P. italicum fucini* Fischer plus their association with *Uptonia* suggests an age as early as the zone of *Uptonia jamesoni* (Fischer, 1971, p. 118, 123) or as the lower part of the *Tragophylloceras ibex* zone.

Figured specimens.—USNM 248031, 248032.

Occurrences.—Upper member of McCarthy Formation on Wrangell Mountains at USGS Mesozoic locs. 28540, 28671, and 28673.

Genus DACTYLIOCERAS Hyatt, 1867

***Dactylioceras cf. D. commune* (J. Sowerby)**

Plate 11, figures 2, 3, 8

Four laterally compressed specimens from the Talkeetna Mountains are characterized by evolute coiling and by sharp, fairly prominent, moderately spaced primary ribs which are inclined slightly forward. Most of the primary ribs pass at about two thirds of the height of the flanks into pairs of slightly weaker secondary ribs that arch gently forward on the venter. Some primary ribs remain simple, and some secondary ribs arise freely on the margins of the venter. Tiny tubercles occur at the furcation points.

These specimens show some resemblance to *Dactylioceras crassiusculosum* (Simpson) (Buckman, 1912, pl. 62; Fischer, 1966, p. 29, pl. 1, figs. 11; pl. 3, fig. 11; Pinna, 1966, p. 94, pl. 5, fig. 3; Sapunov, 1963, p. 120, pl. 2, fig. 3a,b), but they differ by having more closely spaced, weaker ribs that incline forward on the flanks instead of trending nearly radially. Compared with the lectotype of *D. commune* (Sowerby) (Buckman, 1927, pl. 707; Arkell, 1956, pl. 33, figs. 4a,b), they have denser and weaker ribs on their outermost preserved whorl. They do not differ greatly in ribbing, however, from *D. crassibundum* (Simpson), which Howarth (1962a, p. 115, pl. 16, fig. 7a,b) considered to be a synonym of *D. commune* (Sowerby).

These specimens from the Talkeetna Mountains are nearly identical with most of the specimens from arctic Canada that were assigned to *D. commune* (Sowerby) by Frebold (1958, p. 2, pl. 1, figs. 1-7; 1960, p. 18, pl. 5, figs. 4, 6) but have somewhat sparser ribbing than one specimen so assigned (Frebold, 1960, pl. 5, fig. 5).

Figured specimen.—USNM 248047.

Occurrence.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 24787 and 29198.

Subgenus ORTHODACTYLITES Buckman, 1926

***Dactylioceras* (Orthodactylites) *kanense* McLearn**

Plate 11, figures 4, 5, 9

Dactylioceras kanense McLearn, 1930, Royal Soc. Canada Trans., 3d ser., v. 24, sec. 4, p. 4, pl. 1, fig. 2

Dactylioceras kanense McLearn, 1932, Royal Soc. Canada Trans., 3d ser., v. 26, sec. 4, p. 59-62, pl. 3, fig. 5, pl. 4, figs. 1-7, 9 pl. 5, figs. 6-9.

Dactylioceras kanense McLearn. Frebold, 1964b, Canada Geol. Survey Paper 63-4, pl. 7, figs. 1-4.

?*Dactylioceras* cf. *D. kanense* McLearn. Imlay, 1955, U.S. Geol. Survey Prof. Paper 274-D, p. D88, pl. 10, fig. 14.

This species is represented by four molds that are nearly identical in appearance with the holotype. The species is characterized by an evolute, compressed shell, by a narrowly rounded venter, and by fine threadlike ribs that incline gently forward on the flanks and arch forward on the venter. On the inner whorls, furcation of the ribs is fairly common. On the largest whorl, most of the primary ribs are simple and many ribs arise freely on the upper parts of the flanks. The species shows considerable resemblance to *D. attenuatus* (Simpson) (Buckman, 1926, pl. 655) but has fewer forked ribs on its outer whorl.

Hypotype.—USNM 248048.

Occurrence.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic loc. 29198. Possibly represented in northern Alaska in the South Barrow test well No. 3 at the depth of 2,016 feet (614.5 m) (Imlay, 1955, p. 82, 88).

***Dactylioceras* (Orthodactylites) cf. *D. directum* (Buckman)**

Plate 11, figure 6

cf. *Orthodactylites directum* Buckman, 1926, Type ammonites, pl. 654.
cf. *Orthodactylites mitis* Buckman, 1927, Type ammonites, pl. 738.

One specimen, whose outermost preserved whorl is crushed laterally, resembles the holotype specimen of *D. (O.) directum* Buckman in coiling and ribbing. It also resembles *D. (O.) mitis* Buckman, which Howarth (1973, p. 254, 255) considered to be a synonym of *D. (O.) directum*. As in those species, its ribs are fairly strong, nearly straight, rectiradiate, cross the venter nearly transversely and mostly bifurcate high on the flanks. Some ribs, however, remain simple.

Figured specimen.—USNM 248049.

Occurrence.—Kingak Shale in northern Alaska at USGS Mesozoic loc. 29163.

Genus CATACOELOCERAS Buckman, 1923

***Catacoeloceras*? sp. juv.**

Plate 12, figure 6

Coeloceras aff. *C. mucronatum* (d'Orbigny). Imlay, 1955, U.S. Geol.

Survey Prof. Paper 274-D, p. D88, pl. 12, figs. 12-14.

Catacoeloceras sp. "aff. *mucronatum* (d'Orbigny)". Howarth, 1958, London, Palaeontographical Soc. pt. 2, p. xxvi.

The assignment of this species to *Catacoeloceras* by Howarth was probably based on the generic definition published by Arkell and others (1957, p. L254). Later studies by Howarth (1962b, p. 407-410) showed, however, that most of the taxa listed in the treatise as synonyms of *Catacoeloceras* actually had different characteristics than the type species of *Catacoeloceras*, occurred in older beds just above the upper range of *Amaltheus*, and could appropriately be referred to *Nodicoeloceras*. He noted that *Nodicoeloceras* had small tubercles and looped ribs only on its innermost whorls. In contrast *Catacoeloceras* developed small ventrolateral tubercles on its inner and on some intermediate whorls but did not develop looped ribs.

The stratigraphic position of the Alaskan species above *Amaltheus* and below *Dactylioceras* (Imlay, 1955, p. 82; 1968, p. C21) favors an assignment to *Nodicoeloceras* rather than to *Catacoeloceras*. Nonetheless the presence of fairly prominent tubercles and the lack of looped ribs are indicative of *Catacoeloceras* such as *C. confectum* Buckman (1923, pl. 413) and *C. crassum* (Young and Bird) (Buckman, 1918, pl. 119).

Figured specimen.—USNM 108758a-c.

Occurrence.—Kingak Shale in South Barrow test well No. 3 at depth of 2,063 ft (629 m).

Family HILDOCERATIDAE Hyatt, 1867

Subfamily ARIETICERATINAE Howarth, 1955

Genus ARIETICERAS Seguenza, 1885

***Arietoceras* cf. *A. domarense* (Meneghini)**

Plate 10, figures 1, 2, 6-15, 22

This species is represented by 38 molds that match very well with specimens from eastern Oregon described previously under the same name (Imlay, 1968, p. C33, C34, pl. 4, figs. 9-12). Most of the molds also match certain specimens from northern British Columbia and southern Yukon that were assigned to *Arietoceras* cf. *A. algovianum* (Oppel) by Frebold (1964a, p. 13, pl. 3, fig. 3a,b; pl. 5, figs. 2, 3). The species differs from the species herein called *A. cf. A. algovianum* (Oppel) by having a higher, more compressed whorl section and finer, more closely spaced ribs.

Figured specimens.—USNM 248033-248037.

Occurrences.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic locs. 24107, 27586 and 29450. Lubbe Creek Formation in the Wrangell Mountains at Mesozoic loc. 28531. Upper member of the McCarthy Formation in the Wrangell Mountains at USGS Mesozoic loc. 28688.

***Arieticer* cf. *A. algovianum* (Oppel)**

Plate 10, figures 16–20

This species is represented by three fragments that resemble certain specimens from eastern Oregon (Imlay, 1968, p. C34, pl. 4, figs. 1–8) and from northwestern British Columbia (Frebold, 1964a, pl. 3, figs. 4a, b, 5a, b; pl. 4, fig. 2) that were compared, or identified with *A. algovianum* (Oppel). These specimens are characterized by highly evolute coiling, by a low keel that is bordered by shallow furrows, and by high, narrow, widely spaced ribs that trend radially or slightly adapically on the flanks.

Figured specimens.—USNM 248038.

Occurrence.—Upper member of McCarthy Formation, in the Wrangell Mountains at USGS Mesozoic loc. 28688.

***Arieticer* sp.**

Plate 10, figure 21

On one small specimen the innermost whorls are smooth up to a diameter of about 6 mm. The next outer whorl, whose ventral part is not preserved, bears fairly low, moderately spaced ribs that curve slightly backward on the flanks. The outermost whorl bears fairly strong, widely spaced ribs that curve backward strongly on the lower and middle parts of the flanks, curve forward strongly on their upper parts, and then fade out on the ventral margin. The venter bears a low keel that is bordered by very shallow furrows.

This specimen differs from those in Oregon and British Columbia that have been compared or identified with *A. algovianum* (Oppel) (Imlay, 1968, p. C34, pl. 4, figs. 1–8; Frebold, 1964a, p. 13, pl. 3, figs. 4, 5; pl. 4, fig. 2) by having sparser ribs that curve backward much more strongly on the flanks. Its rib plan shows more resemblance to that of *Arieticer* *retrorsicosta* (Oppel) as in Quenstedt (1885, pl. 42, fig. 44) and *A. cf. A. retrorsicosta* (Oppel) in Monestier (1934, pl. 10, figs. 42, 43).

Figured specimen.—USNM 248039.

Occurrence.—Lubbe Creek Formation in the Wrangell Mountains at USGS Mesozoic loc. 28531.

Genus *LEPTALEOCERAS* Buckman, 1918

***Leptaleoceras* cf. *L. pseudoradians* (Reynes)**

Plate 11, figures 12, 13

Ammonites pseudoradians Reynes, 1868, Essai de géologie et de paléontologie aveyronnaises: Paris (Baillière), p. 91, pl. 1, figs. 4a–c.

Arieticer *pseudoradians* (Reynes). Monestier, 1934, Soc. Géol.

France Mém., new ser., v. 10, pt. 3 (Mém. 23), p. 63, pl. 8 figs. 66 and 68.

Leptaleoceras pseudoradians (Reynes). Howarth, 1957, Geol. Soc.

London Quart. Jour., v. 113, p. 198, pl. 17, figs. 1, 2.

Leptaleoceras pseudoradians (Reynes). Frebold, 1964a, Canada Geol. Survey Bull. 116, p. 15, pl. 4, figs. 5–7, pl. 5, figs. 4, 5.

This species is represented by one fairly well preserved specimen that retains some shell material and is mostly septate. Its characteristics are identical with those of specimens from the southern Yukon and British Columbia as illustrated and described by Frebold.

Hypotype.—USNM 248040.

Occurrence.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic loc. 29450.

Genus *FONTANELLICERAS* Fucini, 1931

***Fontanelliceras* cf. *F. fontanellense* (Gemmellaro)**

Plate 11, figures 17–23

(For synonymy see Imlay, 1968, U.S. Geol. Survey Prof. Paper 593-C, p. C36.)

cf. *Fontanelliceras fontanellense* (Gemmellaro). Guex, 1973, Eclogae geol. Helvetiae, v. 66, p. 507, pl. 6, fig. 2.

The genus *Fontanelliceras* is represented by five specimens. The largest specimen (pl. 11, figs. 17, 18, 23) has highly evolute coiling; a tricarinate bisulcate venter; and high, straight, rursiradial ribs that become stronger ventrally and bend forward slightly before terminating on the ventral margin. The smaller specimens have been crushed laterally but resemble the larger in most respects. They differ by having ribs that are sharper, more widely spaced, and nearly radially arranged. The venter, which is well exposed on the largest specimen, bears a prominent keel bordered by furrows that in turn are bordered by weak ridges.

At a diameter of 59 mm, the largest specimen has a whorl height of 14 mm, a whorl thickness of 11 mm, and an umbilical width of 35 mm. Its suture line is not exposed.

Figured specimens.—USNM 248041–248043.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 25941, 27586, and 29450.

Subfamily *HARPOCERATINAE* Neumayr, 1875

Genus *HARPOCERAS* Waagen, 1869

***Harpoceras* cf. *H. exaratum* (Young and Bird)**

Plate 11, figure 11

cf. *Harpoceras exaratum* (Young and Bird). Wright, 1882, London, Palaeontographical Soc. pl. 62, figs. 1–3.

Harpoceras cf. *H. exaratum* (Young and Bird). Imlay, 1955, U.S. Geol. Survey Prof. Paper 274-D, p. 88, pl. 11, figs. 12, 13, 15.

Harpoceras cf. *H. exaratum* (Young and Bird). Frebold and Little, 1962, Canada Geol. Survey Bull. 81, p. 17, 18, pl. 2, figs. 1–9, pl. 3, fig. 5.

Harpoceras cf. *H. exaratum* (Young and Bird). Frebold, 1964 a, Canada Geol. Survey Bull. 116, p. 16, pl. 6, figs. 2–4 (not 1 and 5).

Harpoceras cf. *H. exaratum* (Young and Bird). Frebold, 1957, Canada Geol. Survey Mem. 287, p. 47, pl. 17, fig. 1; pl. 18, figs. 2, 3.

This species is represented by five fragmentary specimens from northern Alaska and one from the Talkeetna Mountains in southern Alaska. Their features match very well with those on specimens from western and Arctic Canada as described and illustrated by Frebold in the papers listed above.

Figured specimen.—USNM 248044.

Occurrences.—Kingak Shale in northern Alaska at USGS Mesozoic localities 22081, 29159, 29160, 29161, and 29776. Talkeetna Formation in Talkeetna Mountains at Mesozoic loc. 29198.

Genus *PROTOGRAMMOCERAS* Spath, 1913

Protogrammoceras cf. *P. paltum* (Buckman)

Plate 12, figures 11, 12

cf. *Paltarpites paltus* Buckman, 1922, Type ammonites, v. 4, pl. 362a and 1923, pl. 362b.

Harpoceras cf. *H. exaratum* (Young and Bird). Frebold, 1964a, Canada Geol. Survey Bull. 116, pl. 6, figs. 1, 5 (not 2-4).

cf. *Paltarpites paltus* Buckman. Frebold, 1970, Canadian Jour. Earth Sci., v. 7, no. 2, p. 443, pl. 4, figs. 5-7.

Three fragmentary specimens from Alaska bear ribbing similar to that on *Protogrammoceras paltum* (Buckman). They differ from *P. argutum* (Buckman) by having fewer ribs that are broader and flat topped. Nonetheless, as discussed by Frebold (1970, p. 443), the ribs during growth on some specimens of *P. paltum* vary considerably in strength from fine to coarse to fine, or even to striae. Consequently fragments of whorls bearing only striae, or only very fine ribs, cannot reliably be assigned to either species.

The assignment of both species to *Protogrammoceras* Spath rather than to *Paltarpites* Buckman, is based on examination of the type specimens by Howarth (1973, p. 265).

Figured specimen.—USNM 248045.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 24108, 25941, and 29449.

Protogrammoceras cf. *P. argutum* (Buckman)

Plate 11, figure 14

cf. *Argutarpites argutus* Buckman, 1923, Type ammonites, v. 4, pl. 363.

Paltarpites cf. *P. argutus* Buckman. Imlay, 1968, U.S. Geol. Survey Prof. Paper 593-C, p. C37, pl. 5, figs. 20-22, 24, 25.

cf. *Paltarpites argutus* (Buckman). Frebold, 1970, Canadian Jour. Earth Sci., v. 7, no. 2, p. 444, pl. 4, figs. 8-10.

This species is represented by two fragments, that are characterized by very fine falcoid ribs and striae and by a rounded umbilical edge. It could be interpreted as a finely ribbed variant of *P. paltum* (Buckman), which species is associated with *P. argutum* (Buckman) in western Canada (Frebold, 1970, p. 444).

Figured specimen.—USNM 248046.

Occurrences.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic locs. 24108 and 25941.

Genus *ELEGANTICERAS* Buckman, 1913

Eleganticeras sp. juv.

Plate 11, figures 10, 15, 16

This species is represented by 12 small internal and external molds. They have a moderately compressed planulate form; a low vertical umbilical wall that rounds fairly abruptly into somewhat flattened flanks; a rather sharp keel on the venter; and an incomplete body chamber that occupies at least half a whorl. The innermost whorls are smooth. Weak, widely spaced ribs appear at a diameter of 7 to 9 mm and gradually become stronger during growth. The largest preserved septate whorl and the adapical end of the nonseptate whorl bear low falcoid ribs that are faint on the lower third of the flanks, become stronger ventrally, and extend almost to the keel. Such ribs are replaced rather abruptly on the adoral half of the largest preserved whorls by much weaker falcoid ribs and then by striae that are apparent only where some of the shell is preserved. The outermost whorl embraces about half of the preceding whorl.

In shape, coiling, and ribbing, this species greatly resembles the holotype of *Elegantuliceras elegantulum* (Young and Bird) (Buckman, 1914, pl. 93). It differs by being only half as large and by its umbilical wall being vertical instead of concave.

Figured specimen.—USNM 248050.

Occurrence.—Kingak Shale in northern Alaska at USGS Mesozoic loc. 29776.

Genus *PSEUDOLIOCERAS* Buckman, 1913

Pseudolioceras cf. *P. compactile* (Simpson)

(See Imlay, 1955, U.S. Geol. Survey Prof. Paper 274-D, p. 89, pl. 12, figs. 17, 18, 21.)

Occurrence.—Kingak Shale at USGS Mesozoic loc. 23772.

Pseudolioceras cf. *P. lythense* (Young and Bird)

(See Imlay, 1955, U.S. Geol. Survey Prof. Paper 274-D, p. D89, pl. 12, fig. 20.)

Occurrence.—Kingak Shale at USGS Mesozoic loc. 23772.

Pseudolioceras sp.

Plate 11, figure 1; Plate 12, figure 13

Two laterally compressed specimens from southern Alaska resemble *P. macIntocki* (Haughton) from arctic Canada and arctic Alaska (Frebold, 1964b, pl. 10, figs. 4-8, 12; Imlay, 1976, pl. 1, figs. 1-5, 7.) They appear to have a smaller umbilicus and sparser, weaker ribbing, but their preservation is not sufficient for a positive specific determination.

Figured specimens.—USNM 248051, 248052.

Occurrences.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic loc. 25317; Kialagvik Formation at Puale Bay on the Alaska Peninsula at USGS Mesozoic loc. 19804.

Subfamily GRAMMOCERATINAE Buckman, 1904
Genus GRAMMOCERAS Hyatt, 1867

***Grammoceras* sp.**

Plate 11, figure 7

One laterally crushed ammonite is characterized by evolute coiling by simple, strong, gently sigmoid ribs, and by a low keel as in *Grammoceras*. It has slightly coarser ribbing than does the ammonite from the southern Yukon described by Frebold (1964a, p. 17, pl. 7, figs. 1, 2) as *G. aff. G. fallaciosum* (Bayle).

Figured specimen.—USNM 248053.

Occurrence.—Talkeetna Formation, upper part, in the Talkeetna Mountains at USGS Mesozoic loc. 25939.

Family HAMMATOCERATIDAE Buckman, 1887
Genus PHYMATOCERAS Hyatt, 1867

***Phymatoceras?* sp.**

Plate 12, figures 14, 16–19

The genus is possibly represented by three specimens. The smallest specimen (pl. 12, fig. 18) bears twinned ribs and a carinate venter similar to that on *Phymatoceras binodata* (Buckman) (1898, Supplement, p. XVI, pl. 1, figs. 11, 12). In each twin the adapical rib is the stronger and is marked near the umbilical edge by a swelling. All ribs curve forward on the venter but are separated from the keel by a smooth area.

The next larger specimen is septate and bears alternating long and short ribs. The long ribs begin on the umbilical wall and are slightly swollen on the umbilical margin. The short ribs begin at or a little above the middle of the flanks. All ribs incline forward on the flanks and venter, become stronger ventrally, become swollen on the venter, and then terminate abruptly before reaching the keel.

The largest specimen is a nonseptate fragment that shows the venter and part of the flank of an adult whorl. It bears broad coarse ribs that are wider than the interspaces, that curve forward slightly on the flanks and more strongly on the venter, and that then terminate abruptly before reaching a rather low keel. The ventral aspect of this specimen is similar to that of *P. tumefacta* (Buckman) (1898, Supplement, p. XIX, pl. 1, figs. 7–10).

Figured specimens.—USNM 248054–248056.

Occurrences.—Talkeetna Formation, upper part, in the Talkeetna Mountains at USGS Mesozoic locs. 25317 and 25342.

Genus BRODIEIA Buckman, 1898

***Brodieia* cf. *B. tenuicostata* var. *nodosa* (Jaworski)**

Plate 12, figure 8

cf. *Hildoceras* (*Brodiceras*) *tenuicostatum* var. *nodosa* Jaworski, 1926, Actas Acad. Nac. Cienc. Cordoba, v. 9, p. 245, pl. 4, fig. 8a,b.

One laterally crushed ammonite consists mostly of an incomplete body chamber that occupies at least three-fourths of a whorl and embraces about two-thirds of the preceding whorl. The venter bears a single low keel. The ribs on the outer whorl are moderately strong, widely spaced, gently falcate, and terminate ventrally near the keel. The primary ribs begin low on the umbilical wall, are fairly strong near the umbilicus, weaken a little near the middle of the flanks, and are strongest on the margin of the venter. Some rib furcation occurs at various heights ranging from the umbilical margin to a little above the middle of the flanks. Other ribs arise freely near or above the middle of the flanks. All ribs are essentially of the same strength on the venter. The ribbing on the penultimate whorl as revealed in the umbilicus is also fairly strong.

This ammonite from Alaska bear ribbing almost identical with that on *Brodieia tenuicostata* var. *nodosa* (Jaworski) (1926, pl. 4, fig. 8a) from Argentina, but it differs by being a little more involute. In that respect it resembles the typical forms of *B. tenuicostata* (Jaworski) (1926, pl. 4, figs. 1–4, 6, 7) but differs by having slightly coarser and sparser ribbing. Overall the resemblances are remarkable.

Figured specimen.—USNM 248057.

Occurrence.—Talkeetna Formation in Talkeetna Mountains at USGS Mesozoic loc. 25940.

Genus HAUGIA Buckman, 1888

***Haugia* cf. *H. grandis* Buckman**

Plate 12, figures 4, 10, 15

cf. *Haugia grandis* Buckman, 1898, Supplement, p. XXVI, pl. s, fig. 11; 1888, pl. 23, figs. 14, 15; 1889, pl. 24, pl. 25, fig. 1

Ten specimens belong to a fairly involute species that has sharp, moderately spaced, flexuous ribs. The largest specimen is similar in size, involution, and ornamentation to a small specimen of *H. grandis* Buckman (1888, pl. 23, figs. 14, 15), but its tubercles are smaller and its ribs arch forward more strongly on its ventral margin. *H. dumortieri* Buckman (1888, pl. 23, figs. 16, 17) has fewer tubercles and ribs at a comparable size.

Figured specimen.—USNM 248058.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 24111 and 24114; Kialagvik Formation at Puale Bay on the Alaska Peninsula at Mesozoic loc. 19804.

***Haugia* cf. *H. variabilis* (d'Orbigny)**

Plate 12, figures 1, 2, 5

cf. *Ammonites variabilis* d'Orbigny, 1845, *Paléontologie française, Terrains Jurassiques*, v. 1, Cephalopods, p. 350, p. 113, figs. 1 and 2 only.

cf. *Harpoceras variable* (d'Orbigny). Wright, 1882, London, *Palaeontographical Soc.*, p. 455, pl. 67, figs. 5, 6 only.

cf. *Haugia* aff. *H. variabilis* (d'Orbigny). Buckman, 1890, London, *Palaeontographical Soc.*, p. 146, pl. 25, fig. 2.

This species is represented by nine crushed molds. It has fairly evolute coiling. The ornamentation on its inner whorls consists of umbilical tubercles, of sharp, nearly radially trending ribs that arise mostly by two's and three's from the tubercles, and of a few ribs that arise freely between the tubercles. During subsequent growth, both tubercles and ribs gradually become stronger and more widely spaced. The ribs become gently flexuous on the flanks, arch forward slightly on the venter, and arise mainly in pairs from prominent conical tubercles. The characteristics of the adult body chamber are unknown.

The Alaskan species may be within the range of variation of *H. variabilis* (d'Orbigny), but it appears to be more evolute and to develop sparser ribbing. In these respects it shows more resemblance to *H. aff. H. japonica* (Neumayr) as illustrated by Matsumoto and Ono (1947, pl. 2, figs. 5). It has much coarser ornamentation than the holotype of *H. japonica* (Neumayr) (Kobayashi, 1935, pl. 12, figs. 3, 4).

Figured specimens.—USNM 248059.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic locs. 25318 and 27508.

***Haugia* cf. *H. compressa* Buckman**

Plate 12, figures 3, 7, 9

cf. *Haugia compressa* Buckman, 1898, London, *Palaeontographical Soc.*, Supplement, p. XXVII, pl. 2, figs. 8–10.

Nine molds represent a fairly involute and densely ribbed species. The largest specimen at a diameter of 66 mm bears about 21 umbilical tubercles and about 65 gently flexuous ribs that arise in two's and three's from the tubercles. Many ribs arise freely along the zone of tuberculation.

This species closely resembles *H. compressa* Buckman in involution and in density of ribbing but at a comparable size has slightly finer ribbing and smaller tubercles than the holotype.

Figured specimens.—USNM 248060, 248061.

Occurrences.—Talkeetna Formation in the Talkeetna Mountains at USGS Mesozoic loc. 24114; basal part of Kialagvik Formation at Puale Bay on Alaska Peninsula at Mesozoic loc. 19804.

REFERENCES CITED

- Arkell, W. J., 1956, *Jurassic geology of the world*: London, Oliver and Boyd, 806 p., 46 pls., 102 figs.
- Arkell, W. J., and others, 1957, *Treatise on invertebrate paleontology*, R. C. Moore, ed.; pt. L, Mollusca 4—Cephalopoda, Ammonoidea: New York, Geol. Soc. America, and Lawrence, Univ. Kansas Press, 490 p.
- Bettoni, Andrea, 1900, *Fossili Domeriani della Provincia di Brescia*: Schweizer. palaeont. Gesell. Abh., v. 27, p. 1–88, pls. 1–9.
- Bremer, Heinrich, 1965, *Zur Ammoniten Fauna und Stratigraphie des unteren Lias (Sinemurium bis Carixium) in der Umgebung von Ankara (Turkei)*: Neues Jahrbuch Geologie Paläontologie Abh., v. 122, no. 2, p. 127–221, pls. 12–16, 12 figs.
- Brosigé, W. P., and Reiser, H. N., 1964, *Geologic map and section of the Chandalar quadrangle, Alaska*: U.S. Geol. Survey Misc. Geol. Inv. Map I-375.
- 1969, *Preliminary geologic map of the Coleen quadrangle, Alaska*: U.S. Geol. Survey open-file report, scale 1:250,000.
- Buckman, S. S., 1887–1907, *Monograph of the ammonites of the Inferior Oolite Series*: London, Palaeontographical Soc., 456 p., 103 pls. (1887–1894); Supplement, 262 p., 24 pls. (1893–1907).
- 1909–1930, *Yorkshire type ammonites*, v. 1 and 2 (1909–1919); *Type ammonites*, v. 3–7 (1920–1930), ed. A. M. Davies (1930).
- Crickmay, C. H., 1928, *The stratigraphy of Parson Bay, British Columbia*: California Univ. Pub. Dept. Geol. Sci., Bull., v. 18, no. 2, p. 51–70, 4 pls., 1 map.
- 1933, *Some of Alpheus Hyatt's unfigured types from the Jurassic of California*: U.S. Geol. Survey Prof. Paper 175-B, p. 51–58, pl. 14–18.
- Dagis, A. A., 1976, [Late Pliensbachian ammonites (Amaltheidae) of north Siberia]: Novosibirsk, Publishing House Nauk USSR, Siberian Branch, p. 5–60, pls. 1–17, 8 figs. (In Russian)
- Dean, W. T., Donovan, D. T., and Howarth, M. K., 1961, *The Liassic ammonite zones and subzones of the north-west European province*: British Mus. (Nat. History) Bull., v. 4, no. 10, p. 437–505, pls. 63–75.
- Detterman, R. L., Reiser, H. N., Brosigé, W. P., and Dutro, J. T., Jr. 1975, *Post-Carboniferous stratigraphy, northeastern Alaska*: U.S. Geol. Survey Prof. Paper. 886, 46 p., 14 figs.
- Donovan, D. T., 1952, *The ammonites of the Blue Lias of the Bristol district*: Annals and Mag. Nat. History, ser. 12, v. 5, p. 629–655, 717–752, pls. 22–23, 27–29.
- 1955, *Revision des espèces décrites dans la Monographie des Ammonites (Lias inférieur) de P. Reynes*: Soc. géol. France, Mém., new ser., v. 73, p. 1–45, 2 pls.
- Donovan, D. T., and Forsey, G. F., 1973, *Systematics of lower Liassic Ammonitina*: Kansas Univ. Paleont. Contr., Paper 64, 18 p., 4 pls.
- Erben, H. N., 1956, *El Jurasico Inferior de Mexico y sus amonitas*: Internat. Geol. Cong., 20th, Mexico, D. F., 1956, 393 p., 41 pls., 5 maps.
- Fischer, Rudolf, 1966, *Die Dactylioceratidae (Ammonoidea) der Kammerker (Nordtirol) und die zonengliederung des Alpenen Toarcian*: Bayerische Akad. Wiss. Math.—Naturw. Kl., Abt. Abh., new ser., No. 126, p. 1–83, pls. 1–6, 4 figs.
- 1971, *Die Coeloceratinen (Ammonoidea) des Monte Cetona (Prov. Siena)—Paläontologie, Geologie und Stratigraphie eines Profils im Mediterranen Carizium und Lotharingium*: Geologica et Palaeontologica, v. 5, p. 93–129, 2 pls., 12 figs.
- Frebold, Hans, 1951, *Ammonite fauna and stratigraphy of the lower Lias in Tyauthton Lake map-area, British Columbia*: Canada Geol. Survey Bull. 18, p. 1–14, pls. 1–14, 16–18.
- 1957, *The Jurassic Fernie group in the Canadian Rocky Mountains and foothills*: Canada Geol. Survey Mem. 287, 197 p., 44 pls., 5 figs.

- 1958, Fauna, age and correlation of the Jurassic rocks of Prince Patrick Island: Canada Geol. Survey Bull. 49, 32 p., pls. 1-18, 2 figs.
- 1960, The Jurassic faunas of the Canadian Arctic—Lower Jurassic and lowermost Middle Jurassic ammonites: Canada Geol. Survey Bull. 59, 33 p., 15 pls., 8 figs.
- 1964a, Lower Jurassic and Bajocian ammonoid faunas of north-western British Columbia and southern Yukon: Canada Geol. Survey Bull. 116, 31 p., 8 pls., 1 fig.
- 1964b, Illustrations of Canadian fossils—Jurassic of western and arctic Canada: Canada Geol. Survey Paper 63-4, 106 p., 51 pls.
- 1967a, Position of the Lower Jurassic genus *Fanninoceras* McLearn and the age of the Maude Formation on Queen Charlotte Islands: Canadian Jour. Earth Sci., v. 4, no. 6, p. 1145-1149, 1 pl.
- 1967b, Hettangian ammonite faunas of the Taseko Lakes area, British Columbia: Canada Geol. Survey Bull. 158, 35, p., 9 pls. 6 figs.
- 1970, Pliensbachian ammonoids from British Columbia and southern Yukon: Canadian Jour. Earth Sci., v. 7, no. 2, p. 435-456, pls. 1-4.
- Frebold, Hans, and Little, H. W., 1962, Paleontology, stratigraphy, and structure of the Jurassic rocks in Salmo map-area, British Columbia: Canada Geol. Survey Bull. 81, 31 p., 5 pls., 1 text-fig.
- Frebold, Hans, and Poulton, T. P., 1977, Hettangian (Lower Jurassic) rocks and faunas, northern Yukon Territory: Canadian Jour. Earth Sci., v. 14, no. 1, p. 89-101, 2 pls., 3 figs.
- Fucini, Alberto, 1900, Ammoniti del lias medio dell Appennino centrale: Palaeontographica Italica, v. 6, p. 17-78, pls. 7-13.
- 1905 Cefalopoda liassici del Monte di Cetona: Palaeontographica Italica, v. 11, p. 93-146, pls. 3-11.
- Geczy, Barnabas, 1976, Les ammonitines du Carixien de la montagne du Bakony: Budapest, Akademiai Kiado, 223 p., 39 pls., 138 figs.
- Getty, T. A., 1973, A revision of the generic classification of the family Echioceratidae (Cephalopoda, ammonioidea) (Lower Jurassic): Kansas Univ. Paleont. Contr. Paper 63, p. 1-32, 5 pls., 4 figs.
- Grantz, Arthur, Thomas, Herman, Stern, T. W., and Sheffey, N. B., 1963, Potassium-argon and lead-alpha ages for stratigraphically bracketed plutonic rocks in the Talkeetna Mountains, Alaska: U.S. Geol. Survey Prof. Paper 475-B, p. B56-B59.
- Guerin-Franiatte, S., 1966, Ammonites du Lias Inferieur de France—Psiloceratidae, Arietitidae: Paris, Centre natl. recherche sci., 2 v., 231 pls., 190 figs., 10 maps.
- Guex, Jean, 1973, Aperçu biostratigraphique sur le Toarcien inferieur du Moyen-Atlas marocain et discussion sur la zonation de ce sous-étage dans les séries méditerranéennes: Eclogae geol. Helvetiae, v. 66, no. 3, p. 493-523, 15 pls., 4 figs.
- Guex, Jean and Taylor, David, 1976, La limite Hettangian-Sinemurian des Prealpes romandes au Nevada: Eclogae geol. Helvetiae, v. 69, no. 2, p. 521-526.
- Hallam, Anthony, 1965, Observations on marine Lower Jurassic stratigraphy of North America, with special reference to United States: Am. Assoc. Petroleum Geologists Bull., v. 49, p. 1481-1501, 5 figs.
- Howarth, M. K., 1957, The Middle Lias of the Dorset Coast: Geol. Soc. London Quart. Jour., v. 113, p. 185-203, pl. 17.
- 1958, A monograph of the ammonites of the Liassic family Amaltheidae in Britain: London Palaeontographical Soc., pt. 1, p. i-xiv, 1-26, pls. 1-4; pt. 2, p. xv-xxxvii, 27-53, pls. 5-10.
- 1962a, The Yorkshire type ammonites and nautiloids of Young and Bird, Phillips, and Martin Simpson: Palaeontology, v. 5, pt. 1, p. 93-136, pls. 13-19.
- 1962b, The Jet Rock Series and the Alum Shale Series of the Yorkshire Coast: Yorkshire Geol. Soc. Proc., v. 33, pt. 4, p. 381-422, pls. 24-28.
- 1973, The stratigraphy and ammonite fauna of the upper Liassic Grey Shales of the Yorkshire Coast: British Mus. (Natural History) Bull., v. 24, no. 4, p. 237-277, pls. 1-9, 6 figs.
- Imlay, R. W., 1955, Characteristic Jurassic mollusks from northern Alaska: U.S. Geol. Survey Prof. Paper 274-D, p. 69-96, pls. 8-13, 1 fig., 4 tables.
- 1967, The Mesozoic pelecypods *Otapiria* Marwick and *Lupherella* Imlay, new genus, in the United States: U.S. Geol. Survey Prof. Paper 573-B, p. B1-B11, pls. 1, 2, 2 figs.
- 1968, Lower Jurassic (Pliensbachian and Toarcian) ammonites from eastern Oregon and California: U.S. Geol. Survey Prof. Paper 593-C, p. C1-C51, 9 pls., 8 figs.
- 1976, Middle Jurassic (Bajocian and Bathonian) ammonites from northern Alaska: U.S. Geol. Survey Prof. Paper 854, 22 p., 4 pls. 5 figs.
- Imlay, R. W., and Detterman, R. L., 1973, Jurassic paleobiogeography of Alaska: U.S. Geol. Survey Prof. Paper 801, 34 p., 15 figs.
- 1977, Some Lower and Middle Jurassic beds in Puale Bay-Alinchak Bay area, Alaska: Am. Assoc. Petroleum Geologists Bull., v. 61, p. 607-611, 2 figs.
- Jaworski, Erich, 1926, La fauna del Lias y Dogger de la Cordillera Argentina en la parte meridional de la Provincia de Mendoza: Acad. Nac. Cienc. Cordoba Actas, v. 9, nos. 3, 4, p. 139-319, pls. 1-4.
- Jones, D. L., Silberling, N. J., Csejtey, Bela, Nelson, W. T., and Blome, C. C., 1980, Age and structural significance of ophiolite and adjoining rocks in the upper Chulitna district, south-central Alaska: U.S. Geol. Survey Prof. Paper 1121-A, 21 p., 12 pls.
- Kelley, J. S., 1978, Lithofacies in Lower Jurassic volcanoclastic rocks adjacent to lower Cook Inlet, southwestern Kenai Peninsula, Alaska: Am. Assoc. Petroleum Geologists Bull., v. 62, no. 11, p. 2356.
- Kobayashi, Teiichi, 1935, Contribution to the Jurassic Torinosu series of Japan: Japanese Jour. Geology and Geography, v. 12, nos. 3-4, p. 59-91, pls. 12, 13.
- Lange, Warner, 1941, Die Ammoniten fauna der *Psiloceras*-stufe Norddeutschlands: Palaeontographica (Stuttgart), v. 93, Abt. A, 192 p., 20 pls.
- 1951, Die Schlotheimiinae aus dem Lias alpha Norddeutschlands: Palaeontographica (Stuttgart), v. 100, Abt. A, p. 1-128, 20 pls.
- MacKevett, E. M., Jr., 1969, Three newly named Jurassic formations in the McCarthy C-5 quadrangle, Alaska: U.S. Geol. Survey Bull. 1274A, p. 35-49, figs. 3-8.
- 1970, Geologic map of the McCarthy C-4 quadrangle, Alaska: U.S. Geol. Survey Geol. Quad. Map GQ-844.
- 1971, Stratigraphy and general geology of the McCarthy C-5 quadrangle, Alaska: U.S. Geol. Survey Bull. 1323, 35 p., 3 figs.
- Martin, G. C., 1926, The Mesozoic stratigraphy of Alaska: U.S. Geol. Survey Bull. 776, 493 p., 13 figs.
- Matsumoto, Tatsuro, and Ono, Akira, 1947, A biostratigraphic study of the Toyora group, with special reference to ammonites: Kyushu Univ. Fac. Sci., Sci. Repts. Geology, v. 2, no. 1, p. 20-33, pls. 1, 2.
- McLearn, F. H., 1930, Notes on some Canadian Mesozoic faunas: Roayl Soc. Canada Trans., ser. 3, v. 24, sec. 4, p. 1-8, 2 pls.
- 1932, Contributions to the stratigraphy and paleontology of Skidgate Inlet, Queen Charlotte Islands, British Columbia: Royal. Soc. Canada Trans., ser. 3, v. 26, sec. 4, p. 51-80, pls. 1-10.
- Monestier, Joseph, 1934, Ammonites du Domerien de la region sud-est de l'Aveyron et de quelques regions de las Lozere à l'exclusion des Amaltheides: Soc. Géol. France Mém., new ser., v. 10, pt. 3 (Mém. 23), 102 p., 111 pls.
- Mousterde, Rene, 1951, Ammonites du Lias Portugais: Soc. géol. Portugal Bol., v. 9, no. 3, p. 175-188, 2 pls., 5 figs.

- Mouterde, Rene, and Ruget, Ch., 1970, Le Lias moyen de Sao Pedro de Muel, 1, Cephalopodes: Portugal Com. Serv. Géol., v. 54, p. 39-74, pls. 1-8.
- Neumayr, Melchior, 1879, Zur Kenntnis der Fauna des unteren Lias in den Nordalpen: Geol. Reichsanst. Wien Abh., v. 7, pt. 5, 46 p., 7 pls.
- Orbigny, Alcide d', 1842-1851, Paléontologie française—Terrains Jurassiques, v. 1, Cephalopods: Paris, 642 p., 234 pls. (For details on dates see Geol. Mag., new ser. v. 6, p. 223-225. Davies Sherborn 1899.)
- Pinna, Giovanni, 1966, Ammoniti del Lias Superiore (Toarcian) dell'Alpe Turati (Erba, Como) Famiglia Dactylioceratidae: Soc. Ital. Sci. Nat. Mem., v. 14, pt. 2, p. 85-136, 4 pls.
- Pinna, Giovanni, and Levi-Setti, Franco, 1971, I Dactylioceratidae della Provincia Mediterranea (Cephalopoda Ammonoidea): Soc. Italiana Sci. Nat. e Mus. Civico Storia Nat. Milano Mem., v. 19, pt. 2, p. 49-136, pls. 1-12, 21 text-figs.
- Quenstedt, F. A., 1856-1858, Der Jura: Tübingen, p. 1-208, pls. 1-24 (1856); p. 209-842, pls. 25-100 (1858).
- 1882-1885, Die Ammoniten des Schwabischen Jura., v. 1, Der Schwarze Jura (Lias): Tübingen, p. 1-48, pls. 1-6 (1882); p. 49-96, pls. 7-12 (1883); p. 97-240, pls. 13-30 (1884); p. 241-440, pls. 31-54 (1885).
- Reynes, P., 1868, Essai de géologie et de paléontologie aveyronnaises: Paris (Baillière), 110 p., 7 pls.
- 1879, Monographie des ammonites Lias: Marseilles and Paris, atlas, 58 pls. [text never published].
- Sapunov, I. G., 1963, [Toarcian ammonites of the family Dactylioceratidae from western Bulgaria]: Bulgar. Akad. Nauk, Geol. Inst., Trudove Geol. Bulgar., Ser. Paleontologiya, v. 5, p. 109-136, 6 pls. (In Bulgarian; Russian and English summaries.)
- Schlegelmilch, Rudolf, 1976, Die Ammonitendes sueddeutschen Lias: Stuttgart, Gustav Fischer Verlag, 212 p., 52 pls., 21 figs.
- Schmidt-Effing, Reinhard, 1972, Die Dactylioceratidae, eine Ammoniten-familie des unteren Jura: Münstersche Forsch. Geologie und Paläontologie, No. 25 and 26, 216 p., 19 pls., 31 figs.
- Sowerby, James, and Sowerby, J. de C., 1812-46, Mineral conchology of Great Britain: London, B. Meredith, 7 v., pls. 1-383 (1812-22) by J. Sowerby; pls. 384-648 (1823-46) by J. de C. Sowerby.
- Spath, L. F., 1925, Notes on Yorkshire ammonites: Naturalist, p. 107-112, 137-141, 167-172, 201-206, 263-269, 299-306, 327-331, 359-364.
- Trueman, A. E., and Williams, D. M., 1925, Studies in the family Echioceratidae: Royal Soc. Edinburgh Trans., v. 53, p. 699-739, pls. 1-4.
- Waechner, Franz, 1882-98, Beiträge zur Kenntnis der tieferen Zonen des unteren Lias in den nordöstlichen Alpen: Beitr. Paläont. Geol. Oster.-Ungarns v. des Orients, v. 2-11, 291 p., 66 pls.
- Wright, Thomas, 1878-86, Monograph on the Lias ammonites of the British Islands: London, Palaeontographical Soc., p. 1-48, pls. 1-8 (1878); p. 49-164, pls. 9-18 (1879); p. 165-264, pls. 19-40 (1880); p. 265-328 (pls. 22A, B, 41-48 (1881); p. 329-400, pls. 49-52, 52A, 53-69 (1882); p. 401-440, pl. 70-77 (1883); p. 441-480, pls. 78-87 (1884); p. 481-503, pl. 88 (1886).

INDEX

[Italic page numbers indicate descriptions and major references]

A	
<i>actaeon</i> , <i>Tropidoceras</i>	18, 20, 35, 37; pl. 8
<i>Aegasteroceras</i>	2, 3
<i>simile</i>	3
<i>Aegoceras extracostatum</i>	30
<i>portlocki</i>	30
<i>tenerum</i>	30
Aichilik River	5
Alaska Peninsula	12, 28
Alaska Range	10, 12, 31, 33, 34
<i>algovianum</i> , <i>Arietoceras</i>	20, 39, 40; pl. 10
Alinchak Bay	11, 12, 28, 29, 30
<i>Alsatites liasicus</i> zone	12
Amaltheidae	4, 37
<i>Amaltheus</i>	17, 18, 37, 39
<i>margaritatus</i>	17, 19, 37; pl. 10
zone	38
<i>nudus</i>	37
<i>stokesi</i>	7, 17, 18, 19, 21, 37; pl. 10
sp	17, 19, 38
(<i>Pseudoamaltheus</i>)	2
<i>engelhardti</i>	17, 38
Ammonite zones:	
Hettangian	11
Pliensbachian	17
Sinemurian	12
Toarcian	18
<i>Ammonites pseudoradians</i>	40
<i>variabilis</i>	43
Anatuvuk River	17
<i>angulata</i> , <i>Schlotheimia</i>	12
<i>aplanatum</i> , <i>Paltechioceras</i>	34
<i>Apoderoceras</i>	2, 35, 37
<i>subtriangulare</i>	18, 21, 35; pl. 8
Arctic Coastal Plain	12, 18
<i>Arctoasteroceras</i>	2, 3, 16, 33
<i>jeletzkyi</i>	3, 16, 22, 31, 33; pl. 5
<i>Argutarpites argutus</i>	41
<i>argutum</i> , <i>Protogrammoceras</i>	18, 21, 41; pl. 11
<i>argutus</i> , <i>Argutarpites</i>	41
<i>argutus</i> , <i>Paltarpites</i>	41
<i>Arietoceras</i>	2, 18, 39
<i>algovianum</i>	20, 39, 40; pl. 10
<i>domarense</i>	18, 20, 21, 39; pl. 10
<i>pseudoradians</i>	40
<i>retroscicosta</i>	40
sp	20, 40; pl. 10
<i>Arietoceratinae</i>	39
<i>Arietites</i>	2
<i>bucklandi</i>	12, 13, 17, 32; pl. 4
zone	13
<i>Arietitid</i> genus undetermined	20, 22
<i>Arietitidae</i>	2, 4
<i>Arnioceras</i>	2, 13, 15, 33
<i>densicosta</i>	15, 20, 22, 33; pl. 5
<i>semicostatum</i>	12, 15
sp	17, 33; pl. 5
<i>Asteroceras obtusum</i>	3
<i>obusum</i> Zone	12, 15
<i>attenuatus</i> , <i>Dactylioceras</i>	39
(<i>Aveyroniceras</i>) <i>italicum</i> , <i>Prodactylioceras</i>	38
B	
<i>Badouxia</i>	2, 15
<i>canadensis</i>	22, 31; pl. 2
<i>columbiae</i>	22, 31; pl. 2
Bajocian	7, 10, 11

<i>balteatum</i> , <i>Entolium</i>	17, 34
<i>bavaricum mexicanum</i> , <i>Vermiceras</i>	34
<i>binodata</i> , <i>Phymatoceras</i>	42
(<i>Brodiceras</i>) <i>tenuicostatum nodosa</i> , <i>Hildoceras</i>	42
<i>Brodieia</i>	2, 18, 42
<i>tenuicostata</i>	21, 42
<i>nodosa</i>	42; pl. 12
<i>bucklandi</i> , <i>Arietites</i>	12, 13, 17, 32; pl. 4

C	
<i>Caenisites</i>	32
<i>turneri</i> zone	12
<i>calcimontanum</i> , <i>Discamphiceras</i>	29, 30
(<i>Caloceras</i>), <i>Psiloceras</i>	29
<i>canadense</i> , <i>Psiloceras</i>	15, 31
<i>canadensis</i> , <i>Badouxia</i>	22, 31; pl. 2
Cape Simpson area	12
<i>carlottense</i> , <i>Fanninoceras</i>	21, 36
<i>Catacoeloceras</i>	2, 39
<i>confectum</i>	39
<i>crassum</i>	39
sp	17, 39; pl. 12
<i>charlesi</i> , <i>Coroniceras</i> (<i>Paracorniceras</i>)	32
<i>Charmasseiceras</i>	2, 13, 15
<i>charmassei</i>	32
<i>marmoreum</i>	17, 22, 31; pl. 3
<i>postaurinum</i>	32
sp	17, 31; pl. 3
<i>charmassei</i> , <i>Charmasseiceras</i>	32
Chitina Valley	34
Chulitna River	12
Chulitna River area	10, 16
Chulitna Valley	2, 33
<i>Coeloceras italicum</i>	38
<i>lorioli</i>	38
<i>mucronatum</i>	39
<i>Coeloceratidae</i>	35
Coleen quadrangle	29
<i>colubriforme</i> , <i>Prodactylioceras</i>	38
<i>columbiae</i> , <i>Badouxia</i>	31; pl. 2
<i>columbiae</i> , <i>Psiloceras</i> (<i>Curviceras</i>)	31
<i>commune</i> , <i>Dactylioceras</i>	21, 38; pl. 11
<i>compactile</i> , <i>Pseudohioceras</i>	18, 19, 41
<i>compressa</i> , <i>Haugia</i>	21, 22, 43; pl. 12
<i>confectum</i> , <i>Catacoeloceras</i>	39
<i>coregonense</i> , <i>Paracaloceras</i>	33
<i>Coroniceras haueri</i>	32
<i>multicostatum</i>	32
sp	20, 22
sp. A	22, 32; pl. 4
sp. B	17, 32; pl. 4
sp. C	13, 17, 32; pl. 4
(<i>Paracorniceras</i>)	2, 15
<i>charlesi</i>	32
sp	22, 32; pl. 4
<i>crassibundum</i> , <i>Dactylioceras</i>	38
<i>crassiusculosum</i> , <i>Dactylioceras</i>	38
<i>crassum</i> , <i>Catacoeloceras</i>	39
<i>crucilobatum</i> , <i>Cruciloboceras</i>	17, 20, 34; pl. 5
<i>Cruciloboceras</i>	2, 15, 17, 34, 36
<i>crucilobatum</i>	17, 20, 34; pl. 5
<i>densinodulum</i>	21, 34; pl. 7
<i>densinodum</i>	35
<i>muticum</i>	21, 35; pl. 7
<i>pacificum</i>	18, 20, 35; pl. 8
<i>submuticum</i>	21, 35; pl. 7
sp	21
(<i>Curviceras</i>) <i>columbiae</i> , <i>Psiloceras</i>	31

<i>curviornatum</i> , <i>Waehneroceras</i>	30
--	----

D	
<i>Dactylioceras</i>	2, 18, 38, 39
<i>attenuatus</i>	39
<i>commune</i>	21, 38; pl. 11
<i>crassibundum</i>	38
<i>crassiusculosum</i>	38
<i>kanense</i>	39
sp	17
(<i>Orthodactylites</i>)	2, 39
<i>directum</i>	19, 39; pl. 11
<i>kanense</i>	17, 21, 39; pl. 11
<i>Dactylioceratidae</i>	4, 38
<i>dayiceroides</i> , <i>Uptonia</i>	18, 20, 36; pl. 9
DeLong Mountains	5, 18, 37
<i>densicosta</i> , <i>Arnioceras</i>	15, 20, 22, 33; pl. 5
<i>densinodulum</i> , <i>Cruciloboceras</i>	21, 34; pl. 7
<i>densinodum</i> , <i>Cruciloboceras</i>	35
<i>Deroceras subtriangulare</i>	35
<i>directum</i> , <i>Dactylioceras</i> (<i>Orthodactylites</i>)	19, 39; pl. 11
<i>Discamphiceras</i>	2, 12, 29
<i>calcimontanum</i>	29, 30
<i>kammerkahrense</i>	29
<i>toxophorum</i>	22, 29; pl. 1
sp	20, 29; pl. 1
<i>domarense</i> , <i>Arietoceras</i>	18, 20, 21, 39; pl. 10
<i>donar</i> , <i>Schlotheimia</i>	31
<i>dumortieri</i> , <i>Haugia</i>	42

E	
<i>Echioceras</i>	34
<i>rarecostatum</i>	4, 12, 15, 34, 35
<i>Echioceratidae</i>	4
<i>Eleganticeras</i>	41
sp	19, 41; pl. 11
<i>Elegantuliceras</i>	2
<i>elegantulum</i>	41
<i>elegantulum</i> , <i>Elegantuliceras</i>	41
<i>elicitum</i> , <i>Paltechioceras</i> (<i>Paltechioceras</i>)	34
<i>engelhardti</i> , <i>Amaltheus</i> (<i>Pseudoamaltheus</i>)	17, 38
<i>Entolium balteatum</i>	17, 34
<i>semiplicatum</i>	16, 17, 34
<i>Eoderoceratidae</i>	2, 34
Etivluk River	17
<i>exaratum</i> , <i>Harpoceras</i>	18, 19, 21, 40; pl. 11
<i>extracostatum</i> , <i>Aegoceras</i>	30

F	
<i>fallaciosum</i> , <i>Grammoceras</i>	42
<i>Fanninoceras</i>	2, 36
<i>carlottense</i>	21, 36
<i>kunae</i>	21, 36; pl. 7
<i>fontanellense</i> , <i>Fontanelliceras</i>	18, 21, 40; pl. 11
<i>Fontanelliceras</i>	2, 18, 40
<i>fontanellense</i>	18, 21, 40; pl. 11
(<i>Franziceras</i>), <i>Psiloceras</i>	2, 7, 12, 29, 31
<i>ruidum</i> , <i>Psiloceras</i>	12, 19, 22, 29; pl. 1
sp., <i>Psiloceras</i>	17, 19, 29; pl. 1
<i>fucini</i> , <i>Prodactylioceras italicum</i>	20, 38; pl. 10

G	
Glenn Shale	6, 29, 37
<i>grammicus</i> , <i>Macrogrammites</i>	30

<i>Grammoceras</i>	2, 18, 42
<i>fallaciosum</i>	42
sp	21, 42; pl. 11
<i>Grammoceratinae</i>	42
<i>grandis</i> , <i>Haugia</i>	21, 22, 42; pl. 12

H

<i>Hammatoceratidae</i>	2, 42
<i>harbledownense</i> , <i>Paltehioceras</i>	15, 17, 20, 21, 34; pl. 4
<i>harbledownensis</i> , <i>Melanhippites</i>	34
<i>Harpoceras</i>	2, 18, 40
<i>exaratum</i>	18, 19, 21, 40; pl. 11
<i>variabile</i>	43
<i>Harpoceratinae</i>	40
<i>huxeri</i> , <i>Coroniceras</i>	32
<i>Haugia</i>	2, 11, 18, 42
<i>compressa</i>	21, 22, 43; pl. 12
<i>dumortieri</i>	42
<i>grandis</i>	21, 22, 42; pl. 12
<i>japonica</i>	43
<i>variabilis</i>	21, 43; pl. 12
Zone	18
sp	21
Healy (A-6) quadrangle	10, 31, 33, 34
Hettangian	7, 11, 15, 16, 28
<i>ammonites</i>	11
<i>Hildoceras</i> (<i>Brodiceras</i>) <i>tenuicostatum nodosa</i>	42
<i>Hildoceratidae</i>	2, 4, 39
Howard Pass quadrangle	17

I

<i>iber</i> Zone, <i>Tragophylloceras</i>	18, 35, 37, 38
Ignek Creek	18
Ipsavik River	5, 16, 17
<i>italicum</i> , <i>Coeloceras</i>	38
<i>Prodactylioceras</i>	18, 38
<i>Prodactylioceras italicum</i>	20, 38; pl. 10
<i>Prodactylioceras</i> (<i>Aveyronicer</i>)	38
<i>fucini</i> , <i>Prodactylioceras</i>	20, 38; pl. 10
<i>italicum</i> , <i>Prodactylioceras</i>	20, 38; pl. 10
Itkillik River	5

J

<i>jamesoni</i> , <i>Uptonia</i>	16, 19, 36; pl. 9
<i>japonica</i> , <i>Haugia</i>	43
<i>jeltzkyi</i> , <i>Arctoastroceras</i>	3, 16, 22, 31, 33; pl. 5

K

<i>kammerkahrense</i> , <i>Discamphiceras</i>	29
<i>kanense</i> , <i>Dactylioceras</i>	39
<i>Dactylioceras</i> (<i>Orthodactylites</i>)	17, 21, 39; pl. 11
Kavik River	5, 16
Kenai	33
Kenai Peninsula	10, 12, 29, 32
Kialagvik Formation	11, 42, 43
Kingak Shale,	5, 29, 31, 32, 33, 37, 39, 41
<i>kunae</i> , <i>Fanninoceras</i>	21, 36; pl. 7

L

<i>Laqueoceras</i>	2, 12, 30
<i>sublaqueus</i>	22, 30; pl. 3
<i>Leptaleoceras</i>	2, 18, 40;
<i>pseudoradians</i>	18, 21, 40; pl. 11
<i>liasicus</i> , Zone <i>Alsatis</i>	12
Lisburne Ridge	17
<i>lorioli</i> , <i>Coeloceras</i>	38
Lubbe Creek Formation	7, 18, 40
<i>lythense</i> , <i>Pseudolioceras</i>	18, 19, 41
<i>Lytoceras</i>	2
sp	22

M

<i>maclintocki</i> , <i>Pseudolioceras</i>	41
<i>Macrogrammites grammicus</i>	30
<i>margaritatus</i> , <i>Amaltheus</i>	17, 19, 37; pl. 10

<i>marmoreum</i> , <i>Charmasseiceras</i>	17, 22, 31; pl. 3
Matanuska River Valley	7, 9
McCarthy Formation	7, 17, 18, 28, 30, 33
34, 35, 36, 37, 38, 40	
<i>Melanhippites harbledownensis</i>	34
<i>mexicanum</i> , <i>Vermiceras bavaricum</i>	34
<i>Microderoceras</i>	34
<i>mitis</i> , <i>Orthodactylites</i>	39
<i>montana</i> , <i>Schlotheimia</i>	31
<i>mucronatum</i> , <i>Coeloceras</i>	39
<i>multicostatum</i> , <i>Coroniceras</i>	32
<i>Vermiceras</i>	15
<i>muticum</i> , <i>Cruciloboceras</i>	21, 35; pl. 7

N

Nelchina	9
Nelchina area	7
<i>Nodicoceras</i>	39
<i>nodosa</i> , <i>Brodieia tenuicostata</i>	42; pl. 12
<i>Hildoceras</i> (<i>Brodiceras</i>) <i>tenuicostatum</i>	42
North Fork of the American River	17
<i>nudus</i> , <i>Amaltheus</i>	37

O

<i>obtusum</i> , <i>Asteroceras</i>	3
Old Rampart	6, 11, 17
(<i>Orthocheioceras</i>), <i>Paltehioceras</i>	2, 4, 34
<i>radiatum</i> , <i>Paltehioceras</i>	34
sp., <i>Paltehioceras</i>	34; pl. 3
<i>Orthodactylites directum</i>	19, 39
<i>mitis</i>	39
(<i>Orthodactylites</i>), <i>Dactylioceras</i>	2
<i>directum</i> , <i>Dactylioceras</i>	19, 39; pl. 11
<i>kanense</i> , <i>Dactylioceras</i>	17, 21, 39; pl. 11
Otapiria	17
<i>tailleuri</i>	5, 16, 17
<i>Oxyntoceras oxyntotum</i>	3, 16
<i>oxyntotum</i> Zone	12
<i>Oxyntoceratidae</i>	36
<i>oxyntotum</i> , <i>Oxyntoceras</i>	3, 16

P

<i>pacificum</i> , <i>Cruciloboceras</i>	18, 20, 35; pl. 8
<i>Paltarpites</i>	41
<i>argutus</i>	41
<i>paltus</i>	41
<i>Paltehioceras</i>	16, 17, 34
<i>aplanatum</i>	34
<i>harbledownense</i>	15, 17, 20, 21, 34; pl. 4
sp	20
(<i>Orthocheioceras</i>)	2, 4, 34
<i>radiatum</i>	34
sp	34; pl. 3
(<i>Paltehioceras</i>) <i>elicitum</i>	34
(<i>Paltehioceras</i>) <i>elicitum</i> , <i>Paltehioceras</i>	34
<i>paltum</i> , <i>Protogrammoceras</i>	18, 21, 41; pl. 12
<i>paltus</i> , <i>Paltarpites</i>	41
<i>Paracaloceras</i>	2, 15
<i>coregonense</i>	33
<i>rursicostatum</i>	13, 22, 33; pl. 6
(<i>Paracoronicer</i>) <i>Coroniceras</i>	2, 15
Parson Bay	17
<i>Phymatoceras</i>	2, 42
<i>binodata</i>	42
<i>tumefacta</i>	42
sp	21, 42; pl. 12
<i>planorbis</i> , <i>Psiloceras</i>	20, 22, 28; pl. 1
<i>Platyleuroceras</i> sp	35
<i>Pleuroceras spinatum</i> Zone	38
<i>plicatum</i> , <i>Psiloceras</i> (<i>Psiloceras</i>)	29
<i>Pliensbachian</i>	2, 4, 5, 7, 18
<i>ammonites</i>	17
Point Barrow area	12, 37
<i>Polymorphitidae</i>	2, 4, 36
<i>portlocki</i> , <i>Aegoceras</i>	30
<i>Schlotheimia</i> (<i>Waehneroceras</i>)	30
<i>Waehneroceras</i>	22, 30; pl. 2
<i>postaurium</i> , <i>Charmasseiceras</i>	32

<i>Prodactylioceras</i>	2, 38
<i>colubriforme</i>	38
<i>italicum</i>	18, 38
<i>fucini</i>	20, 38; pl. 10
<i>italicum</i>	20, 38; pl. 10
sp	20
(<i>Aveyronicer</i>) <i>italicum</i>	38
<i>Protogrammoceras</i>	2, 18, 41
<i>argutum</i>	18, 21, 41; pl. 11
<i>paltum</i>	18, 21, 41; pl. 12
(<i>Pseudoamalthus</i>) <i>engelhardti</i> , <i>Amaltheus</i>	17, 38
<i>Pseudolioceras</i>	2, 18, 41
<i>compactile</i>	18, 19, 41
<i>lythense</i>	18, 19, 41
<i>maclintocki</i>	41
<i>whiteavesi</i>	11
sp	21, 22, 41; pl. 11, 12
<i>pseudoradians</i> , <i>Ammonites</i>	40
<i>Arietoceras</i>	40
<i>Leptaleoceras</i>	18, 21, 40; pl. 11
<i>Psiloceras canadense</i>	15, 31
<i>planorbis</i>	20, 22, 28; pl. 1
<i>plicatum</i>	29
<i>Psiloceras</i> (<i>Psiloceras</i>)	6, 12, 28, 29, 31
sp	19
(<i>Caloceras</i>)	29
(<i>Curvicer</i>) <i>columbiae</i>	31
(<i>Franzicer</i>)	7, 12, 31
<i>ruidum</i>	12, 22, 29; pl. 1
sp	17, 19, 29; pl. 1
(<i>Psiloceras</i>), <i>Psiloceras</i>	6, 12, 28, 29, 31
<i>Psiloceratidae</i>	2, 28
Puale Bay	2, 11, 12, 13, 18, 28
29, 30, 31, 33, 42, 43	
Puale Bay area	33

Q

Queen Charlotte Islands	2, 35, 36, 37
-------------------------------	---------------

R

<i>radiatum</i> , <i>Paltehioceras</i> (<i>Orthocheioceras</i>)	34
<i>Radstockiceras</i>	2
Rainy Pass	10
<i>rarsicostatum</i> zone, <i>Echioceras</i>	4, 13, 15, 34, 35
Red Hill	18
<i>retroscosta</i> , <i>Arietoceras</i>	40
<i>ruidum</i> , <i>Psiloceras</i> (<i>Franzicer</i>)	12, 19, 22, 29; pl. 2
<i>rursicostatum</i> , <i>Paracaloceras</i>	13, 22, 33; pl. 6

S

Sadlerochit River	18
Sagavanirktok River	4, 5
<i>Schlotheimia</i>	2, 12, 31
<i>angulata</i> Zone	12
<i>donar</i>	31
<i>montana</i>	31
sp	22, 31; pl. 2
(<i>Waehneroceras</i>) <i>portlocki</i>	30
<i>Schlotheimiidae</i>	2, 4
Seldovia	13
Seldovia area	12, 15, 29, 32, 33
Seldovia Bay	10
<i>semicostatum</i> , <i>Arnioceras</i>	12, 15
<i>semiplicatum</i> , <i>Entolium</i>	16, 17, 34
Shellabarger and Rainy Pass areas	10
Shellabarger Pass	10
<i>simile</i> , <i>Aegasteroceras</i>	3
Simpson test well No. 1	37
Sinemurian	4, 5, 7, 10, 11, 12, 13
<i>ammonites</i>	12
South Barrow No. 12 well	29
South Barrow test well No. 12	31, 32, 33
South Barrow test well No. 3	32, 37, 38, 39
Spike Mountain	6
<i>spinatum</i> , <i>Pleuroceras</i>	38
<i>stokesi</i> , <i>Amaltheus</i>	7, 17, 18, 19, 21, 37; pl. 10
<i>sublaqueus</i> , <i>Laqueoceras</i>	22, 30; pl. 3
<i>submuticum</i> , <i>Cruciloboceras</i>	21, 35; pl. 7

subtriangulare, Apoderoceras 18, 21, 35; pl. 8
Deroceras 35

T

tailleuri, Otapiria 5, 16, 17
 Talkeetna (B-6) quadrangle 10
 Talkeetna (C-6) quadrangle 10, 32
 Talkeetna Formation 8, 10, 33, 34, 35, 36,
 37, 38, 39, 40, 41, 42, 43
 Talkeetna Mountains 2, 7, 8, 12, 15, 17,
 18, 33, 34, 35, 36, 37,
 38, 39, 40, 41, 42, 43
tenerum, Aegoceras 30
Waehneroceras 22, 30, 31; pl. 2
tenuicostata, Brodieia 21, 42
nodosa, Brodieia 42; pl. 12
Hildoceras
 (*Brodiceras*) 42
 Toarcian 4, 5, 7, 10, 11, 18
 ammonites 18
toxophorum, Discamphiceras 22, 29; pl. 1
Tragophylloceras
ibex Zone 18, 35, 37, 38

Triassic 11
Tropidoceras 2, 37
actaeon 18, 20, 35, 37; pl. 8
tumefacta, Phymatoceras 42
turneri, Caenisisites 12

U

Uptonia 2, 5, 17, 18, 36, 37, 38
dayiceroides 18, 20, 36; pl. 9
jamesoni 16, 19 36; pl. 9
 Zone 17, 18, 35, 37, 38
 sp 17, 19, 36; pl. 9
 sp. A 20, 36; pl. 8
 sp. B 21, 37; pl. 9

V

variabile, Harpoceras 43
variabilis, Ammonites 43
Haugia 21, 43; pl. 12
Vermiceras bavaricum mexicanum 34
multicostatum 15

W

Waehneroceras 2, 12, 28, 29, 30
curviornatum 30
portlocki 22, 30; pl. 2
tenerum 22, 30; pl. 2
 sp 17, 30; pl. 2
 (*Waehneroceras*) *portlocki, Schlothemia* 30
whiteavesi, Pseudolioceras 11
 Wide Bay 30
 Wide Bay areas 12
 Wide Bay test well No. 1 30
 Wrangell Mountains 7, 12, 13, 15, 17, 18, 28,
 30, 33, 34, 35, 36, 37, 38, 40

X

Xiphoceratinae 34

Y

Yakutat District 17
 Yakutat D-4 quadrangle 37

PLATES 1-12

Contact photographs of the plates in this report are available, at
cost, from U.S. Geological Survey Library, Federal Center,
Denver, Colorado 80225

PLATE 1

[Figures natural size unless otherwise indicated]

FIGURES 1, 2. *Psiloceras* cf. *P. planorbis* (J. de C. Sowerby) (p. 28).

1. Specimen, USNM 247950, from USGS Mesozoic loc. 29891.

2. Specimen, USNM 247951, from USGS Mesozoic loc. M1738.

3, 4, 8-10. *Discamphiceras* cf. *D. toxophorum* Waehner (p. 29).

3. Specimen, USNM 247960, from Richfield Oil Co. loc. 3002.

4. Specimen, USNM 247958, from USGS Mesozoic loc. 3110.

8. Specimen, USNM 247961, from Richfield Oil Co. loc. 3002.

9, 10. Crushed fragment and rubber imprint of external mold of one specimen, USNM 247959, from USGS Mesozoic loc. 25694.

5-7. *Discamphiceras* sp. (p. 29).

Rubber imprints of three external molds, USNM 247962, from USGS Mesozoic loc. 29890.

11, 15-17. *Psiloceras* (*Franziceras*) sp. ind. (p. 29).

11, 15, 16. Ventral and lateral views of specimen, USNM 247956, from South Barrow test well No. 12 at depth of 2,170.5 ft (661.4 m).

17. Specimen, USNM 247957 from USGS Mesozoic loc. 29742.


12-14, 18-24. *Psiloceras* (*Franziceras*) cf. *P. (F.) ruidum* Buckman (p. 29).

12. Rubber imprint of external mold ($\times 2$) of a specimen, USNM 247952, from USGS Mesozoic loc. 29738.

13, 14. Rubber imprint of external mold ($\times 2$), specimen, USNM 247954, from USGS Mesozoic loc. 29738.

18-20. Rubber imprint of external mold and views of internal mold of a specimen, USNM 247955, from USGS Mesozoic loc. 21242.

21-24. Rubber imprints ($\times 2$) and suture line ($\times 4$) of one specimen, USNM 247953, from USGS Mesozoic loc. 29738.



PSILOCERAS, *P. (FRANZICERAS)*, AND *DISCAMPHICERAS*

PLATE 2

[Figures natural size unless otherwise indicated]

FIGURES 1–6. *Waehneroceras* cf. *W. tenerum* (Neumayr) (p. 30).

1–3. Three specimens, USNM 247964, from Richfield Oil Co. loc. 3002. Figure 1 is a rubber imprint of an external mold.

4, 5. Specimen, USNM 247965, from USGS Mesozoic loc. 19803. Figure 5 below fracture is an external mold.

6. Specimen, USNM 247966, from USGS Mesozoic loc. 25694.

7, 10–15. *Waehneroceras* cf. *W. portlocki* (Wright) (p. 30).

7. Specimen, USNM 247971, from USGS Mesozoic loc. 25694.

10. Specimen, USNM 247969, from USGS Mesozoic loc. 12075.

11. Specimen, USNM 247970, from USGS Mesozoic loc. 12394.

12. Specimen, USNM 247967, from Richfield Oil Co. loc. 3002.

13. Specimen, USNM 247968, from USGS Mesozoic loc. 3110.

14, 15. Two specimens, USNM 247972, from Richfield Oil Co., Wide Bay test No. 1 at depth of 2,235–2,236 ft (681 m).

8, 9. *Waehneroceras*? sp. (p. 30).

Ventral and lateral views ($\times 2$) of specimen, USNM 247973, from the South Barrow test well No. 12 at depth of 2,181.5 ft (665 m).

16, 17. *Schlotheimia* sp. (p. 31).

Specimen, USNM 247974, from USGS Mesozoic loc. 10820.

18–21, 24–28. *Badouxia canadensis* (Frebold) (p. 31)

18, 19. Hypotype, USNM 247977, from Richfield Oil Co. loc. 1240.

20, 21. Hypotype, USNM 247976, from USGS Mesozoic loc. 31266.

24, 28. Lateral views of hypotype, USNM 247975, from USGS Mesozoic loc. 31372. Figure 24 is a rubber imprint of the external mold.

25–27. Specimen, USNM 247975, from USGS Mesozoic loc. 31266.

22, 23. *Badouxia columbiae* (Frebold) (p. 31).

Hypotype, USNM 247978, from USGS Mesozoic loc. 31264.

22, 23. Hypotype, USNM 247978, from USGS Mesozoic loc. 31264.



WAEHNEROCERAS, SCHLOTHEIMIA, AND BADOUXIA

PLATE 3

[Figures natural size unless otherwise indicated]

FIGURES 1-3. *Charmasseiceras* cf. *C. marmoreum* (Oppel) (p. 31).

Lateral views of three specimens, USNM 247979, from the South Barrow test well No. 12 at depth of 2,061.5 ft (628 m).

4-10. *Charmasseiceras* sp. (p. 31).

4-6, 9, 10. Suture line ($\times 4$) and lateral and ventral views of same specimen ($\times 2$ and $\times 6$), USNM 247981, from the South Barrow test well No. 3 at depth of 2,412 ft (735 m).

7, 8. Lateral and ventral views of specimen, USNM 247980, from the South Barrow test well No. 12 at depth of 2,056 ft (627 m).

11, 12. *Paltechioceras* (*Orthechioceras*?) sp. (p. 34).

Rubber imprint of external mold (fig. 11) and ventral view of specimen, USNM 248003, from USGS Mesozoic loc. 31261.

13. *Laqueoceras* cf. *L. sublaqueus* (Waehner) (p. 30).

Rubber imprint of external mold, USNM 247963, from Richfield Oil Co. loc. 3002.



CHARMASSEICERAS, PALTECHIOCERAS, AND LAQUEOCERAS

PLATE 4

[All figures are natural size]

FIGURES 1–5. *Coroniceras* sp. A (p. 32).

Three specimens, USNM 247982, from USGS Mesozoic loc. 30908. Ventral and lateral views of one specimen are shown in figures 1 and 3. Same views of another specimen are shown in figures 2 and 4.

6–10. *Coroniceras* sp. B (p. 32).

6. Rubber imprint of external mold of specimen, USNM 247983, from the South Barrow test well No. 12 at depth of 1,987.5 ft (605.8 m).

7–10. Two specimens, USNM 247984, from the South Barrow test well No. 12 at depth of 1,987.6 feet (605.8 m). Ventral and lateral views of one specimen are shown in figures 7 and 10. Same views of another specimen are shown in figures 8 and 9.

11–13. *Arietites* cf. *A. bucklandi* (Sowerby) (p. 32).

Figure 13 probably represents an inner whorl of specimen shown in figures 11 and 12, USNM 108778, from the Avak test well No. 1 at depth of 1,836 ft (560 m).

14. *Coroniceras* sp. C (p. 32).

Rubber imprint of external mold, USNM 247985, from the South Barrow test well No. 3 at depth of 2,470 ft (753 m).

15–22. *Paltechioceras* cf. *P. harbledownense* (Crickmay) (p. 34).


15, 16. Rubber imprint and external mold of specimen, USNM 248000, from USGS Mesozoic loc. 31174.

17, 22. Specimens, USNM 248001, from USGS Mesozoic loc. 30139.

18–21. Ventral and lateral views of two specimens, USNM 248002, from USGS Mesozoic loc. 28661.

23–25. *Coroniceras* (*Paracorniceras*) sp. (p. 32).

Two specimens, USNM 247986, from USGS Mesozoic loc. 31128. Figure 22 represents a rubber imprint of the inner whorls. Figures 23 and 24 show lateral and ventral views of an outer whorl.



CORONICERAS, C. (PARACORONICERAS), ARIETITES, AND PALTECHIOCERAS

PLATE 5

[Figures natural size unless otherwise indicated]

FIGURES 1–4. *Arctoasteroceras jeletzkyi* Frebold (p. 33).

1, 2. Hypotype, USNM 247999, from USGS Mesozoic loc. 31261.

3, 4. Hypotype, USNM 247998, from USGS Mesozoic loc. 31261.

5, 6, 12–15. *Arnioceras* sp. juv. (p. 33).

5, 6. Ventral and lateral views.

12, 13. Ventral and lateral views.

14, 15. Rubber imprints of two external molds showing smooth inner whorls. Specimens, USNM 247990, from the South Barrow test well No. 12 at depth of 2,056 ft (626.7 m).

7, 8. *Cruciloboceras* cf. *C. crucilobatum* Buckman (p. 34).

7. Laterally crushed small specimen ($\times 2$), USNM 248005, from USGS Mesozoic loc. 14030.


8. Rubber imprint of external mold of large specimen, USNM 248004, from USGS Mesozoic loc. 14472.

9–11, 16–24. *Arnioceras* cf. *A. densicosta* (Quenstedt) (p. 33).

9, 16, 17. Rubber imprint of inner whorls (fig. 9) and suture line ($\times 2$) of one specimen, USNM 247987, from USGS Mesozoic loc. 16229.

10, 11, 18, 19, 22–24. Five specimens, USNM 247988, from USGS Mesozoic loc. 16229. Ventral and lateral views of one specimen are shown in figures 10 and 11. Same views of another specimen are shown in figures 18 and 19. Figures 22–24 represent rubber imprints of external molds.

20, 21. Lateral and ventral view of specimen, USNM 247989, from USGS Mesozoic loc. 21237.




ARCTOASTEROCERAS, ARNIOCERAS, AND CRUCILOBICERAS

PLATE 6

[All figures are natural size]

FIGURES 1–11. *Paracaloceras rursicostatum* Frebold (p. 33).

- 1, 2. Hypotype, USNM 247991, from USGS Mesozoic loc. 12396.
- 3, 5. Hypotype, USNM 248070, from USGS Mesozoic loc. 31372.
4. Hypotype, USNM 247996, from USGS Mesozoic loc. 10820.
6. Hypotype, USNM 247997, from USGS Mesozoic loc. 10820.
- 7, 11. Hypotype, USNM 247993, from Richfield Oil Co. loc. 1241.
8. Hypotype, USNM 247995, from USGS Mesozoic loc. 31266.
9. Hypotype, USNM 247994, from USGS Mesozoic loc. 2981.
10. Hypotype, USNM 247992, from USGS Mesozoic loc. 12396.



PARACALOCERAS

PLATE 7

[All figures are natural size]

FIGURES 1-3. *Crucilobiceras* cf. *C. submuticum* (Oppel) (p. 33).

1, 2. Specimen, USNM 248008, from USGS Mesozoic loc. 6706.

3. Specimen, USNM 248009, from USGS Mesozoic loc. 28661.

4, 5. *Crucilobiceras* cf. *C. densinodulum* Buckman (p. 34).

Specimen, USNM 248006, from USGS Mesozoic loc. 28661.

6-10, 12-15. *Crucilobiceras* cf. *C. muticum* (d'Orbigny) (p. 35).

Six specimens, USNM 248007, from USGS Mesozoic loc. 28661. Figures 9 and 10 represent ventral and lateral views of smallest specimen. Figures 12 and 13 represent same views of a fairly large specimen. Figures 14 and 15 represent same views of the largest specimen.

11. *Fanninoceras kunae* McLearn (p. 36).

Hypotype, USNM 248013, from USGS Mesozoic loc. 27586.



CRUCILOBICERAS AND FANNINOCERAS

PLATE 8

[Figures natural size unless otherwise indicated]

FIGURES 1–9. *Tropidoceras actaeon* (d'Orbigny) (p. 37).

1–3. Hypotype, USNM 248021, from USGS Mesozoic loc. 28534.

4–6. Hypotype, USNM 248020, from USGS Mesozoic loc. 28534. Figure 4 is a rubber imprint of the external mold.

7. Hypotype, USNM 248022, from USGS Mesozoic loc. 28534.

8, 9. Hypotype, USNM 248023, from USGS Mesozoic loc. 28534.

10–12, 15–17. *Cruciloboceras* cf. *C. pacificum* Frebold (p. 35).

One specimen, USNM 248010, from USGS Mesozoic loc. 28534. Figures 15–17 ($\times 2$). Figure 12 is a rubber imprint of an external mold of the same specimen.


13. *Uptonia*? sp. A. (p. 36).

Specimen, USNM 248018, from USGS Mesozoic loc. 29773.

14, 18–23. *Apoderoceras* cf. *A. subtriangulare* (Young and Bird) (p. 35).

14, 18–21. Specimen, USNM 248011, from USGS Mesozoic loc. 6697. Figures 18 and 19 represent the adapical part of the middle whorl shown on figure 21. Figure 14 does not include the cross section of the outermost whorl. Figure 20 is a rubber imprint of the external mold of the same specimen.

22, 23. Lateral and ventral views of a small specimen, USNM 248012, from USGS Mesozoic loc. M6171.



TROPIDOCERAS, CRUCILOBICERAS, UPTONIA?, AND APODEROCERAS

PLATE 9

[Figures natural size unless otherwise indicated]

FIGURES 1-4, 8, 12-16. *Uptonia* cf. *U. dayiceroides* Mouterde (p. 36).

1-3. Specimens, USNM 248014, from USGS Mesozoic loc. 28675.

4, 8, 12-16. Specimens, USNM 248015, from USGS Mesozoic loc. 28671.

5-7. *Uptonia* sp. B (p. 37).


Specimen, USNM 248019, from USGS Mesozoic loc. 6697. Figures 5 and 7 are ($\times 2$).

9-11. *Uptonia?* sp. (p. 36).

Specimen, USNM 248017, from USGS Mesozoic loc. 29775.

17. *Uptonia* cf. *U. jamesoni* (J. de C. Sowerby) (p. 36).

Laterally crushed specimen, USNM 248016, from USGS Mesozoic loc. 29774.



UPTONIA

PLATE 10

[Figures natural size unless otherwise indicated]

FIGURES 1, 2, 6–15, 22. *Arieticerias* cf. *A. domarense* (Meneghini) (p. 39).

1, 2, 8–10, 15. Specimens, USNM 248033, from USGS Mesozoic loc. 28688. Ventral and lateral views of one specimen are shown in figures 1 and 2. Same views of another specimen are shown in figures 9 and 10.

6. Specimen, USNM 248034, from USGS Mesozoic loc. 29450.

7. Specimen, USNM 248036, from USGS Mesozoic loc. 27586.

11–14. Two specimens, USNM 248035, from USGS Mesozoic loc. 24107. Ventral and lateral views of one specimen are shown in figures 11 and 12. Same views of another specimen are shown in figures 13 and 14.

22. Specimen, USNM 248037, from USGS Mesozoic loc. 28531.

3. *Prodactylioceras italicum italicum* (Fucini) (p. 38).

Hypotype, USNM 248030, from USGS Mesozoic loc. 28671.

4, 5. *Prodactylioceras* cf. *P. italicum fucini* Fischer (p. 38).

4. Specimen, USNM 248032, from USGS Mesozoic loc. 28673.

5. Specimen, USNM 248031, from USGS Mesozoic loc. 28671.

16–20. *Arieticerias* cf. *A. algovianum* (Oppel) (p. 40).

Specimens, USNM 248038, from USGS Mesozoic loc. 28688. Ventral and lateral views are shown in figures 16 and 17, and also in figures 19 and 20.

21. *Arieticerias* sp. (p. 40).

Specimen, USNM 248039, from USGS Mesozoic loc. 28531.

23, 24. *Amaltheus* cf. *A. stokesi* (J. Sowerby) (p. 37).

23. Specimen, USNM 248029, from USGS Mesozoic loc. 30074.

24. Specimen, USNM 248028, from USGS Mesozoic loc. 25941.

25, 26. *Amaltheus margaritatus* (Montfort) (p. 37).


25. Hypotype, USNM 248025, from Simpson test well No. 1 at depth of 5,677 ft (1,730m).

26. Hypotype, USNM 248024, from USGS Mesozoic loc. M2441.

27, 28. *Amaltheus stokesi* (J. Sowerby) (p. 37).

27. Hypotype, USNM 248026, from USGS Mesozoic loc. 29340.

28. Hypotype, USNM 248027, from USGS Mesozoic loc. 29165.




ARIETICERAS, PRODICTYLOCERAS, ARIETICERAS, AND AMALTHEUS

PLATE 11

[Figures natural size unless otherwise indicated]

- FIGURE 1. *Pseudolioceras* sp. (p. 41).
Specimen, USNM 248052, from USGS Mesozoic loc. 19804.
- 2, 3, 8. *Dactylioceras* cf. *D. commune* (Sowerby) (p. 38).
Specimens, USNM 248047, from USGS Mesozoic loc. 24787.
- 4, 5, 9. *Dactylioceras* (*Orthodactylites*) *kanense* McLearn. (p. 39).
Hypotype, USNM 248048, from USGS Mesozoic loc. 29198. Figure 5 is a rubber imprint of an external mold.
6. *Dactylioceras* (*Orthodactylites*) cf. *D. (O.) directum* Buckman (p. 39).
Specimen, USNM 248049, from USGS Mesozoic loc. 29163.
7. *Grammoceras* sp. (p. 42).
Specimen, USNM 248053, from USGS Mesozoic loc. 25939.
- 10, 15, 16. *Eleganticeras* sp. juv. (p. 41).
Specimens ($\times 2$), USNM 248050, from USGS Mesozoic loc. 29776. Figure 16 is a ventral view of figure 15.
11. *Harpoceras* cf. *H. exaratum* (Young and Bird) (p. 40).
Specimen, USNM 248044, from USGS Mesozoic loc. 29161.
- 12, 13. *Leptaleoceras* cf. *L. pseudoradians* (Reynes) (p. 40).
Specimen, USNM 248040, from USGS Mesozoic loc. 29450.
14. *Protogrammoceras* cf. *P. argutum* (Buckman) (p. 41).
Specimen, USNM 248046, from USGS Mesozoic loc. 25941.
- 17-23. *Fontanelliceras* cf. *F. fontanellense* (Gemmellaro) (p. 40).
7, 18, 23. Specimen, USNM 248043, from USGS Mesozoic loc. 29450.
19, 20. Specimen, USNM 248041, from USGS Mesozoic loc. 25941.
21, 22. Specimen, USNM 248042, from USGS Mesozoic loc. 27586.



PSEUDOLIOCERAS, DACTYLIOCERAS, D. (ORTHODACTYLITES), GRAMMOCERAS, ELEGANTICERAS, HARPOCERAS, LEPTALEOCERAS, PROTOGRAMMOCERAS, AND FONTANELLICERAS

PLATE 12

[All figures are natural size]

FIGURES 1, 2, 5. *Haugia* cf. *H. variabilis* (d'Orbigny) (p. 43).

Specimens, USNM 248059, from USGS Mesozoic loc. 25318.

3, 7, 9. *Haugia* cf. *H. compressa* Buckman (p. 43).

3, 9. Specimens, USNM 248060, from USGS Mesozoic loc. 24114.

7. Specimen, USNM 248061, from USGS Mesozoic loc. 19804.

6. *Catacoeloceras?* sp. juv. (p. 39).

Specimen, USNM 108758, from South Barrow No. 3 test well at depth of 2,063 ft (629 m).

4, 10, 15. *Haugia* cf. *H. grandis* Buckman (p. 42).

Lateral views of three specimens, USNM 248058, from USGS Mesozoic loc. 24114. Adoral part of figure 4 is an internal mold.

8. *Brodieia* cf. *B. tenuicostata* var. *nodosa* Jaworski (p. 42).

Specimen, USNM 248057, from USGS Mesozoic loc. 25940.

11, 12. *Protogrammoceras* cf. *P. paltum* Buckman (p. 41).

Specimen, USNM 248045, from USGS Mesozoic loc. 25941. Figure 12 is a rubber imprint of an external mold.

13. *Pseudolioceras* sp. (p. 41).


Specimen, USNM 248051, from USGS Mesozoic loc. 25317.

14, 16–19. *Phymatoceras?* sp. (p. 42).

14, 16. Specimen, USNM 248055, from USGS Mesozoic loc. 25317.

17, 19. Specimen, USNM 248054, from USGS Mesozoic loc. 25317.

18. Specimen, USNM 248056, from USGS Mesozoic loc. 25342.



*HAUGIA, CATACOELOCERAS, BRODIEIA, PROTOGRAMMOCERAS,
PSEUDOLIOCERAS, AND PHYMATOCERAS?*