

Visualizing the Geology of Lake Trout Spawning Sites: Northern Lake Michigan

By
Peter Dartnell,¹ Peter Barnes,¹ James V. Gardner,² and Kristen Lee¹
2004



DISCUSSION

Geologists and biologists are working together to understand the links between lake floor geology (composition and shape) and the distribution of lake trout throughout their life cycle. Lake floor geology is one of the main factors determining where lake trout spawn, feed, and hide. In support of ongoing research to study Lake Michigan trout habitats, the U.S. Geological Survey in cooperation with the U.S. Army Corps of Engineers mapped the morphology of principle lake trout spawning sites (Dawson and others, 1997). Using the Army Corps of Engineer's SHOALS airborne lidar (Light Detection and Ranging) system we mapped six regions in Northern Lake Michigan (fig. 1) in order to identify ideal spawning regions composed of shallow, clean, gravel/cobble substrate, adjacent to deeper water (Barnes and others, 2003).

Lidar mapping systems, which use laser pulses to measure water depths from an airplane, are now available to map the nearshore lake morphology at meter-scale detail. Maps generated from the bathymetric data are used to define regions with smooth homogeneous substrate, regions with higher relief, and mixed regions with both smooth and rough relief. This morphologic information combined with sediment samples and direct bottom observations enable geologists to map areas with rougher relief composed of rock outcrop, boulders, and cobbles, as well as smooth regions covered with sand or mud. This information helps biologists, fishery managers, and ecologists visualize the lake floor in significant detail which promotes better fishery management, species protection, and habitat identification.

These posters present the maps and discuss the geology of the six lake trout spawning sites mapped by the lidar system. Where the mapping approached land, aerial photography of the land is combined with the bathymetric data to help visualize the scale of the offshore features. Map and perspective views of Boulder Reef, Hog Island Reef, and Little Traverse Bay are shown on sheet 1, whereas map and perspective views of Trout and High Island Shoal, Gull Island Reef, and Dablia Shoal are shown on sheet 2. Additional information, bathymetric data, imagery, and metadata are available online at <http://geopubs.wr.usgs.gov/open-file/0f03-120/>.




Figure 1. Location map of six sites mapped with the U.S. Army Corps of Engineer's SHOALS lidar system along with U.S. Geological Survey 30-m DEM topography and regional shaded bathymetry developed from Holcombe and others, 1996.

REFERENCES

- Barnes, P., Fleisher, G., Gardner, J.V., and Lee, K., 2003. Bathymetry and selected perspective views of 6 reef and coastal areas in Northern Lake Michigan. U.S. Geological Survey Open-File Report OF03-120. [Available online at <http://geopubs.wr.usgs.gov/open-file/0f03-120/>.]
- Dawson, K.A., Eshenroder, R.L., Holy, M.E., and Ward, C., 1997. Quantification of historic lake trout (*Salvelinus namaycush*) spawning aggregations in Lake Michigan. *Canadian Journal of Fish Aquatic Science*, vol. 54, no. 10, p. 2290-2302.
- Edsall, T.A., Poe, T.P., Nester, R.T., and Brown, C.L., 1989. Side-scan sonar mapping of lake trout spawning habitat in Northern Lake Michigan. *North American Journal of Fisheries Management*, vol. 9, p. 269-279.
- Holcombe, T.L., Reid, D.F., Virden, W.T., Niemeyer, T.C., De la Sierra, R., and Divins, D.L., 1996. Bathymetry of Lake Michigan. U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Geophysical Data Center, Boulder, Colorado. Report MGG-11 [A color poster with descriptive text and digital data available on CD-ROM].
- Somers, L.H., 1968. Preliminary report on geological studies in northern Lake Michigan using underwater observation techniques. *International Association for Great Lakes Research, Proceedings, 11th conference on Great Lakes Research*, vol. 239-244.

Lake floor images generated from SHOALS lidar data below 178.80 m (ref. IGLD 85). Data acquired on a four-meter grid in August 2001 were collected and processed by J.E. Chance & Associates under contract to the Army Corps of Engineers and the U.S. Geological Survey.

Land imagery created from Digital Orthophoto Quadrangles (DOQ) modified from http://www.michigan.gov/dnr/0,1607,7-153-10371_14546--,00.html

Data, imagery, and metadata available online at <http://geopubs.wr.usgs.gov/open-file/0f03-120/>. Data collection field activity metadata URL: <http://wdr.usgs.gov/info/bnk/b/101/m/m/1-01-1m.meta.html>. PDF versions of these posters are available at <http://pubs.usgs.gov/imap/2930/>

Author Affiliation
¹U.S. Geological Survey, Menlo Park, California
²University of New Hampshire, Durham, New Hampshire




Figure 2.1. Color-coded shaded relief bathymetry of Boulder Reef. The rough relief in the southern and shallowest portion of the reef (6 to 22 m water depth) is covered with gravel, cobbles, and boulders, with rare patches of sand (Somers, 1968; Edsall and others, 1989; and video data). This coarse bottom material is roughly organized into highs and lows less than a meter high. The arcuate ridge at the south end of the reef (A) is likely a remnant depositional glacial feature from a small lobe of ice originating from the north. The coarse-texture material extends beyond the depths of the survey to the west and south, but transitions to a sand substrate to the east (B) and north (video observations by Greg Kennedy, U.S. Geological Survey, Ann Arbor, Michigan, personal commun., 2002). The strong east-west lineations at depths of 15-20 m in the north half of the image (C) could be related to glacial movement but this is transverse to movement suggested by the ridge to the south or to older glacial events. Sediment samples show that these lineations are draped with clean sand. Note: The faint N-S lineations seen in the central part of the image are artifacts of the data gathering flight lines.




Figure 2.2. Oblique view of the Boulder Reef area looking southwest. This view emphasizes the east-west lineations (C) and smooth morphology of the sand draped surface north of Boulder Reef. The vertical exaggeration is 4x and the distance across the bottom of the image is about 4.0 km.

BOULDER REEF




Figure 2.3. Oblique view of Boulder Reef looking northwest. The relatively steep slopes of the southern end of the reef and the different rough morphologic character of the cobble and gravel substrate fill the foreground. A small depositional apron of sand (B) seen in underwater video collected at this site is forming off the southeast part of the reef. The vertical exaggeration is 4x and the distance across the bottom of the image is about 3.5 km.





Figure 3.1. Color-coded shaded relief bathymetry of the area southeast of Hog Island including Hog Island Reef (A). The reef is about 1200 by 1400 m and ranges in water depth from 2 to 18 m. Underwater observations on the reef report flat-lying outcrops of limestone (Somers, 1968). This mapped area is marked by strong northwest-southeast ridges and lineations (B) about a meter high and a few meters wide. An indistinct northeast-southwest lineation is also present, especially in the northeastern quadrant of the image (C). Draped over this mapped region are patches of smoother relief, possibly sand, which forms north-south bedforms (D).




Figure 3.2. Oblique view looking west toward Hog Island. The pronounced northwest-southeast trending lineations in the center of the mapped area (B) die out to the north and appear buried by a coarse morphology that looks similar to the boulder and cobble areas seen at Boulder Reef (sheet 1, fig. 2.1) and Gull Island Reef (sheet 2, fig. 6.1). The vertical exaggeration is 4x and the distance across the bottom of the image is about 3.5 km.

HOG ISLAND REEF




Figure 3.3. Oblique view looking north at Hog Island Reef (A). North-south trending bedforms (D) are associated with patches of modern lake sediment composed of sand aligned east-west. The fine and linearly continuous morphology of glacial or bedrock outcrop underlie both the bedform and sand patches. The vertical exaggeration is 4x and the distance across the bottom of the image is about 3.5 km.

Figure 4.4. Oblique view of the eastern portion of the Little Traverse Bay data set looking west along the coast. This view illustrates the nearshore, rough relief of outcrops and coarser substrate and the offshore, sediment covered substrate, separated by the shore-parallel trench (C). The blue rectangular areas close to shore are bathymetry data gaps and show the water column from the aerial photographs. The vertical exaggeration is 4x and the distance across the bottom of the image is about 800 meters.





Figure 4.3. Oblique view looking toward the east over Nine Mile Point. This view illustrates the onshore (G) and offshore expression of a glacially created drumlin trending northwest-southeast. The view also shows the sharp trough-like boundary (B) between the offshore sediment covered region and the nearshore, irregular relief of coarser material and outcrop. The vertical exaggeration is 4x and the distance across the bottom of the image is about 1.0 km.



LITTLE TRAVERSE BAY




Figure 4.2. Oblique view looking south towards Big Rock Point. Rough irregular morphology (E) composed of bedrock outcrop extends from the coast offshore to the edge of the survey, except in the left foreground (F), which appears depositional. The vertical exaggeration is 4x and the distance across the bottom of the image is about 1.3 km.




Figure 4.1. Color-coded shaded relief bathymetry of the nearshore region along the south coast of Little Traverse Bay (fig. 1). Onshore aerial photography and Digital Elevation Models (DEMs) show that the surface morphology of this region is marked by glacial drumlins aligned northwest-southeast. Nearshore, mainly in water depths from 0 to 10 m (A), the morphology is irregular and blocky, suggesting outcrops and coarse substrate. Offshore of the outcrops, a shore-parallel trench (B), 1-3 m deep, separates the outcrops from the deeper sediment covered region (C).