Figure 1. Thermal maturity map of Devonian shale in the Illinois, Michigan, and Appalachian basins. Abbreviations are as follows: %R, vitrinite reflectance reported as mean percent reflectance in oil; CAI, conodont color alteration index; SCI, spore color index. The basin outlines are compiled from Thomas and others (1989) and Swzezy (2002, 2008, 2009). In the Illinois basin, the thermal maturity values are based on vitrinite reflectance (VR) measurements from the basal interval of the Middle Devonian to Lower Mississippian New Albany Shale (Barrows and Cluff, 1984; Nuccio and Hatch, 1996; Strapoc and others, 2010). In the Michigan basin, the thermal maturity values are based on VR measurements from the basal interval of the Upper Devonian Antrim Shale (Moyer, 1982; Cercone, 1984; Cercone and Pollack, 1991; Wang and others, 2010). In the Appalachian basin, the thermal maturity values are based on VR measurements from Devonian shale and conodont CAI measurements from Devonian limestone that is stratigraphically near the Middle Devonian Marcellus Shale and correlative units (Harris and others, 1978; Harris, 1979; Bayer, 1982; Repetski and others, 2008). These values from Middle Devonian strata in the Appalachian basin are generally consistent with VR values from the Upper Devonian West Falls Formation (Gerlach and Cercone, 1993; Curtis and Faure, 1997) and from the Upper Devonian Ohio Shale (Rimmer and others, 1993; Curtis and Faure, 1997).

Figure 2. Diagram that shows the relations among various thermal maturity indicators including conodont CAI and %R, and associated zones of hydrocarbon generation (modified from Fisher and others, 1980; Repetski and others, 2008). The zone boundaries are approximate and are based on oil generation from Type II kerogen. The color scheme corresponds to the thermal maturity values in Figure 1.

DISCUSSION

Much of the oil and gas in the Illinois, Michigan, and Appalachian basins of eastern North America is thought to be derived from Devonian shale that is within these basins (for example, Miller and others, 2003; Swzezy, 2002, 2008, 2009; Swzezy and others, 2005, 2010). As the Devonian strata were buried by younger sediments, the Devonian shale was subjected to great temperature and pressure, and in some areas the shale crossed a thermal maturity threshold and began to generate oil. With increasing burial (increasing temperature and pressure), some of this oil-generating shale crossed another thermal maturity threshold and began to generate natural gas. Knowledge of the thermal maturity of the Devonian shale is therefore useful for predicting the occurrence and the spatial distribution of oil and gas within these basins.

This publication presents a thermal maturity map of Devonian shale in the Illinois, Michigan, and Appalachian basins (fig. 1). The map shows outlines of the three basins (dashed black lines) and an outline of Devonian shale (solid black lines). The basin outlines are compiled from Thomas and others (1989) and Swzezy (2008, 2009). The outline of Devonian shale is a compilation from Freeman (1978), Thomas and others (1989), de Witt and others (1993), Dart (1995), Nicholson and others (2004), Dicken and others (2005a,b), and Stoesser and others (2005).

Thermal maturity of the Devonian shale is depicted in figure 1 as based on measurements of vitrinite reflectance reported as mean percent reflectance in oil (%R), conodont color alteration, and sporopollen coloration. Maturity values are depicted using the following four colors: (1) yellow, %R = 0.6 to 0.8 (thermally immature for oil generation); (2) green, %R = 0.6 to 1.3 (thermal maturity for oil generation); (3) red, %R = 1.3 to 2.0 (thermal maturity for gas generation); and (4) gray, %R = > 2.0 (thermal maturity beyond gas generation). The information depicted in figure 1, however, is only a general approximation of the thermal maturity because of uncertainty in interpretation of thermal maturity indices and uncertainty in correlation among different thermal maturity indices. There is also uncertainty associated with the thresholds for hydrocarbon generation, which vary with kerogen type (Dow, 1977; Marchion, 1982). Despite these uncertainties, the thermal maturity interval for oil generation is defined in this study (figs. 1 and 2) as vitrinite reflectance values of 0.6 to 1.3 %R, as based on the work of Stach and others (1975), Dow (1977), Harris and others (1978), Fischer and others (1980), Cole and others (1987), Horsfield and Rullkötter (1994), Peters and Cassa (1994), Hunt (1996), Tobin and Claxton (2000), Peters and others (2005a,b), and Repetski and others (2008).

Vitrinite reflectance (VR) is the most widely used technique to estimate thermal maturity of organic matter in post-Silurian rocks (Dow, 1977; Teichmüller, 1987; Tissot and others, 1978). This technique is based on measurements of the reflective properties of terrestrial organic matter and values are reported as mean percent reflectance in oil. Conventional VR measurements are conducted on randomly oriented phyllosilicates in nonpolarized light under oil immersion. The most reliable VR measurements are obtained from the vitrinite maceral collustrite (Dow, 1977; Baskin, 1983). This maceral can be identified readily in rocks such as coal that contain relatively high concentrations of organic matter, but the maceral may be difficult to identify in rocks such as shale that contain relatively low concentrations of dispersed organic matter (Bostick and Alpern, 1977; Barker, 1996). Other characteristics of rocks that may contribute to difficulties in measurement of VR include the complexity of organic matter, the compositional variety of organic matter, and the presence of dispersed organic matter mixed with indigenous vitrinite (Bostick, 1979; Dembicki, 1984; Utting and others, 2004). Furthermore, in the use of conventional VR techniques, some types of organic matter (for example, perhydrous vitrinite) tend to yield suppressed VR values, whereas other types of organic matter (for example, subhydrous vitrinite) tend to yield enhanced VR values (McTavish, 1978; Newman and Newman, 1982; Baskin, 2000, 1983; Price and Barker, 1985; Wenger and Baker, 1987; Marchion and others, 1991; Fang and Jansen, 1992; Lo, 1993; Carr, 2000a,b). Vitrinite suppression may be particularly pronounced in alginite-rich shales (Hutton and Cook, 1989; Kalkreuth, 1982; Price and Barker, 1985; Peters and others, 2006). Several explanations have been suggested for causes of suppressed and enhanced vitrinite values, and multiple causes are likely (for example, Price and Barker, 1985; Raymond and Marchion, 1991; Carr, 2000a,b; Wilkins and others, 2002; Fedor and Hämmer-Vidó, 2003; Quick and Tabet, 2003).

The conodont color alteration index (CAI) is an index of thermal maturity that is often used for marine rocks of Ordovician to Triassic age (Epstein and others, 1977; Harris and others, 1978; Harris, 1979; Rejzhan and others, 1987). Conodonts are phosphatic marine microfossils that are thought to be the teeth elements of an eel-like marine vertebrate (Briggs and others, 1983; Briggs, 1992; Gabbot and others, 1995). These teeth elements (conodonts) contain trace amounts of organic matter that changes color with increasing thermal maturity. Charts showing the correlation of conodont CAI data with other thermal maturity indices have been published by Harris (1979) and Repetski and others (2008).

Spores and pollen exhibit changes in color and opacity with increasing temperature and depth of burial (Grayson, 1975; Lerche and McKenna, 1991). This phenomenon has been quantified using a number of indices, including the Etai de Conservation (EC) index of Correa (1967), the organic matter alteration index (TAI) of Staplin (1969), the spore color index (SCI) of Fisher and others (1980), the palynomorph color index of Batten (1980), and the thermal alteration scale (TAS) of Batten (1982). Several publications include charts that show comparisons among the various spore and pollen color indices (for example, Strapoc, 1983) and comparisons of spore and pollen color indices with other thermal maturity indices such as vitrinite reflectance and conodont CAI data (Fisher and others, 1980; Batten, 1982; Utting and others, 2004; Marshall, 1991).

Stratigraphic nomenclature for Devonian shale in the three basins is shown in figure 3. In the Illinois basin, the thermal maturity values are based on VR measurements from the basal interval of the Middle Devonian to Lower Mississippian New Albany Shale (Barrows and Cluff, 1984; Nuccio and Hatch, 1996; Strapoc and others, 2010). In the Michigan basin, the thermal maturity values are based on VR measurements and SCI measurements from the basal interval of the Upper Devonian Antrim Shale (Moyer, 1982; Cercone, 1984; Cercone and Pollack, 1991; Wang and others, 2010). In the Appalachian basin, the thermal maturity values are based on vitrinite reflectance measurements from Devonian shale and conodont CAI measurements from Devonian limestone that is stratigraphically near the Middle Devonian Marcellus Shale and correlative units (Harris and others, 1978; Harris, 1979; Bayer, 1982; Repetski and others, 2008). These values from Middle Devonian strata in the Appalachian basin are generally consistent with VR values from the Upper Devonian West Falls Formation (Gerlach and Cercone, 1993; Curtis and Faure, 1997) and from the Upper Devonian Ohio Shale (Rimmer and others, 1993; Curtis and Faure, 1997).

Although there is uncertainty in the thermal maturity data presented in figure 1 (and the interpretations from these data), some general patterns may be discerned. In the Illinois basin, the Devonian shale is within the oil window, and the thermal maturity in the central and southern portions of the basin, and the shale is immature elsewhere within the basin (Barrows and Cluff, 1984). This pattern suggests that oil in Devonian or younger strata along the margins of the Illinois basin may have migrated from the central or southern portions of the basin, and natural gas within Devonian or younger strata is likely to be of biogenic (rather than thermogenic) origin because there is very little Devonian shale that is within the gas generation window. In support of these conclusions, various geochemical studies have documented the presence of biogenic gas in the Devonian shale of the Illinois basin (McIntosh and Martini, 2008; Martini and others, 2008). In the Michigan basin, the Devonian shale is within the oil window in the central part of the basin, and the shale is immature elsewhere in the basin. This pattern suggests that oil in Devonian or younger strata along the margins of the Michigan basin may have migrated from the central part of

the basin, and natural gas within Devonian or younger strata is likely to be of biogenic (rather than thermogenic) origin because there is no Devonian shale that is within the gas generation window. In support of these conclusions, various geochemical studies have documented the presence of biogenic gas in the Devonian shale of the Michigan basin (Martini and others, 1996, 1998, 2003, 2008; McIntosh and others, 2004, 2011). In the Appalachian basin, the Devonian shale is immature on the western margin of the basin and overmature on the eastern margin of the basin. This observation suggests that oil in Devonian or younger strata on the western margin of the basin (west of the area where Devonian shale is in the oil window) is likely to have migrated from the east. Furthermore, in Devonian and younger strata, oil should be more common in the western portion of the basin (where the Devonian shale is in the oil window), and thermogenic gas should be more common in the eastern portion of the basin (where the Devonian shale is in the gas window). Also, gas-prone kerogen is more common in the eastern portion of the basin, and oil-prone kerogen is more common in the western portion of the basin (Zielinski and Melver, 1982). Thus, any natural gas within Devonian or younger strata on the western margin of the basin is likely to have migrated from the eastern part of the basin or to be of biogenic (rather than thermogenic) origin because of the low thermal maturity on the western margin of the basin. In support of these conclusions, various geochemical studies have documented some small accumulations of biogenic gas in the Devonian shale of the northern Appalachian basin, along the southern shore of Lake Erie and in western New York (Osborn and McIntosh, 2010).

REFERENCES

Barker, C.E., 1996, A comparison of vitrinite reflectance measurements made on whole-rock and dispersed organic matter concentrate mounts: *Organic Geochemistry*, v. 24, no. 2, p. 251-256.

Barrows, M.H., and Cluff, R.M., 1984, New Albany Shale Group (Devonian-Mississippian) source rocks and hydrocarbon generation: Illinois basin, in: Demas, Gerard, and Morris, R.J., eds., *Petroleum geochemistry and basin evaluation: American Association of Petroleum Geologists Memoir* 35, p. 111-138.

Batten, D.J., 1980, Use of transmitted light microscopy of sedimentary organic matter for evaluation of hydrocarbon source potential: Fourth International Palynological Conference, Lucknow, India, Proceedings, v. 2, p. 589-594.

Batten, D.J., 1982, Palynofacies, palaeoenvironment and petroleum: *Journal of Micropalaeontology*, v. 1, no. 1, p. 107-114.

Bayer, K.C., 1982, Map showing approximate eastern limit of commercial oil and gas fields in relation to structural features and physiographic provinces in the Appalachian region: U.S. Geological Survey Oil and Gas Investigations Chart OC-121, 1 sheet.

Bostick, N.H., 1979, Microscopic measurement of the level of catagenesis of solid organic matter in sedimentary rocks to aid exploration for petroleum and to determine former burial temperatures—A review, in: Scholle, P.A., and Schlager, P.R., eds., *Aspects of diagnosis: Society for Sedimentary Geology (SEPM) Special Publication* 26, p. 17-43.

Bostick, N.H., and Alpern, B., 1977, Principles of sampling, preparation and constituent selection for microphotometry in measurement of maturation of sedimentary organic matter: *Journal of Microscopy*, v. 109, no. 1, p. 41-47.

Briggs, D.E.G., 1992, Conodonts: A major extinct group added to the vertebrates: *Science*, v. 256, p. 1285-1286.

Briggs, D.E.G., Clarkson, E.N.K., and Aldridge, R.J., 1983, The conodont animal: *Lethaia*, v. 16, no. 1, p. 1-14.

Baskin, J.M., 1983, Selection criteria for the use of vitrinite reflectance as a maturity tool, in: Brooks, J., ed., *Petroleum geochemistry and exploration in Europe*: Oxford, Blackwell Scientific Publications, The Geological Society (London) Special Publication 12, p. 295-307.

Carr, A.D., 2000a, Suppression and retardation of vitrinite reflectance, part 1: Formation and significance for hydrocarbon generation: *Journal of Petroleum Geology*, v. 23, no. 3, p. 313-343.

Carr, A.D., 2000b, Suppression and retardation of vitrinite reflectance, part 2: Derivation and testing of a kinetic model for suppression: *Journal of Petroleum Geology*, v. 23, no. 4, p. 475-496.

Cercone, K.R., 1984, Thermal history of Michigan basin: American Association of Petroleum Geologists Bulletin, v. 68, no. 2, p. 130-136.

Cercone, K.R., and Pollack, H.N., 1991, Thermal maturity of the Michigan basin, in: Catocinos, P.A., and Daniels, P.A., eds., *Early sedimentary evolution of the Michigan basin: Geological Society of America Special Paper* 256, p. 1-11.

Cole, G.A., Drozd, R.J., Sedivy, R.A., and Halpern, H.I., 1987, Organic geochemistry and oil-source correlations, Paleozoic of Ohio: American Association of Petroleum Geologists Bulletin, v. 71, no. 7, p. 788-809.

Correa, Michel, 1967, Relations possible entre l'état de conservation des éléments figures de la matière organique (microfossiles palynologiques) et l'existence de gisements d'hydrocarbures: *Revue de l'Institut Français du Pétrole*, v. 22, no. 9, p. 1285-1306.

Curtis, J.B., and Faure, G., 1997, Accumulation of organic matter in the Rome trough of the Appalachian basin and its subsequent thermal history: American Association of Petroleum Geologists Bulletin, v. 81, no. 3, p. 424-437.

Dart, R.L., 1995, Maps of upper Mississippian embayment Paleozoic and Precambrian rocks: U.S. Geological Survey Miscellaneous Field Studies Map MF-2284, 1 sheet.

de Witt, Wallace, Jr., Roen, J.B., and Wallace, L.G., 1993, Stratigraphy of Devonian black shales and associated rocks in the Appalachian basin, in: Roen, J.B., and Kerpelke, R.C., *Petroleum geology of the Devonian and Mississippian black shale of eastern North America*: U.S. Geological Survey Bulletin 1909, p. 81-847.

Dembicki, Harry, Jr., 1984, An interlaboratory comparison of source rock data: *Geochimica et Cosmochimica Acta*, v. 48, no. 12, p. 2641-2649.

Devera, J.A., and Hasenmueller, N.R., 1990, Kaskaskia Sequence—Middle and Upper Devonian Series through Mississippian Kinderhookian Series, in: Leighton, M.W., Kolata, D.R., O'Leary, D.F., and Eidel, J.J., eds., *Interior carbonate basins: American Association of Petroleum Geologists Memoir* 51, p. 113-123.

Dicken, C.L., Nicholson, S.W., Horton, J.D., Foose, M.P., and Mueller, J.A.L., 2005a, Preliminary integrated geologic map databases for the United States: Alabama, Florida, Georgia, Mississippi, North Carolina, and South Carolina (version 1.1, updated December 2007): U.S. Geological Survey Open-File Report 2005-1323, available online at <http://pubs.usgs.gov/of/2005/1323/>.

Dicken, C.L., Nicholson, S.W., Horton, J.D., Kinney, S.A., Gunther, Gregory, Foose, M.P., and Mueller, J.A.L., 2005b, Preliminary integrated geologic map databases for the United States: Delaware, Maryland, New York, Pennsylvania, and Virginia (version 1.1, updated August 2008): U.S. Geological Survey Open-File Report 2005-1325, available online at <http://pubs.usgs.gov/of/2005/1325/>.

Dow, W.G., 1977, Kerogen studies and geological interpretations: *Journal of Geochemical Exploration*, v. 7, p. 79-99.

Epstein, A.G., Epstein, J.B., and Harris, L.D., 1977, Conodont color alteration—An index to organic metamorphism: U.S. Geological Survey Professional Paper 995, 27 p.

Everham, W.D., 2004, Thermal history of the Michigan basin: Ph.D. dissertation, Michigan Technological University, Houghton, Michigan, 251 p.

Fang, H., and Jianyu, C., 1992, The cause and mechanism of vitrinite reflectance anomalies: *Journal of Petroleum Geology*, v. 15, no. 4, p. 419-434.

Fedor, F., and Hämmer-Vidó, M., 2003, Statistical analysis of vitrinite reflectance data—A new approach: *International Journal of Coal Geology*, v. 56, no. 3-4, p. 277-294.

Fisher, M.J., Barnard, P.C., and Cooper, B.S., 1980, Organic maturation and hydrocarbon generation in the Mesozoic sediments of the Sverdrup basin, Arctic Canada: Fourth International Palynological Conference, Lucknow, India, Proceedings, v. 2, p. 581-588.

Freeman, E.B., ed., 1978, Geological highway map, southern Ontario: Ontario Geological Survey Map 2418, 1 sheet.

Gabbott, S.E., Aldridge, R.J., and Theron, J.N., 1995, A giant conodont with preserved muscle tissue from the Upper Ordovician of South Africa: *Nature*, v. 374, p. 800-801.

Gerlach, J.B., and Cercone, K.R., 1993, Former Carboniferous overburden in the northern Appalachian basin: A reconstruction based on vitrinite reflectance: *Organic Geochemistry*, v. 20, no. 2, p. 223-232.

Gradstein, F., Ogg, J.G., and Smith, A.G., eds., 2004, *A geological time scale 2004*: Cambridge University Press, Cambridge, UK, 589 p.

Grayson, J.F., 1975, Pétrographie de la matière organique des sédiments, relations avec la paléotempérature et le potentiel pétrolier [Relations of palynomorph transluency to carbon and hydrocarbon in clastic sediments]: Centre National de la Recherche Scientifique, Paris, France, p. 261-273.

Harris, A.G., 1979, Conodont color alteration, an organo-mineral metamorphic index, and its application to Appalachian basin geology, in: Scholle, P.A., and Schlager, P.R., eds., *Aspects of diagnosis: Society for Sedimentary Geology (SEPM) Special Publication* 26, p. 3-16.

Harris, A.G., Harris, L.D., and Epstein, J.B., 1978, Oil and gas data from Paleozoic rocks in the Appalachian basin: Maps for assessing hydrocarbon potential and thermal maturity (conodont color alteration isograds and overburden isograds): U.S. Geological Survey Miscellaneous Investigations Series Map I-917-E, 4 sheets, scale 1:250,000.

Hasenmueller, N.R., 1993, New Albany Shale (Devonian and Mississippian) of the Illinois basin, in: Roen, J.B., and Kerpelke, R.C., eds., *Petroleum geology of the Devonian and Mississippian black shale of eastern North America*: U.S. Geological Survey Bulletin 1909, p. C1-C19.

Hayes, D.O., 2005, Thermal evolution of the Michigan basin, in: *Petroleum Technology Transfer Council (PTTC), Part II—Undiscovered oil and gas in the Michigan basin*: Petroleum Technology Transfer Council (PTTC) Workshop, Mount Pleasant, Michigan, October 20, 2005, p. 195-229.

Horsfield, B., and Rullkötter, J., 1994, Diagenesis, catagenesis, and metagenesis of organic matter, in: Magoon, L.B., and Dow, W.G., eds., *The petroleum system—A review to trap*: American Association of Petroleum Geologists Memoir 60, p. 189-199.

Hunt, J.M., 1996, *Petroleum geochemistry and geology* (2d ed.): New York, W.H. Freeman and Company, 743 p.

Hutton, A.C., and Cook, A.G., 1980, Influence of angle on the reflectance of vitrinite from Joadja, NSW, and some other coals and oil shales containing alginite: *Fuel*, v. 59, no. 10, p. 711-714.

Kalkreuth, W.D., 1982, Rank and petrographic composition of selected Jurassic—Lower Cretaceous coals of British Columbia, Canada: *Bulletin of Canadian Petroleum Geology*, v. 30, no. 2, p. 112-139.

Lerche, I., and McKenna, T.E., 1991, Pollen translucency as a thermal maturation indicator: *Journal of Metamorphic Geology*, v. 9, no. 1, p. 19-36.

Lo, H.B., 1993, Correction criteria for the suppression of vitrinite reflectance in hydrogen-rich kerogens: Preliminary guidelines: *Organic Geochemistry*, v. 20, no. 6, p. 653-657.

Marshall, L.E., 1991, Quantitative spore colour: *Journal of the Geological Society (London)*, v. 148, p. 223-233.

Martini, A.M., Budai, J.M., Walter, L.M., and Schoell, Martin, 1996, Microbial generation of economic accumulations of methane within a shallow organic-rich shale: *Nature*, v. 383, p. 155-158.

Martini, A.M., Walter, L.M., Budai, J.M., Ku, T.C.W., Kaiser, C.J., and Schoell, Martin, 1998, Genetic and temporal relations between formation waters and biogenic methane: Upper Devonian Antrim Shale, Michigan basin, USA—CO₂ reduction vs. acetate fermentation— isotopic evidence: *Geochimica et Cosmochimica Acta*, v. 62, no. 10, p. 1699-1720.

Martini, A.M., Walter, L.M., Ku, T.C.W., Budai, J.M., McIntosh, J.C., and Schoell, Martin, 2003, Microbial production and modification of gases in sedimentary basins: A geochemical case study from a Devonian shale play, Michigan basin: American Association of Petroleum Geologists Bulletin, v. 87, no. 8, p. 1355-1375.

Martini, A.M., Walter, L.M., and McIntosh, J.C., 2008, Identification of microbial and thermogenic gas components from Upper Devonian black shale cores, Illinois and Michigan basins: American Association of Petroleum Geologists Bulletin, v. 92, no. 3, p. 327-339.

Matthews, R.D., 1993, Review and revision of the Devonian-Mississippian stratigraphy in the Michigan basin, in: Roen, J.B., and Kerpelke, R.C., eds., *Petroleum geology of the Devonian and Mississippian black shale of eastern North America*: U.S. Geological Survey Bulletin 1909, p. D1-D85.

McIntosh, J.C., Garven, G., and Hanor, J.S., 2011, Impacts of Pleistocene glaciations on large-scale groundwater flow and salinity in the Michigan basin: *Geofluids*, v. 11, no. 1, p. 18-33.

McIntosh, J.C., and Martini, A.M., 2008, Hydrogeological indicators for microbial methane in fractured black shales: Case studies of the Antrim, New Albany, and Ohio Shales, in: Hill, D.G., Lillis, P.G., and Curtis, J.B., eds., *Gas shale in the Rocky Mountains and beyond: Rocky Mountain Association of Geologists 2008 Guidebook*, p. 162-174.

McIntosh, J.C., Walter, L.M., and Martini, A.M., 2004, Extensive microbial modification of formation water geochemistry: Case study from a midcontinent sedimentary basin, United States: *Geological Society of America Bulletin*, v. 116, no. 5-6, p. 743-759.

McTavish, R.A., 1978, Pressure retardation of vitrinite diagenesis, offshore north-west Europe: *Nature*, v. 271, p. 648-650.

Milici, R.C., Ryder, R.T., Swzezy, C.S., Charpentier, R.R., Cook, T.A., Crovelli, R.A., Klett, T.R., Pollastro, R.M., and Schenk, C.J., 2003, Assessment of undiscovered oil and gas resources of the Appalachian Basin Province, 2002: U.S. Geological Survey Fact Sheet 2003-03, 2 p.

Moyer, R.B., 1982, Thermal maturity and organic content of selected Paleozoic formations—Michigan basin: Master's thesis, Michigan State University, East Lansing, 62 p.

Marchion, D.G., 1987, Recent advances in organic petrology and organic geochemistry: An overview with some reference to 'oil' from coal: in: Scott, A.C., ed., *Coal and coal-bearing strata: Recent advances: Geological Society (London) Special Publication* 22, p. 257-302.

Marchion, D.G., Pearson, J., and Raymond, A.C., 1991, Anomalies in vitrinite reflectance gradients: Bulletin de la Société Géologique de France, v. 162, no. 2, p. 183-191.

Newman, J., and Newman, N.A., 1982, Reflectance anomalies in Pike River coals: Evidence of variation in vitrinite type, with implications for maturation studies and "gagete" rank: *New Zealand Journal of Geology and Geophysics*, v. 25, no. 2, p. 233-243.

Nicholson, S.W., Dicken, C.L., Foose, M.P., and Mueller, J.A.L., 2004, Preliminary integrated geologic map databases for the United States: Minnesota, Wisconsin, Michigan, Illinois, and Indiana (version 1.1, updated December 2007): U.S. Geological Survey Open-File Report 2004-1355, available at <http://pubs.usgs.gov/of/2004/1355/>.

Nuccio, R.C., and Hatch, R.J., 1996, Vitrinite reflectance suppression in the New Albany Shale, Illinois basin—Vitrinite reflectance and Rock-Eval data: U.S. Geological Survey Open-File Report 96-665, 37 p.

Osborn, S.G., and McIntosh, J.C., 2010, Chemical and isotope tracers of the contribution of microbial gas in Devonian organic-rich shales and reservoir sandstones, northern Appalachian basin: *Applied Geochemistry*, v. 25, no. 3, p. 456-471.

Peters, K.E., and Cassa, M.R., 1994, Applied source rock geochemistry, in: Magoon, L.B., and Dow, W.G., eds., *The petroleum system—A review to trap*: American Association of Petroleum Geologists Memoir 60, p. 93-120.

Peters, K.E., Walters, C.C., and Moldrow, J.M., 2005a, The biomarker guide, volume I: Biomarkers and isotopes in petroleum exploration and earth history: Cambridge University Press, Cambridge, UK, p. 475-1155.

Peters, K.E., Walters, C.C., and Moldrow, J.M., 2005b, The biomarker guide, volume I: Biomarkers and isotopes in petroleum exploration and earth history: Cambridge University Press, Cambridge, UK, 471 p.

Scientific Investigations Map 3214

Petersen, H.I., 2002, A re-consideration of the "oil window" for hamic coal and kerogen type III source rocks: *Journal of Petroleum Geology*, v. 25, no. 4, p. 407-432.

Petersen, H.I., Foopattanakamol, A., and Ratnasathian, B., 2006, Petroleum potential, thermal maturity and the oil window of oil shales and coals in Cenozoic rift basins, central and northern Thailand: *Journal of Petroleum Geology*, v. 29, no. 4, p. 337-360.

Price, L.C., and Barker, C.E., 1985, Suppression of vitrinite reflectance in amorphous rich kerogen—A major unrecognized problem: *Journal of Petroleum Geology*, v. 8, no. 1, p. 59-84.

Quiver, J.C., and Tabet, D.E., 2003, Suppressed vitrinite reflectance in the Ferron coalbed gas fairway, central Utah: Possible influence of overpressure: *International Journal of Coal Geology*, v. 56, no. 1-2, p. 49-67.

Raymond, A.C., and Marchion, D.G., 1991, The relationship between organic maturation, the widths of thermal aureoles and the thicknesses of sills in the Midland Valley of Scotland and northern England: *Journal of the Geological Society (London)*, v. 148, no. 2, p. 215-218.

Rejzhan, V.A., Harris, A.G., and Huebert, J.S., 1987, Conodont color and textural alteration: An index to regional metamorphism, contact metamorphism, and hydrothermal alteration: *Geological Society of America Bulletin*, v. 99, no. 4, p. 471-479.

Repetski, J.E., Ryder, R.T., Woary, D.J., Harris, A.G., and Trippi, M.H., 2008, Thermal maturity patterns (CAI and %R) in Upper Ordovician and Devonian rocks of the Appalachian Basin: A major revision of USGS Map I-917-E using new subsurface collections: U.S. Geological Survey Scientific Investigations Map SIM 3006, 1 CD-ROM. (Also available at <http://pubs.usgs.gov/sim/3006/>.)

Rimmer, S.M., Canell, D.J., and Gooding, P.J., 1993, Rock-Eval pyrolysis and vitrinite reflectance trends in the Cleveland Shale Member of the Ohio Shale, eastern Kentucky: *Organic Geochemistry*, v. 20, no. 6, p. 735-745.

Ryder, R.T., Crangle, R.D., Jr., Trippi, M.H., Swzezy, C.S., Lentz, E.E., Rowan, E.L., and Hope, R.S., 2009, Geologic cross section D-D' through the central Appalachian basin from the Findlay arch, Sandusky County, Ohio, to the Valley and Ridge province, Hardy County, West Virginia: U.S. Geological Survey Scientific Investigations Map SIM 3007, 2 sheets with 52-page pamphlet, available at <http://pubs.usgs.gov/sim/>.

Shaver, R.H., Ault, C.H., Burger, A.M., Carr, D.J., Droste, J.B., Eggert, D.L., Gray, H.H., Harper, D., Hasenmueller, N.R., Hasenmueller, W.A., Horowitz, A.S., Hutchinson, H.C., Keith, B.D., Keller, S.J., Patton, J.B., Rexroad, C.B., and Wier, C.E., 1986, Compendium of Paleozoic rock-unit stratigraphy in Indiana: A revision: *Indiana Geological Survey Bulletin* 59, 203 p.

Smith, P.M.R., 1983, Spectral correlation of spore coloration standards, in: Brooks, J., ed., *Petroleum geochemistry and exploration in Europe: The Geological Society (London) Special Publication* 12, p. 289-294.

Stach, E., Mackowsky, M.-Th., Teichmüller, M., Taylor, G.H., Chandra, D., and Teichmüller, R., 1975, Stach's textbook of coal petrology (2d ed.): Gebrüder Bornier, Berlin, 428 p.

Staplin, F.J., 1969, Sedimentary organic matter, organic metamorphism, and oil and gas occurrence: *Bulletin of Canadian Petroleum Geology*, v. 17, no. 1, p. 47-66.

Stoesser, D.B., Green, G.N., Morath, L.C., Heran, W.D., Wilson, A.B., Moore, D.W., and Van Gosen, B.S., 2005, Preliminary integrated geologic map databases for the United States: Montana, Wyoming, Colorado, New Mexico, North Dakota, South Dakota, Nebraska, Kansas, Oklahoma, Texas, Iowa, Missouri, Arkansas, and Louisiana (version 1.2, updated December 2007): U.S. Geological Survey Open-File Report 2005-1351, available at <http://pubs.usgs.gov/of/2005/1351/>.

Strapoc, D., Mastalerz, M., Schimmelmann, A., Drobnik, A., and Hasenmueller, N.R., 2010, Geochemical constraints on the origin and volume of gas in the New Albany Shale (Devonian-Mississippian), eastern Illinois basin: American Association of Petroleum Geologists Bulletin, v. 94, no. 11, p. 1713-1740.

Swzezy, C.S., 2002, Regional stratigraphy and petroleum systems of the Appalachian basin, North America: U.S. Geological Survey Geological Investigations Series Map I-2768, 1 sheet, available at <http://pubs.usgs.gov/vmap/i-2768/>.

Swzezy, C.S., 2008, Regional stratigraphy and petroleum systems of the Michigan basin, North America: U.S. Geological Survey Scientific Investigations Map 2978, 1 sheet, available at <http://pubs.usgs.gov/vmap/2978/>.

Swzezy, C.S., 2009, Regional stratigraphy and petroleum systems of the Illinois basin, U.S.A.: U.S. Geological Survey Scientific Investigations Map 3068, 1 sheet, available at <http://pubs.usgs.gov/vmap/3068/>.

Swzezy, C.S., Hatch, J.R., Brennan, S.T., East, J.A., Rowan, E.L., Repetski, J.E., Charpentier, R.R., Cook, T.A., Klett, T.R., Pollastro, R.M., and Schenk, C.J., 2007, Assessment of undiscovered oil and gas resources of the Illinois basin, 2007: U.S. Geological Survey Fact Sheet 2007-3058, available at <http://pubs.usgs.gov/fact/2007/3058/>.

Swzezy, C.S., Hatch, J.R., East, J.A., Hayba, D.O., and Repetski, J.E., in press, Petroleum systems and assessment of the U.S. portion of the Michigan basin: U.S. Geological Survey Digital Data Series DDS-69-AA, 1 CD-ROM.

Swzezy, C.S., Hatch, J.R., Hayba, D.O., Repetski, J.E., Charpentier, R.R., Cook, T.A., Klett, T.R., Pollastro, R.M., and Schenk, C.J., 2005, Assessment of undiscovered oil and gas resources of the U.S. portion of the Michigan basin, 2004: U.S. Geological Survey Fact Sheet 2005-3070, available at <http://pubs.usgs.gov/fact/2005/3070/>.

Teichmüller, M., 1987, Recent advances in coalification studies and their application to geology, in: Scott, A.C., ed., *Coal and coal-bearing strata: Recent advances: Geological Society (London) Special Publication* 32, p. 127-169.

Thomas, W.A., Chown, T.M., Daniels, D.L., Neathery, T.L., Glover, L. III, and Gleason, R.J., 1989, The subsurface Appalachians beneath the Atlantic and Gulf Coastal Plains, in: Hatch, R.D., Jr., Thomas, W.A., and Vile, G.W., eds., *The Appalachian-Ouachita region in the United States: The geology of North America*, v. F-2, Geological Society of America, Denver, p. 445-458.

Tissot, B.P., Pelet, R., and Ungerer, P.H., 1987, Thermal history of sedimentary basins, maturation indices, and kinetics of oil and gas generation: American Association of Petroleum Geologists Bulletin, v. 71, no. 12, p. 1445-1466.

Tobin, R.C., and Claxton, B.L., 2000, Multidisciplinary thermal maturity studies using vitrinite reflectance and fluid inclusion microthermometry: A new calibration of old techniques: American Association of Petroleum Geologists Bulletin, v. 84, no. 10, p. 1647-1665.

Utting, J., Goodarzi, F., Dougherty, B.J., and Henderson, C.M., 1989, Thermal maturity of Carboniferous and Permian rocks of the Sverdrup basin, Canadian Arctic Archipelago: *Geological Survey of Canada Paper* 89-19, 29 p.

Utting, John Spina, Amalia, Jamieson, Ian, McGregor, D.C., and Marshall, J.E., 2004, Revoked miopores in the upper Paleozoic and Lower Triassic of the northern circum-polar area and selected localities: *Palynology*, v. 28, no. 1, p. 75-119.

Wang, H.F., Crowley, K.D., and Nadel, G.C., 1994, Thermal history of the Michigan basin from apatite fission-track analysis and vitrinite reflectance, in: Ortoleva, P.J., ed., *Basin compartments and seals: American Association of Petroleum Geologists Memoir* 61, p. 167-177.

Wenger, L.M., and Baker, D.R., 1987, Variations in vitrinite reflectance with organic facies—Examples from Pennsylvania cyclocoenoses of the Midcontinent, U.S.A.: *Organic Geochemistry*, v. 11, no. 5, p. 411-416.

Wilkins, R.W.T., Dessel, C.F., and Bucknigham, C.P., 2002, Comparison of two petrographic methods for determining the degree of anomalous vitrinite reflectance: *International Journal of Coal Geology*, v. 52, no. 1-4, p. 45-62.

William, H.B., Atherton, E., Buchhorn, T.C., Gies, G., Hopkins, J.C., Hopkins, M.E., Linkswage, J.A., and Simon, J.A., 1975, Handbook of Illinois stratigraphy: Illinois State Geological Survey Bulletin 95, 2