



## Local (Offshore of Ventura Map Area) and Regional (Offshore from Refugio Beach to Hueneme Canyon) Shallow-Subsurface Geology and Structure, Santa Barbara Channel, California

By  
Samuel Y. Johnson, Eleyne L. Phillips, Andrew C. Ritchie, Florence L. Wong, Ray W. Sliter, Amy E. Draut, and Patrick E. Hart  
2013

**DISCUSSION**

This sheet includes maps that show the thickness and the depth to base of uppermost Pleistocene and Holocene (in other words, post-Last Glacial Maximum) deposits for the Offshore of Ventura map area (Maps A, B), as well as for a larger area that extends about 115 km along the coast from Hueneme Canyon to Refugio Beach (Maps D, E) to establish a regional context. To make these maps, water bottom and depth to base of the post-Last Glacial Maximum horizons were mapped from seismic-reflection profiles. The difference in the two horizons was reported for every shot point as XY coordinates (UTM zone 11) and two-way travel time (TWT). The thickness of the post-Last Glacial Maximum unit (Maps B, E) was determined by applying a sound velocity of 1,600 m/s to the TWT. The thickness points were interpolated to a preliminary continuous surface, overlaid with zero-thickness bedrock outcrops (sheet 10 of this report), and contoured (Wong and others, 2012). Data within Hueneme Canyon were excluded from the contouring because the seismic-reflection data are too sparse to adequately image the highly variable changes in sediment thickness that characterize the canyon (Maps A, B, D, E).

Several factors required manual editing of the preliminary thickness maps to make the final products. The Red Mountain Fault Zone (RMFZ), Pitas Point Fault (PPF), and Oak Ridge Fault (ORF) along the sediment sequence in the region (Maps D, E). The data points also are dense along tracklines (about 1 m apart) and sparse between tracklines (1–2 km apart), resulting in contouring artifacts. To incorporate the geologic information and complexity, the resulting interpolated contours were modified. Contour modifications and regrading were repeated several times to produce the final regional sediment-thickness map (Wong and others, 2012).

Data to determine the depth to base of the post-Last Glacial Maximum unit was similarly processed and contoured. However, this preliminary data set was set aside in favor of a surface determined by subtracting the modified thickness data from multi-beam bathymetry collected separately (see sheet 1). The depth of this surface in the Hueneme Canyon to Refugio Beach area ranges from 12 to 190 m (Map E; see also Wong and others, 2012).

Five different "domains" of sediment thickness, which are bounded either by faults or by Hueneme Canyon, are recognized on the regional maps (Maps D, E): (1) Refugio Beach to the southern strand of the Red Mountain Fault Zone (RMFZ); (2) between the southern strand of the Red Mountain Fault Zone and the Pitas Point Fault (PPF); (3) between the Pitas Point Fault and Oak Ridge Fault (ORF); (4) between the Oak Ridge Fault and Hueneme Canyon; and (5) south of Hueneme Canyon. These data highlight the contrast among three general zones of sediment thickness: (1) the uplifted, sediment-poor Santa Barbara shelf (domain 1; mean sediment thickness of 3.5 m); (2) a transitional zone (domain 2; mean sediment thickness of 18.0 m); and (3) the subsiding, sediment-rich delta and shelf offshore of the Ventura and Santa Clara Rivers and Calleguas Creek (domains 3, 4, and 5; mean sediment thicknesses of 39.3, 38.9, and 28.3 m, respectively).

In the Offshore of Ventura map area, sediment thickness ranges from 0 to 53 m, with a mean thickness of 34.1 m. The two major reasons for this large thickness are that (1) the map area lies offshore of the mouths of the Santa Clara and Ventura Rivers, both major sediment sources, and (2) much of the area lies within the sedimentary basin in the south flank of the active, north-dipping Pitas Point Fault (sheet 8, 10). The north flank of the Pitas Point Fault is rapidly subsiding, and a steep gradient of sediment thickness exists where crossing the fault from north (mean thickness, 17.0 m, thinning rapidly northward) to south (mean thickness, about 40 m). Thicker sediment (18 to 53 m) in the basin south of the Pitas Point Fault is cut by the active, south-

**REFERENCES CITED**

Caldwell, R.J., Taylor, L.A., Eakin, B.W., Carignan, K.S., Grobe, P.R., Lim, E., and Friday, D.Z., 2010, Digital elevation models of Santa Monica, California—Procedures, data sources and analysis: NOAA Technical Memorandum NESDIS NGDC-46, NOAA National Geophysical Data Center, available at <http://www.ngk.noaa.gov/dem/squareCalGrid/download/603>.

Carignan, K.S., Taylor, L.A., Eakin, B.W., Warrick, R.R., Lim, E., and Medley, P.R., 2009, Digital elevation model of Santa Barbara, California—Procedures, data sources, and analysis: NOAA Technical Memorandum NESDIS NGDC-29, NOAA National Geophysical Data Center, available at <http://www.ngk.noaa.gov/dem/squareCEGrid/download/603>.

Hack, R.G., 1998, Santa Barbara Channel regional foreline map, topography, formation, in Kunitomi, D.S., Hopp, T.E., and Galloway, J.M., eds., Structure and petroleum geology, Santa Barbara Channel, California: American Association of Petroleum Geologists, Pacific Section, Miscellaneous Publication 46, 1 plate.

Jennings, C.W., and Bryant, W.A., 2010, Fault activity map of California: Geological Survey Geologic Data Map no. 6, scale 1:750,000.

Lee, W.H., and Vukobrat, J.G., 1973, Recent earthquake activity in the Santa Barbara Channel region: Bulletin of the Seismological Society of America, v. 63, p. 1,757–1,773.

Mior, S.A., Kellogg, K.S., Stanley, R.G., Gormila, L.D., Keller, E.A., and Brandt, T.R., 2009, Geologic map of the Santa Barbara coastal plain area, Santa Barbara County, California: U.S. Geological Survey Scientific Investigations Map 3001, scale 1:25,000, 1 sheet, pamphlet 38 p., available at <http://pubs.usgs.gov/sim/3001/>.

Southern California Earthquake Data Center, 2010, Southern California Earthquake Catalog: Southern California Earthquake Data Center database, accessed April 5, 2011, at <http://www.data.scec.org/catalog/index.html>.

Wong, F.L., Phillips, E.L., Johnson, S.Y., and Sliter, R.W., 2012, Modeling of depth to base of Last Glacial Maximum and seafloor sediment thickness for the California State Waters Map Series, eastern Santa Barbara Channel, California: U.S. Geological Survey Open-File Report 2012-1161, 16 p., available at <http://pubs.usgs.gov/ofr/2012/1161/>.



Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

This map was printed on an electronic photo directly from digital file. Downloaded coloration may vary between electronic displays and between X and Y directions on the same plate, and paper may change color due to atmospheric conditions, therefore, color and proportions may vary from one plate to another.

For sale by U.S. Geological Survey, Information Systems, Box 25086, Federal Center, Denver, CO 80225-1888-4855

Digital file available at <http://pubs.usgs.gov/sim/3254/>

Suggested Citation: Johnson, S.Y., Phillips, E.L., Ritchie, A.C., Wong, F.L., Sliter, R.W., Draut, A.E., and Hart, P.E., 2013, Local (Offshore of Ventura map area) and regional (Offshore from Refugio Beach to Hueneme Canyon) Shallow-Subsurface Geology and Structure, Santa Barbara Channel, California, in: Hart, P.E., Johnson, S.Y., Draut, A.E., Phillips, E.L., Ritchie, A.C., Wong, F.L., Sliter, R.W., and Hart, P.E., eds., U.S. Geological Survey Scientific Investigations Map 3254, sheet 9 of 11, available at <http://pubs.usgs.gov/sim/3254/>.