

DISCUSSION

This sheet includes maps that show the thickness and the depth to base of uppermost Pleistocene and Holocene (in other words, post-Late Glacial Maximum) deposits for the Offshore of Santa Barbara map area (Maps A, B), as well as for a larger area that extends about 115 km along the coast from Hueneme Canyon to Refugio Beach (Maps D, E) to establish a regional context. To make these maps, water bottom and depth to base of the post-Late Glacial Maximum horizons were mapped from seismic-reflection profiles. The difference in the two horizons was exported for every shot point as XY coordinates (UTM zone 11) and two-way travel time (TWT). The thickness of the post-Late Glacial Maximum unit (Maps B, E) was determined by applying a sound velocity of 1,600 m/sec to the TWT. The thickness points were interpolated to a preliminary continuous surface, overlaid with zero-thickness bedrock outcrop (sheet 10 of this report), and contoured (Wong and others, 2012). Data within Hueneme Canyon were excluded from the contouring because the seismic-reflection data are too sparse to adequately image the highly variable changes in sediment thickness that characterize the canyon (Maps A, B, D, E).

Several factors required manual editing of the preliminary thickness maps to make the final products. The Red Mountain Fault Zone (RMFZ), Pitas Point Fault (PPF), and Oak Ridge Fault (ORF) during the sediment sequence in the region (Maps D, E). The data points also are along tracklines (about 1 m apart) and sparse between tracklines (1–2 km apart), resulting in contouring artifacts. To incorporate the effect of the faults, to remove irregularities from interpolation, and to reflect other geologic information and complexity, the resulting interpolated contours were modified. Contour modifications and regridding were repeated several times to produce the final regional sediment-thickness map (Wong and others, 2012).

Data to determine the depth to base of the post-Late Glacial Maximum unit was similarly processed and contoured. However, this preliminary data set was not made in favor of a surface determined by subtracting the modified thickness data from multi-beam bathymetry collected separately (see sheet 1). The depth of this surface in the Hueneme Canyon to Refugio Beach area ranges from 12 to 190 m (Map D, see also, Wong and others, 2012).

Five different "domains" of sediment thickness, which are bounded either by faults or by Hueneme Canyon, are recognized on the regional maps (Maps D, E): (1) Refugio Beach to the south strand of the Red Mountain Fault Zone (RMFZ); (2) between the south strand of the Red Mountain Fault Zone and the Pitas Point Fault (PPF); (3) between the Pitas Point Fault and the Oak Ridge Fault (ORF); (4) between the Oak Ridge Fault and Hueneme Canyon; and (5) south of Hueneme Canyon. These data highlight the contrast among three general zones of sediment thickness: (1) the uplifted, sediment-poor Santa Barbara shelf (domain 1, mean sediment thickness of 3.5 m); (2) a transitional zone (domain 2, mean sediment thickness of 18.0 m); and (3) the subsiding, sediment-rich delta and shelf offshore of the Ventura and Santa Clara Rivers and Calleguas Creek (domains 3, 4, and 5; mean sediment thickness of 39.3, 38.9, and 28.3 m, respectively).

In the Offshore of Santa Barbara map area, sediment thickness ranges from 0 to 18 m, with a mean thickness of 3.4 m. The thickest sediment accumulations are in three nearshore to inner-shelf deposystems, which have positive seaward relief characterized by relatively flat upper surfaces and steeper seaward-dipping slopes. These features are correlated with step-up and coastal water-bottoms, and thus, are interpreted as delta-mouth bars. Sediment is notably lacking or forms only a thin veneer over bedrock outcrops in much of the midshelf to outer shelf area.

The regional pattern of faults and of earthquakes occurring between 1922 and

REFERENCES CITED

Caldwell, R.J., Taylor, L.A., Fakus, B.W., Carignan, K.S., Grobe, P.R., Lim, E., and Friday, D.Z., 2010, Digital elevation models of Santa Monica, California—Procedures, data sources, and analysis: NOAA Technical Memorandum NESDIS NGDC-46, NOAA National Geophysical Data Center, available at <http://www.ngdc.noaa.gov/dem/square/cellGrid/download/663>.

Carignan, K.S., Taylor, L.A., Fakus, B.W., Wamken, R.R., Lim, E., and Medley, P.R., 2009, Digital elevation model of Santa Barbara, California—Procedures, data sources, and analysis: NOAA Technical Memorandum NESDIS NGDC-29, NOAA National Geophysical Data Center, available at <http://www.ngdc.noaa.gov/dem/square/cellGrid/download/663>.

Heck, R.G., 1998, Santa Barbara Channel regional foreline map, top Monterey Formation, in Kuntze, D.S., Higgs, J.E., and Galloway, J.M., eds., Structure and petroleum geology, Santa Barbara Channel, California—American Association of Petroleum Geologists, Pacific Section, and Coast Geological Society, Miscellaneous Publication 46, 1 plate.


Jennings, C.W., and Bryant, W.A., 2010, Fault activity map of California: California Geological Survey Geologic Data Map no. 6, scale 1:750,000.

Lee, W.H.K., and Vedder, J.G., 1973, Recent earthquake activity in the Santa Barbara Channel region: Bulletin of the Seismological Society of America, v. 63, p. 1,257–1,273.

Mor, S.A., Kellogg, K.S., Stanley, R.G., Garrolo, L.D., Keller, E.A., and Brandt, T.R., 2009, Geologic map of the Santa Barbara coastal plain area, Santa Barbara County, California: U.S. Geological Survey Scientific Investigations Map 3001, scale 1:25,000, 1 sheet, pamphlet 38 p., available at <http://pubs.usgs.gov/sim/3001/>.

Southern California Earthquake Data Center, 2010, Southern California Earthquake Catalog: Southern California Earthquake Data Center database, available at <http://www.data.scec.org/sc-catalog/index.html>.

Wong, F.L., Phillips, E.L., Johnson, S.Y., and Sliter, R.W., 2012, Modeling of depth to base of Last Glacial Maximum and seafloor sediment thickness for the California State Waters Map Series, eastern Santa Barbara Channel, California: U.S. Geological Survey Open-File Report 2012–161, 16 p., available at <http://pubs.usgs.gov/ofr/2012/161/>.



Local (Offshore of Santa Barbara Map Area) and Regional (Offshore from Refugio Beach to Hueneme Canyon) Shallow-Subsurface Geology and Structure, Santa Barbara Channel, California

By
Samuel Y. Johnson, Eleyne L. Phillips, Andrew C. Ritchie, Florence L. Wong, Ray W. Sliter, Amy E. Draut, James E. Conrad, and Patrick E. Hart
2013

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

This map and report are electronic files, and they are not to be used for reproduction or distribution without the express written permission of the U.S. Geological Survey. The map and report are not to be used for reproduction or distribution without the express written permission of the U.S. Geological Survey.

For sale by U.S. Geological Survey, Information Services, Box 2580, Federal Center, Denver, CO 80225-1888. ASK-USGS

Original files available at <http://data.scec.org/>

Supporting Center: Johnson, S.Y., Phillips, E.L., Ritchie, A.C., Wong, F.L., Sliter, R.W., Draut, A.E., Conrad, J.E., and Hart, P.E., 2013, Local (Offshore of Santa Barbara Map Area) and Regional (Offshore from Refugio Beach to Hueneme Canyon) Shallow-Subsurface Geology and Structure, Santa Barbara Channel, California, report by Johnson, S.Y., Phillips, E.L., Sliter, R.W., Draut, A.E., Conrad, J.E., Hart, P.E., Ritchie, A.C., Wong, F.L., Johnson, S.Y., and Sliter, R.W., eds., Santa Clara, CA: U.S. Geological Survey, 11 sheets, 124,380. <http://dx.doi.org/10.3133/SIRM3281>

USGS Open-File Report 2012–161