

Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Scientific Investigations Report 2004-5151



U.S. Department of the Interior
U.S. Geological Survey
Prepared in cooperation with the
Florida Department of Environmental Protection

The following agencies provided most of the water-use data for this report:

*Florida Department of Environmental Protection; Tallahassee, Fla.
Northwest Florida Water Management District; Havana, Fla.
St. Johns River Water Management District; Palatka, Fla.
South Florida Water Management District; West Palm Beach, Fla.
Southwest Florida Water Management District; Brooksville, Fla.
Suwannee River Water Management District; Live Oak, Fla.*

*Appreciation is extended to the reviewers of this report:
John Trommer, Tampa, Fla., and Whitney Stringfield, Columbia, S.C.*

For Additional Information on the National Water-Use Program or the Florida Water-Use Program Contact:

National Water-Use Program
U.S. Geological Survey
Water Resources Division
1221 Sunrise Valley Drive, MS-414
Reston, VA 20192
URL <http://water.usgs.gov>

U.S. Geological Survey
Florida Integrated Science Center
2010 Levy Avenue
Tallahassee, FL 32310
URL <http://fl.water.usgs.gov>

**Front Cover Photographs — by Richard L. Marella
(clockwise from top left):**

*Limerock mine pit, Miami-Dade County
Portable traveling-gun vegetable irrigation in Miami-Dade County
Powerplant in South Florida
Public-supply water intake in Bay County
Domestic well in Putnam County
Recreational facility in Leon County*

Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

By Richard L. Marella

Prepared in cooperation with the
Florida Department of Environmental Protection

U.S. Department of the Interior
Gale A. Norton, Secretary

U.S. Geological Survey
Charles G. Groat, Director

U.S. Geological Survey, Reston, Virginia: 2004
For sale by U.S. Geological Survey, Information Services
Box 25286, Denver Federal Center
Denver, CO 80225

For more information about the USGS and its products:
Telephone: 1-888-ASK-USGS
World Wide Web: <http://www.usgs.gov/>

Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Contents

Abstract	1
Introduction	2
Purpose and Scope	2
Previous Investigations	2
Data Sources and Limitations	4
Acknowledgments	6
Water Withdrawals and Use	7
Water Source and Use Category	10
Public Supply	13
Domestic Self-Supplied	17
Commercial-Industrial Self-Supplied	19
Agricultural Self-Supplied (Irrigation and Nonirrigation)	22
Recreational Irrigation	26
Power Generation	29
Water Management Districts	31
Water Consumption and Discharges	35
Domestic Wastewater	36
Septic Tanks	38
Water Withdrawal Trends, 1970-2000	40
Selected References	44
Water-Use Bibliography for Florida	46
Florida	46
Northwest Florida Water Management District	47
St. Johns River Water Management District	47
South Florida Water Management District	48
Southwest Florida Water Management District	48
Suwannee River Water Management District	49
Additional Water-Use Reports or Papers	49
Appendix 1 and 2	
Purpose, Data Sources and Limitations	52
Selected References for Appendix 1 and 2	52
Appendix 1. Freshwater withdrawals by category for each county in Florida, 1965-2000	55
Appendix 2. Freshwater withdrawals by category for each Water Management District in Florida, 1975-2000	125
Glossary	132

Figures

1.	Graph showing historical and projected population of Florida, 1950-2020.....	2
2.	Counties and water management districts in Florida.....	3
3-5.	Graphs showing:	
3.	Total water withdrawals in Florida by source, 2000.....	7
4.	Average daily freshwater withdrawals by month in Florida, 2000.....	8
5.	Fresh ground- and surface-water withdrawals in Florida by category, 2000.....	8
6.	Map showing approximate areal extent throughout which principal aquifers in Florida are the primary source of ground water and quantity of ground-water withdrawals, 2000.....	10
7.	Map showing general location of hydrologic units in Florida and fresh ground- and surface-water withdrawals within these units, 2000.....	12
8-40.	Graphs showing:	
8.	Public-supply ground-water withdrawals in Florida by principal aquifer, 2000.....	13
9.	Average daily public-supply freshwater withdrawals by month in Florida, 2000.....	13
10.	Public-supply water use (deliveries) in Florida, 2000.....	15
11.	Historical public supply average daily per capita use in Florida, 1950-2000.....	15
12.	Historical public-supply freshwater withdrawals in Florida by source, 1950-2000..	17
13.	Domestic self-supplied ground-water withdrawals in Florida by principal aquifer, 2000.....	19
14.	Historical domestic self-supplied freshwater withdrawals in Florida, 1950-2000.....	19
15.	Commercial-industrial self-supplied ground-water withdrawals in Florida by principal aquifer, 2000.....	19
16.	Average daily commercial-industrial self-supplied freshwater withdrawals by month in Florida, 2000.....	21
17.	Commercial-industrial self-supplied freshwater use in Florida by major industrial type, 2000.....	21
18.	Historical commercial-industrial self-supplied freshwater withdrawals in Florida by source, 1950-2000.....	21
19.	Agricultural self-supplied ground-water withdrawals in Florida by principal aquifer, 2000.....	22
20.	Average daily agricultural self-supplied freshwater withdrawals by month in Florida, 2000.....	22
21.	Agricultural self-supplied freshwater use in Florida by major crop type, 2000.....	25
22.	Historical agricultural self-supplied freshwater withdrawals in Florida by source, 1950-2000.....	25
23.	Historical agricultural acreage in Florida for selected crops, 1970-2000.....	26
24.	Historical agricultural acreage irrigated in Florida by irrigation system type, 1980-2000.....	26
25.	Recreational irrigation ground-water withdrawals in Florida by principal aquifer, 2000.....	28
26.	Average daily recreational irrigation freshwater withdrawals by month in Florida, 2000.....	28
27.	Historical recreational irrigation freshwater withdrawals in Florida by source, 1985-2000.....	29
28.	Average daily power generation freshwater withdrawals by month in Florida, 2000.....	31
29.	Historical power generation water withdrawals in Florida by source, 1950-2000.....	31

30. Population and population served by public supply in Florida by water management district, 2000	32
31. Freshwater withdrawals in Florida by water management district, 2000.....	32
32. Fresh and saline water withdrawals in Florida by water management district, 2000	34
33. Historical freshwater withdrawals in Florida by water management district, 1975-2000.....	34
34. Historical fresh ground-water withdrawals in Florida by water management district, 1975-2000.....	35
35. Historical fresh surface-water withdrawals in Florida by water management district, 1975-2000.....	35
36. Freshwater withdrawals and estimated water consumption in Florida by category, 2000.....	35
37. Treated domestic wastewater discharges in Florida by primary disposal method, 2000.....	36
38. Historical treated domestic wastewater discharges in Florida, 1985-2000.....	38
39. Historical fresh and saline water withdrawals in Florida, 1950-2000	40
40. Historical freshwater withdrawals in Florida by source, 1950-2000	42

Tables

1. Total water withdrawals in Florida by category, 2000.....	7
2. Total water withdrawals in Florida by county, 2000.....	9
3. Total ground-water withdrawals by principal aquifer in Florida by county, 2000	11
4. Public-supplied population, water use, withdrawals, transfers, and treated water in Florida by county, 2000	14
5. Estimated public supply water use (deliveries), and per capita use in Florida by county, 2000.....	16
6. Domestic self-supplied population and water withdrawals in Florida by county, 2000.....	18
7. Commercial-industrial self-supplied water withdrawals in Florida by county, 2000.....	20
8. Agricultural self-supplied water withdrawals in Florida by county, 2000.....	23
9. Acres irrigated, irrigation system type, and water use in Florida by crop type, 2000	24
10. Recreational irrigation water withdrawals in Florida by county, 2000	27
11. Power generation water withdrawals in Florida by county, 2000	30
12. Water withdrawals by category in Florida by water management district, 2000.....	33
13. Estimated freshwater consumed by category in Florida, 2000.....	36
14. Treated domestic wastewater population served, discharge, and capacity in Florida by county, 2000	37
15. Treated domestic wastewater and septic tank population served and discharge in Florida by county, 2000	39
16. Historical population and water withdrawals in Florida by source, 1950-2000.....	41
17. Historical freshwater withdrawals in Florida by category, 1970-2000.....	43

Conversion Factors, Abbreviations and Acronyms

Multiply	By	To obtain
Area		
acre	4,047	square meter (m ²)
acre	0.00156	square mile (mi ²)
acre feet	0.325851	million gallons (Mgal)
square mile (mi ²)	2.590	square kilometer (km ²)
Flow rate		
gallon per day (gal/d)	3.785	liters per day (L/d)
million gallons per day (Mgal/d)	0.003785	million cubic meters per day (Mm ³ /d)
Energy		
gigawatthour (GWh)	1,000	megawatthour (mWh)
gigawatthour (GWh)	1,000,000	kilowatthour (kWh)

FASS	Florida Agricultural Statistic Service
FDACS	Florida Department of Agriculture and Consumer Services
FDEP	Florida Department of Environmental Protection
IFAS	Institute of Food and Agricultural Science (University of Florida)
NRCS	Natural Resources Conservation Service (U.S. Department of Agriculture)
NWFWMD	Northwest Florida Water Management District
SJRWMD	St. Johns River Water Management District
SFWMD	South Florida Water Management District
SWFWMD	Southwest Florida Water Management District
SRWMD	Suwannee River Water Management District
USGS	U.S. Geological Survey

Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

By Richard L. Marella

Abstract

In 2000, the estimated amount of water withdrawn in Florida was 20,148 million gallons per day (Mgal/d), of which 59 percent was saline and 41 percent was fresh. Ground water accounted for 62 percent of freshwater withdrawals and surface water accounted for the remaining 38 percent. Ninety-two percent of the 15.98 million people in Florida relied on ground water for their drinking water needs in 2000. Almost all of the saline water withdrawals (99.9 percent) were from surface water.

Public supply accounted for 43 percent of ground water withdrawn in 2000, followed by agricultural self-supplied (39 percent), commercial-industrial self-supplied (including mining) (8.5 percent), recreational irrigation (4.5 percent), domestic self-supplied (4 percent), and power generation (1 percent). Agricultural self-supplied accounted for 62 percent of fresh surface water withdrawn in 2000, followed by power generation (20 percent), public supply (8 percent), recreational irrigation (6 percent), and commercial-industrial self-supplied (4 percent). Almost all of saline water withdrawn was used for power generation.

The largest amount of freshwater was withdrawn in Palm Beach County and the largest amount of saline water was withdrawn in Hillsborough County. Significant withdrawals (more than 200 Mgal/d) of fresh ground water occurred in Miami-Dade, Polk, Orange, Palm Beach, Broward, and Collier Counties. Significant withdrawals (more than 200 Mgal/d) of fresh surface water occurred in Palm Beach, Hendry, and Escambia Counties. The South Florida Water Management District accounted for the largest amount of freshwater withdrawn (49 percent).

About 62 percent of the total ground water withdrawn was from the Floridan aquifer system; 17 percent was from the Biscayne aquifer. Most of the surface water used in Florida was from managed and maintained canal systems or large water bodies. Major sources of fresh surface water include the Caloosahatchee River, Deer Point Lake, Hillsborough River, Lake Okeechobee and associated canals, and the canals associated with the headwaters of the Upper St. Johns River.

Freshwater withdrawals increased 46 percent and saline water withdrawals increased 25 percent in Florida between 1970 and 2000. Ground-water withdrawals increased 82 percent, and surface-water withdrawals increased 10 percent during this period. Between 1970 and 2000, total freshwater withdrawals increased for public supply by 176 percent and for agricultural self-supplied by 87 percent; withdrawals for commercial-industrial self-supplied decreased by 37 percent, and power generation (thermoelectric) decreased by 57 percent. Recreational irrigation withdrawals increased 127 percent between 1985 and 2000. Between 1995 and 2000, freshwater withdrawals increased 13 percent, and saline withdrawals increased 9 percent.

An estimated 52 percent of the freshwater withdrawn in Florida was consumed; the remaining 48 percent was returned for further use. Domestic wastewater discharged in 2000 totaled 1,495 Mgal/d, of which 44 percent was discharged to surface waters, 34 percent to the ground through land application systems, and 22 percent to deep injection wells. Domestic wastewater discharge increased by 33 percent between 1985 and 2000, but decreased by 3 percent between 1995 and 2000. An estimated 11.21 million people were served by domestic wastewater systems in 2000, whereas the remaining 4.77 million people discharged wastewater to more than 1.95 million septic tanks. Discharge from the septic tanks was estimated to be 263 Mgal/d in 2000.

Introduction

Water is among Florida's most valued resources. The State has more than 1,700 streams and rivers and 7,800 freshwater lakes, and the entire State is underlain by aquifers capable of yielding significant quantities of freshwater to wells. These water resources provide for human and environmental needs. However, these water resources are finite, and growth in population, tourism, and agriculture are placing an increasing demand on them.

The population of Florida in 2000 totaled nearly 15.98 million (University of Florida, 2001), ranking fourth in the Nation (U.S. Census Bureau, 2001). This represents an increase of about 480 percent from the 1950 population of 2.77 million (Dietrich, 1978), and a 23.5 percent increase from the 1990 population of 12.94 million (University of Florida, 2001) (fig. 1). Florida's population is projected to reach nearly 17.4 million by the year 2005, and 21.7 million by the year 2020 (Smith and Nogle, 2001) (fig. 1). In addition to the resident population in Florida, an estimated 72 million people visited the State in 2000 (University of Florida, 2002). Freshwater is vital for Florida's permanent and seasonal residents and demands will continue to increase.




Figure 1. Historical and projected population of Florida, 1950-2020 (From Dietrich, 1978; Smith and Nogle, 2001; and University of Florida, 2001).

The agricultural sector in Florida also depends heavily on the State's water resources. In 2000, Florida accounted for 76 percent of the total citrus produced in the United States and was fourth in Nation net farm income (Florida Department of Agriculture and Consumer Services, 2002). Agricultural production should continue to flourish because of the subtropical climate and demands from the growing population. Information

on the amount of water required to support future agricultural growth is needed.

To provide accurate information on water use and trends, the U.S. Geological Survey (USGS), in cooperation with the Florida Department of Environmental Protection (FDEP) and the Northwest Florida, St. Johns River, South Florida, Southwest Florida, and Suwannee River Water Management Districts, estimate water use in Florida at 5-year intervals. Water-use estimates for 2000 are presented in this report.

Purpose and Scope

This report summarizes the quantities of water withdrawn, consumed, and discharged in 2000, and indicates trends in water use. Overall, the report provides a basis for estimating water budgets and projecting future water needs. Data are presented on water withdrawals in Florida for each of the following water use categories: public supply, domestic self-supplied, commercial-industrial self-supplied (including mining uses), agricultural self-supplied (including irrigation and nonirrigation uses), recreational irrigation, and power generation. Information concerning instream (nonwithdrawal) water use such as hydroelectric power generation, navigation, water-based recreation, propagation of fish and wildlife, and dilution and conveyance of liquid or solid waste is not included.

Within each category, withdrawal data are presented by source (ground or surface water), and where sufficient data are available, seasonal and historical patterns of water use are described. Data also are presented by county and water management district (fig. 2) for each water use category. Information about specific public-supply water systems and domestic wastewater facilities can be obtained by contacting the USGS office in Tallahassee or by visiting the USGS web site at http://fl.water.usgs.gov/WaterUse/Water_Use_web/index.htm.

Previous Investigations

This report is the ninth in a series of reports documenting the results of water-use investigations in Florida. Statewide water-use data for Florida were published for 1965 and 1970 (Pride, 1973 and 1975); for 1975, 1977, and 1980 (Leach, 1978, 1983; and Leach and Healy, 1980) and for 1985, 1990, and 1995 (Marella, 1988, 1992, 1999). These reports included assessments of all water uses in Florida (including public-supply, domestic self-supplied, commercial-industrial self-supplied, agricultural irrigation and nonirrigation, recreational irrigation, and power generation water uses) by county. Historical water-use data for each county are presented in appendix 1. Individual county tables in appendix 1 detail freshwater withdrawals by category for all data collected between 1965 and 2000.




Figure 2. Counties and water management districts in Florida.

Prior to 1965, state water-use data were only published at the national level. Nationwide summaries of water-use data were published for 1950, 1955, and 1960 (MacKichan, 1951, 1957; and MacKichan and Kammerer, 1961). These reports included detailed water-use data at the state level, but did not include water-use data for counties. Nationwide summaries, including data for Florida, also were published by the USGS for 1965, 1970, 1975 (Murray, 1968; and Murray and Reeves, 1972, 1977); for 1980, 1985, 1990, 1995 (Solley and others,

1983, 1988, 1993, 1998), and for 2000 (Hutson and others, 2004).

Additional water-use reports have been published by selected water management districts between 1975 and 2000 (see water-use bibliography for Florida). The St. Johns River and the Southwest Florida Water Management Districts have published annual water-use reports since 1977; the Northwest Florida, South Florida, and the Suwannee River Water Management Districts have intermittently published reports between 1977 and 1985. Detailed water-use data for 2000 were pub-

4 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

lished by the Southwest Florida Water Management District (2002), and a summary fact sheet for 2000 was published by the St. Johns River Water Management District (2004). Historical water-use data for each water management district are presented in appendix 2. Individual district tables in appendix 2 detail freshwater withdrawals by category for all data collected between 1975 and 2000.

Data Sources and Limitations

As part of the USGS National Water-Use Information Program, water-use data are collected and compiled for each state every 5 years (Solley and others, 1988). Data are reported for each state by water-use category, county, hydrologic unit (basin), and aquifer. Water-use data for Florida were compiled through an ongoing cooperative program with the FDEP, as part of the 2000 National Water-Use Information Program. Data were obtained from the FDEP (Drinking Water and Wastewater Sections), the Northwest Florida Water Management District (NFWMD), the St. Johns River Water Management District (SJRWMD), the South Florida Water Management District (SFWMD), the Southwest Florida Water Management District (SWFWMD), and the Suwannee River Water Management District (SRWMD), as well as from various utilities, industries, and power companies. Specific data sources for each category or source are listed below.

Public supply. Data for public-supply withdrawals were obtained from the FDEP's Drinking Water Section (monthly operating reports), the five water management districts (consumptive water-use permit files or annual reports), or directly from the water supplier. All water use values for this category are from metered data.

The population served value for each public supplier was determined by one of three sources: (1) estimated using the number of service connections (obtained from the FDEP monthly operating reports) multiplied by the number people per household per county (Smith and Cody, 2001); (2) estimated from published population studies for selected cities (University of Florida, 2001); and (3) estimated by the individual public suppliers and reported to FDEP (Drinking Water Quick Look Report, Florida Department of Environmental Protection, written commun., April 2000) or to their respective water management district. Population served values in the Northwest Florida, the South Florida, and the Suwannee River Water Management Districts were estimated by the USGS. Population served estimates for some counties within the South Florida Water Management District also were made by the district using housing counts from the 2000 Census data overlaid by utility service areas. Estimates for population served in the St. Johns River and the Southwest Florida Water Management Districts were made by the respective district (St. Johns River Water Management District, 2004; Southwest Florida Water Management District, 2002), however, values for the Southwest Florida Water Management District were modified to remove seasonal population estimates.

Domestic self-supplied. Domestic self-supplied populations are derived by subtracting the population served by public-supplied systems from the total county population. Domestic self-supplied withdrawals are calculated by multiplying the 2000 statewide domestic per capita use of 106 gallons per day by the self-supplied population served for each county. Withdrawal estimates for the counties within the Southwest Florida Water Management District were calculated by the district (Southwest Florida Water Management District, 2002). All water used for domestic self-supplied is assumed to be obtained from ground water. Values for this category are estimated.

Commercial-industrial self-supplied. Data for commercial, industrial, and mining withdrawals were obtained from the FDEP's Drinking Water Section (monthly operating reports), the five water management districts (consumptive water-use permit files or annual reports), or directly from the user. Most of the water-use values for this category are from metered data.

Agricultural self-supplied. Water withdrawn for irrigation was estimated by the water management districts based on crop acreage multiplied by a use coefficient (usually in inches per acre) generated from selected irrigation models for each crop. Only a small percentage (about 10-15) of the water-use estimates for 2000 were derived from actual metered data.

Acreage data were obtained by the water management districts from a variety of sources including, (1) the University of Florida, Institute of Food and Agriculture Science (IFAS), County Extension Offices; (2) the water management districts' consumptive water-use permit files; (3) the Florida Department of Agriculture and Consumer Services (FDACS); Agricultural Statistics Service, Commercial Citrus Inventory 2000, Field Crops Summary 2000, and the Vegetable Summary 1999-2000 (Florida Agricultural Statistics Service, 2000; 2001a; 2001b); (4) the FDACS, Division of Plant Industry, Annual Report (Florida Department of Agriculture and Consumer Services, 2000); and (5) the U.S. Department of Agriculture, 1997 Census of Agriculture, Florida (U.S. Department of Agriculture, 1999). The total number of acres for selected crops for each county was obtained using these sources, as well as the acreage that was irrigated. The method of irrigation was obtained from many of the same sources or was estimated by the individual water management districts.

The supplemental irrigation coefficient used to determine the amount of water needed per crop was obtained from several irrigation models and publications including (1) the modified Blaney-Criddle irrigation model; "Irrigation Water Requirements, 1970, revised" (U.S. Soil Conservation Service, 1970); (2) the University of Florida "AFSIRS" computer model (Smajsrta, 1986); and, (3) the "Florida Irrigation Guide" (U.S. Soil Conservation Service, 1982). These models or publications either use actual or long-term averages of rainfall, and temperature (Jones and others, 1984). Other sources of supplemental irrigation coefficients used for determining irrigation needs included actual metered data for selected crops, or those coefficients provided by agriculture authorities or the water management districts.

Supplemental irrigation application rates (in inches per acre) generated from these sources were multiplied by the number of irrigated acres to yield water use. From these values, estimates of the water sources (ground or surface) were made by local county extension service personnel or by using information from the water management district's consumptive water-use permit files. Supplemental irrigation rates include the amount of water needed to grow each crop, the amount of water necessary to overcome the inefficiency of the irrigation system, and water used for frost and freeze protection.

Withdrawals were estimated for livestock and fish farming by multiplying the total number of cattle (including dairy cattle), chickens, horses, and swine for each county by a water-use coefficient per animal type. The number of animals per county was obtained from the FDACS, Agricultural Statistics Service, Livestock Summary 2000 (Florida Agricultural Statistics Service, 2001c). Water-use coefficients were obtained from the University of Florida, IFAS (St. Johns River Water Management District, 1984). Livestock estimates were not made for those counties within the St. Johns River Water Management District for 2000.

Estimates of the quantity of ground or surface water used were provided by local county extension service personnel or were determined by using information from the water management district's consumptive water-use data base. If no information or data source was available, a ratio of 90 percent ground water and 10 percent surface water was used based on previous studies. The quantity of water withdrawn for fish farming was obtained from the water management district's consumptive water-use permit files or was estimated based on a coefficient of 50 inches per pond acre. Pond acreage was obtained from the water management districts or the U.S. Department of Agriculture, 1997 Census of Agriculture, Florida (U.S. Department of Agriculture, 1999).

Recreational irrigation. Water withdrawal values for recreational irrigation were estimated based on irrigated acreage multiplied by a coefficient (usually in inches per acre) generated from selected irrigation models for turf grass. Only a small percentage (about 10-15) of the water-use estimates for 2000 were derived from actual metered data.

The total number of acres and acres irrigated were obtained by the water management districts from a variety of sources that included: (1) the University of Florida, IFAS, County Extension Offices; (2) the water management districts' consumptive water-use permit files and the districts needs and sources documents; or were estimated from the number of golf courses or number of holes per county obtained from the Florida Sports Foundation (2000) or the National Golf Foundation (2001).

The supplemental irrigation coefficient used to determine the amount of water needed was obtained from several irrigation models or publications including: (1) the modified Blaney-Criddle irrigation model, and Irrigation Water Requirements, 1970, revised (U.S. Soil Conservation Service, 1970); (2) the University of Florida AFSIRS computer model (Smajstrla, 1986); and (3) the Florida Irrigation Guide (U.S. Soil Conserva-

tion Service, 1982). Other sources of supplemental irrigation coefficients were derived from metered data or were provided by turf-grass authorities.

Supplemental irrigation application rates (in inches per acre) generated from these sources were multiplied by the number of irrigated acres, and a water-use value was calculated. From this value, estimates of the water sources (ground or surface) were provided by the water management district's consumptive water-use permit files.

Power generation. Data for power generation withdrawals were obtained from the water management district's (consumptive water-use permit files or annual reports), or directly from the power companies. Withdrawal data were collected for ground and surface waters, both fresh and saline sources. Information on the amount of water purchased from public supplies was obtained from each facility along with the total gross power generated. Most of the water-use values for this category are from metered data.

Wastewater discharges. Data for domestic wastewater discharges were obtained from the FDEP's Domestic Wastewater Section, 2000 Reuse Inventory (Florida Department of Environmental Protection, 2001), or from data obtained from the monthly operating reports submitted to FDEP (Joseph Doker, Florida Department of Environmental Protection, written commun., October 2003). Most discharge values for this category are from metered data. The population served was estimated for each county by increasing the 1995 county population served (Marella, 1999) by the percent of growth by county between 1995 and 2000 (University of Florida, 2001), or by subtracting out the population served by septic tanks from the total county population. The population served by septic tanks in each county was estimated by multiplying the number of people per household (Smith and Cody, 2001) by the number of septic tanks per county. A few counties showed a decline in domestic wastewater population served between 1995 and 2000, most likely due to overestimating the 1995 population served.

Aquifer withdrawals. Estimates of water withdrawn by aquifer were made for each category. For public supply, commercial-industrial (including mining) self-supplied, and power generation information for the aquifer of withdrawal were obtained from water management district's consumptive water-use permits for each individual user. Estimates were made for domestic self-supplied, agricultural self-supplied, and recreational irrigation using information obtained from several ground-water studies recently conducted throughout the State that detailed estimates of withdrawals for selected aquifers in specific counties or areas (Duerr and others, 1988, Knowles and others, 2002, Sepulveda, 2002, and Shoemaker and Edwards, 2003) or were made from information obtained from local county agencies (County Health Departments) that regulate domestic wells. In counties with no information, estimates were made by assuming 90 percent of the water withdrawn was from the primary aquifer used for public supply and the remaining 10 percent from the local water table or shallow aquifer.

Accuracy. Water withdrawals and use data presented herein represent the average daily quantities used, as derived from annual data, and are expressed in million gallons per day (Mgal/d). Water-use values presented in the tables are reported to two places to the right of the decimal or to the nearest 10,000 gallons per day (gal/d). Water-use values in the text are rounded to the nearest million gallons per day, and percentages are rounded. Water-use data published herein report may not be identical to the water-use data published by the water management districts or FDEP because of differences in data-collection procedures, categories, and methodology.

The accuracy of these values varies from category to category. Public-supply values usually are more accurate because most public-supply systems meter water usage, whereas agricultural self-supplied or domestic self-supplied values usually are estimated. However, a small percentage of the irrigation values (agricultural and recreational) for 2000 were obtained from metered data.

Changes. Some changes in water-use categories have occurred in Florida between 1970 and 2000. Most categories have remained the same since 1970, with the following exceptions. During the 1970's, rural water use consisted of domestic self-supplied and livestock. In the 1980's, livestock was added to the agricultural, and domestic self-supplied became a separate category. For many reports, agricultural water use has included all irrigation, but in 1995, golf course and other recreational irrigation were removed from the agricultural water-use category and placed within a separate category (recreational irrigation). Because estimates of water use for golf courses was differentiated for 1985 and 1990, data for recreational irrigation was subtracted from agriculture for those years. Agricultural self-supplied was called irrigation in the 1970's, then became agricultural irrigation in the 1980's, with the addition of livestock and removal of recreation irrigation. Commercial-industrial self-supplied was previously called industrial water use, and power generation was previously called thermoelectric power generation.

Miscellaneous water withdrawals and uses included in county totals presented in the 1985 water-use report (Marella, 1988) included water withdrawals for residential lawn irrigation, residential heat pumps and air-conditioning units, and water discharged from free-flowing wells. Because of the inconsistency in data from county to county for these uses, they were not included in the 1985 statewide totals (Marella, 1988). Since 1985, some data on residential lawn watering for several counties have been updated and are included in the recreational irrigation section of this report under turf grass. Water used by residential heat pumps and air conditioning units and water discharged from free-flowing wells remain inconsistent between counties and are not included in this report.

Data for Miami-Dade County prior to 2000 are reported under Dade County. In 1997, Dade County officially changed to Miami-Dade County. All data presented herein are listed under Miami-Dade County.

Acknowledgments


The author gratefully acknowledges the Florida Department of Environmental Protection--Office of Water Policy for its cooperation in the State water-use program, and extends a special thanks to the following individuals who provided program support or data and technical assistance; Joseph Doker, Lauren Walker-Coleman, Kathleen Greenwood, Kenna Study, and Thomas Swihart, (Tallahassee), as well as the many individuals in the Drinking Water and Wastewater Programs from the FDEP district offices in Ft. Myers, Jacksonville, Orlando, Pensacola, Tampa, and West Palm Beach.

Special thanks are extended to the Executive Directors of the five water management districts (Northwest Florida Water Management District, St. Johns River Water Management District, South Florida Water Management District, Southwest Florida Water Management District, and the Suwannee River Water Management District) for their participation and staff cooperation in providing water-use data. The participation of the following individuals is acknowledged for their data collection and tabulation efforts, and for their technical advice and support: Jeff Bonekemper, Francis Flores, and Tyler Macmillan at the Northwest Florida Water Management District in Havana; Bruce Florence, Jeanne Thompson, and Mary Beth Wilder at the St. Johns River Water Management District in Palatka; Chad Brcka, Lori Burke, Elizabeth Ellis, John Floyd, Sharon Fowler, Barbara Fronczek, David Gilpin-Hudson, Kurt Leckler, Richard March, William Rasperger, and Donna Rickabus at the South Florida Water Management District in West Palm Beach; Albert Bond, C. Donald Rome, and Kathleen Scott at the Southwest Florida Water Management District in Brooksville; Michael Fuller and Melanie Sardina at the Suwannee River Water Management District in Live Oak.

Additionally, special thanks are extended to the many utility operators, plant managers, and individuals who provided data vital to the completion of this effort. These include staff from: Bay County Public Utilities, Broward County Public Health Unit, Florida Power and Light Company, Florida Power Corporation (Progress Energy Company), Florida Rock Industries, Florida Water Services, Gainesville Regional Utilities, Gulf Power Company, Jacksonville Electric Authority (JEA), City of Key West, Lakeland Electric and Water, Lee County Public Health Unit, Miami-Dade Water and Sewer, Orlando Utilities Commission, Palm Beach County Public Health Unit, Polk County Public Health Unit, Sarasota County Public Health Unit, Tampa Electric, City of Tallahassee, University of Florida (Institute of Food and Agriculture Science), U.S. Department of Agriculture (Natural Resources Conservation Service), and the many others who provided data and technical guidance.

Water Withdrawals and Use

In 2000, the total amount of water withdrawn in Florida was estimated at 20,148 Mgal/d (table 1), of which 59 percent was saline water and 41 percent was freshwater (fig. 3). Ground water accounted for nearly 62 percent of freshwater withdrawals, and surface water accounted for the remaining 38 percent. Almost all of the saline water withdrawals (99.9 percent) were from surface water (table 1), which was used for once-through cooling at 23 power generating plants. The saline water was withdrawn from bays or rivers along the coast, and then returned to those sources. Florida ranked second in the Nation after California in saline water withdrawals in 2000 (Hutson and others, 2004).



Note: Saline ground water was less than 0.1 percent

Figure 3. Total water withdrawals in Florida by source, 2000.

Table 1. Total water withdrawals in Florida by category, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day]

Florida 2000	Freshwater			Saline Water		
	Ground	Surface	Total	Ground	Surface	Total
Public Supply	2,199.36	237.43	2,436.79	0.00	0.00	0.00
Domestic self-supplied	198.68	0.00	198.68	0.00	0.00	0.00
Commercial-industrial self-supplied	430.70	132.60	563.30	0.00	1.18	1.18
Agricultural self-supplied	1,989.95	1,933.06	3,923.01	0.00	0.00	0.00
Recreational irrigation	230.45	181.28	411.73	0.00	0.00	0.00
Power generation	29.53	628.73	658.26	3.82	11,950.82	11,954.64
TOTALS	5,078.67	3,113.10	8,191.77	3.82	11,952.00	11,955.82

The amount of freshwater withdrawn monthly fluctuates because of variations in temperature, precipitation (rainfall), crop production, and tourism. More than one-half (54 percent) of the freshwater withdrawn in 2000 was between February and June (fig. 4). Seasonal variation in withdrawals is normal; however, severe drought conditions were still present in Florida during the first half of 2000 (University of Nebraska, 2004). The dry conditions were present throughout most of the State, and three of the five water management districts (SJRWMD, SFWMD, and SWFWMD) issued water shortage mandates to help curb water use during the spring. Due to the dry conditions, water demands for all irrigation purposes (agriculture, recreation, and residential lawns) were higher-than-normal in 2000.

Public supply (43 percent) and agricultural self-supplied (39 percent) were the largest uses of ground water in 2000, followed by commercial-industrial self-supplied (including mining) (8.5 percent), recreational irrigation (4.5 percent), domestic self-supplied (4 percent), and power generation

(1 percent) (fig. 5 and table 1). Agricultural self-supplied (62 percent) was the largest user of fresh surface water in 2000, followed by power generation (20 percent), public supply (8 percent), and recreational irrigation (6 percent), and commercial-industrial self-supplied (4 percent) (fig. 5 and table 1). Overall, agricultural self-supplied accounted for 48 percent of the total freshwater withdrawn (ground and surface). Power generation accounted for nearly all (99.9 percent) of the saline water withdrawn.

The largest amount of freshwater was withdrawn in Palm Beach County, and the largest amount of saline water was withdrawn in Hillsborough County (table 2). Significant withdrawals (more than 200 Mgal/d) of fresh ground water were in Miami-Dade, Polk, Orange, Palm Beach, Broward, and Collier Counties. Significant withdrawals (more than 200 Mgal/d) of fresh surface water occurred in Palm Beach, Hendry, and Escambia Counties.

8 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000




Figure 4. Average daily freshwater withdrawals by month in Florida, 2000.




Figure 5. Fresh ground- and surface-water withdrawals in Florida by category, 2000.

Table 2. Total water withdrawals in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day]

County	Ground water			Surface water			Total water		
	Fresh	Saline	Total	Fresh	Saline	Total	Fresh	Saline	Total
Alachua	59.58	0.00	59.58	0.57	0.00	0.57	60.15	0.00	60.15
Baker	6.15	0.00	6.15	1.73	0.00	1.73	7.88	0.00	7.88
Bay	12.22	0.00	12.22	45.68	265.05	310.73	57.90	265.05	322.95
Bradford	5.79	0.00	5.79	0.06	0.00	0.06	5.85	0.00	5.85
Brevard	143.53	0.00	143.53	39.20	1,079.28	1,118.48	182.73	1,079.28	1,262.01
Broward	273.46	0.00	273.46	28.35	1,512.39	1,540.74	301.81	1,512.39	1,814.20
Calhoun	4.34	0.00	4.34	1.79	0.00	1.79	6.13	0.00	6.13
Charlotte	35.95	0.00	35.95	28.04	0.00	28.04	63.99	0.00	63.99
Citrus	30.05	0.00	30.05	1.29	393.90	395.19	31.34	393.90	425.24
Clay	33.57	0.00	33.57	0.52	0.00	0.52	34.09	0.00	34.09
Collier	202.49	0.00	202.49	27.55	0.00	27.55	230.04	0.00	230.04
Columbia	13.88	0.00	13.88	0.21	0.00	0.21	14.09	0.00	14.09
DeSoto	123.87	0.00	123.87	9.51	0.00	9.51	133.38	0.00	133.38
Dixie	3.50	0.00	3.50	0.03	0.00	0.03	3.53	0.00	3.53
Duval	154.33	0.00	154.33	1.65	661.15	662.80	155.98	661.15	817.13
Escambia	91.94	0.00	91.94	225.31	0.00	225.31	317.25	0.00	317.25
Flagler	24.55	0.00	24.55	3.60	0.00	3.60	28.15	0.00	28.15
Franklin	2.25	0.00	2.25	0.00	0.00	0.00	2.25	0.00	2.25
Gadsden	8.93	0.00	8.93	7.40	0.00	7.40	16.33	0.00	16.33
Gilchrist	15.83	0.00	15.83	0.32	0.00	0.32	16.15	0.00	16.15
Glades	21.03	0.00	21.03	53.64	0.00	53.64	74.67	0.00	74.67
Gulf	3.10	0.00	3.10	1.65	0.00	1.65	4.75	0.00	4.75
Hamilton	41.61	0.00	41.61	0.11	0.00	0.11	41.72	0.00	41.72
Hardee	90.53	0.00	90.53	0.61	0.00	0.61	91.14	0.00	91.14
Hendry	197.12	0.00	197.12	314.99	0.00	314.99	512.11	0.00	512.11
Hernando	49.46	0.00	49.46	0.99	0.00	0.99	50.45	0.00	50.45
Highlands	157.29	0.00	157.29	17.26	0.00	17.26	174.55	0.00	174.55
Hillsborough	197.44	0.00	197.44	94.28	3,188.00	3,282.28	291.72	3,188.00	3,479.72
Holmes	3.77	0.00	3.77	0.32	0.00	0.32	4.09	0.00	4.09
Indian River	87.36	0.00	87.36	161.08	39.01	200.09	248.44	39.01	287.45
Jackson	21.87	0.00	21.87	91.23	0.00	91.23	113.10	0.00	113.10
Jefferson	8.39	0.00	8.39	0.17	0.00	0.17	8.56	0.00	8.56
Lafayette	6.78	0.00	6.78	0.16	0.00	0.16	6.94	0.00	6.94
Lake	90.84	0.00	90.84	9.77	0.00	9.77	100.61	0.00	100.61
Lee	120.80	0.00	120.80	39.77	492.27	532.04	160.57	492.27	652.84
Leon	45.36	0.00	45.36	0.30	0.00	0.30	45.66	0.00	45.66
Levy	28.44	0.00	28.44	2.63	0.00	2.63	31.07	0.00	31.07
Liberty	1.82	0.00	1.82	0.00	0.00	0.00	1.82	0.00	1.82
Madison	9.06	0.00	9.06	0.17	0.00	0.17	9.23	0.00	9.23
Manatee	122.30	0.00	122.30	64.65	0.00	64.65	186.95	0.00	186.95
Marion	67.68	0.00	67.68	2.10	0.00	2.10	69.78	0.00	69.78
Martin	43.60	0.00	43.60	155.27	0.00	155.27	198.87	0.00	198.87
Miami-Dade	537.91	3.68	541.59	28.68	33.19	61.87	566.59	36.87	603.46
Monroe	1.44	0.14	1.58	0.62	0.00	0.62	2.06	0.14	2.20
Nassau	46.53	0.00	46.53	0.48	1.18	1.66	47.01	1.18	48.19
Okaloosa	32.15	0.00	32.15	0.46	0.00	0.46	32.61	0.00	32.61
Okeechobee	59.80	0.00	59.80	12.03	0.00	12.03	71.83	0.00	71.83
Orange	281.36	0.00	281.36	16.72	0.00	16.72	298.08	0.00	298.08
Osceola	117.52	0.00	117.52	26.03	0.00	26.03	143.55	0.00	143.55
Palm Beach	273.70	0.00	273.70	1,005.48	433.02	1,438.50	1,279.18	433.02	1,712.20
Pasco	139.69	0.00	139.69	2.24	1,956.50	1,958.74	141.93	1,956.50	2,098.43
Pinellas	43.20	0.00	43.20	3.58	419.11	422.69	46.78	419.11	465.89
Polk	330.53	0.00	330.53	33.73	0.00	33.73	364.26	0.00	364.26
Putnam	40.35	0.00	40.35	48.92	0.00	48.92	89.27	0.00	89.27
St. Johns	52.55	0.00	52.55	3.16	0.00	3.16	55.71	0.00	55.71
St. Lucie	80.84	0.00	80.84	198.92	1,477.95	1,676.87	279.76	1,477.95	1,757.71
Santa Rosa	30.85	0.00	30.85	0.35	0.00	0.35	31.20	0.00	31.20
Sarasota	40.37	0.00	40.37	5.91	0.00	5.91	46.28	0.00	46.28
Seminole	88.47	0.00	88.47	1.78	0.00	1.78	90.25	0.00	90.25
Sumter	26.80	0.00	26.80	17.69	0.00	17.69	44.49	0.00	44.49
Suwannee	26.41	0.00	26.41	101.41	0.00	101.41	127.82	0.00	127.82
Taylor	46.79	0.00	46.79	3.02	0.00	3.02	49.81	0.00	49.81
Union	2.91	0.00	2.91	0.02	0.00	0.02	2.93	0.00	2.93
Volusia	97.02	0.00	97.02	137.72	0.00	137.72	234.74	0.00	234.74
Wakulla	4.79	0.00	4.79	28.38	0.00	28.38	33.17	0.00	33.17
Walton	10.51	0.00	10.51	1.81	0.00	1.81	12.32	0.00	12.32
Washington	4.32	0.00	4.32	0.00	0.00	0.00	4.32	0.00	4.32
State totals	5,078.67	3.82	5,082.49	3,113.10	11,952.00	15,065.10	8,191.77	11,955.82	20,147.59

Water Source and Use Category

Ground water was the source of drinking water for 14.75 million people (92 percent) in Florida during 2000. Florida, the largest user of ground water east of the Mississippi River, ranked fifth in the Nation in ground-water withdrawals in 2000 (Hutson and others, 2004). Ground water is available throughout the State, and generally needs little or no treatment prior to use. Florida accounted for about 78 percent of the total water withdrawn from the Floridan aquifer system (4,020 Mgal/d) in the southeastern United States (Marella and Berndt, 2005). About 62 percent of the ground water withdrawn in

Florida was obtained from the Floridan aquifer system (3,125 Mgal/d), which underlies most of the State (fig. 6). Polk, Orange, Hillsborough, and Duval Counties were the largest users of water from the Floridan aquifer system (table 3). The Biscayne aquifer was the source for about 17 percent of ground water, whereas the remaining 21 percent was obtained from unnamed surficial aquifers, the intermediate aquifer, and the sand-and-gravel aquifer (fig. 6 and table 3). Surficial aquifers are primarily used for household wells (domestic self-supplied) or in areas where the Floridan aquifer system is nonpotable or too deep to be used economically.




Figure 6. Approximate areal extent throughout which principal aquifers in Florida are the primary source of ground water and quantity of ground-water withdrawals, 2000. (Map modified from Vecchioli and Foose, 1985.)

Table 3. Total ground-water withdrawals by principal aquifer in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day]

County	Biscayne Aquifer	Floridan Aquifer	Intermediate Aquifer	Sand-Gravel Aquifer	Surficial Aquifer	Total
Alachua	0.00	59.16	0.00	0.00	0.42	59.58
Baker	0.00	5.95	0.00	0.00	0.20	6.15
Bay	0.00	12.02	0.00	0.00	0.20	12.22
Bradford	0.00	5.60	0.00	0.00	0.19	5.79
Brevard	0.00	123.55	0.00	0.00	19.98	143.53
Broward	273.37	0.09	0.00	0.00	0.00	273.46
Calhoun	0.00	4.25	0.00	0.00	0.09	4.34
Charlotte	0.00	7.32	27.11	0.00	1.52	35.95
Citrus	0.00	29.35	0.00	0.00	0.70	30.05
Clay	0.00	32.97	0.00	0.00	0.60	33.57
Collier	0.00	4.37	115.86	0.00	82.26	202.49
Columbia	0.00	13.51	0.00	0.00	0.37	13.88
DeSoto	0.00	110.01	13.86	0.00	0.00	123.87
Dixie	0.00	3.42	0.00	0.00	0.08	3.50
Duval	0.00	150.00	0.00	0.00	4.33	154.33
Escambia	0.00	0.00	0.00	91.94	0.00	91.94
Flagler	0.00	20.58	0.00	0.00	3.97	24.55
Franklin	0.00	2.23	0.00	0.00	0.02	2.25
Gadsden	0.00	8.75	0.00	0.00	0.18	8.93
Gilchrist	0.00	15.70	0.00	0.00	0.13	15.83
Glades	0.00	7.65	0.71	0.00	12.67	21.03
Gulf	0.00	2.57	0.00	0.00	0.53	3.10
Hamilton	0.00	41.54	0.00	0.00	0.07	41.61
Hardee	0.00	81.73	8.80	0.00	0.00	90.53
Hendry	0.00	0.25	126.17	0.00	70.70	197.12
Hernando	0.00	49.32	0.00	0.00	0.14	49.46
Highlands	0.00	133.77	23.43	0.00	0.09	157.29
Hillsborough	0.00	194.86	2.34	0.00	0.24	197.44
Holmes	0.00	3.64	0.00	0.00	0.13	3.77
Indian River	0.00	74.01	0.00	0.00	13.35	87.36
Jackson	0.00	21.55	0.00	0.00	0.32	21.87
Jefferson	0.00	8.31	0.00	0.00	0.08	8.39
Lafayette	0.00	6.72	0.00	0.00	0.06	6.78
Lake	0.00	90.41	0.00	0.00	0.43	90.84
Lee	0.00	19.40	32.71	0.00	68.69	120.80
Leon	0.00	44.93	0.00	0.00	0.43	45.36
Levy	0.00	28.04	0.00	0.00	0.40	28.44
Liberty	0.00	1.77	0.00	0.00	0.05	1.82
Madison	0.00	8.94	0.00	0.00	0.12	9.06
Manatee	0.00	118.25	4.05	0.00	0.00	122.30
Marion	0.00	66.04	0.00	0.00	1.64	67.68
Martin	0.00	5.79	0.00	0.00	37.81	43.60
Miami-Dade	537.91	3.68	0.00	0.00	0.00	541.59
Monroe	0.11	1.37	0.00	0.00	0.10	1.58
Nassau	0.00	43.62	0.00	0.00	2.91	46.53
Okaloosa	0.00	32.02	0.00	0.00	0.13	32.15
Okeechobee	0.00	47.76	0.00	0.00	12.04	59.80
Orange	0.00	280.48	0.00	0.00	0.88	281.36
Osceola	0.00	114.65	0.00	0.00	2.87	117.52
Palm Beach	63.24	6.45	0.00	0.00	204.01	273.70
Pasco	0.00	139.24	0.00	0.00	0.45	139.69
Pinellas	0.00	43.16	0.00	0.00	0.04	43.20
Polk	0.00	317.73	11.56	0.00	1.24	330.53
Putnam	0.00	39.85	0.00	0.00	0.50	40.35
St. Johns	0.00	46.69	0.00	0.00	5.86	52.55
St. Lucie	0.00	46.98	0.00	0.00	33.86	80.84
Santa Rosa	0.00	9.03	0.00	21.82	0.00	30.85
Sarasota	0.00	27.23	13.10	0.00	0.04	40.37
Seminole	0.00	88.20	0.00	0.00	0.27	88.47
Sumter	0.00	26.34	0.00	0.00	0.46	26.80
Suwannee	0.00	26.14	0.00	0.00	0.27	26.41
Taylor	0.00	46.70	0.00	0.00	0.09	46.79
Union	0.00	2.80	0.00	0.00	0.11	2.91
Volusia	0.00	96.72	0.00	0.00	0.30	97.02
Wakulla	0.00	4.79	0.00	0.00	0.00	4.79
Walton	0.00	10.49	0.00	0.00	0.02	10.51
Washington	0.00	4.18	0.00	0.00	0.14	4.32
State totals	874.63	3,124.62	379.70	113.76	589.78	5,082.49

12 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Florida ranked twenty-seventh in the Nation in fresh surface-water withdrawals in 2000 (Hutson and others, 2004). The primary users of surface water throughout Florida are irrigators and power plants, which used about 82 percent of the fresh water. The majority of fresh surface water used is in the Southern Florida subregion (fig. 7) and is associated with Lake Okeechobee and the Everglades Agricultural Area of Glades, Hendry, and Palm Beach Counties (fig. 7). This area is intensively irrigated for sugarcane and vegetables and accounted for nearly two-thirds of the surface-water withdrawals during 2000.

Most of the surface water used in Florida is from managed and maintained canal systems or large water bodies. Major sources of fresh surface water for irrigation include Lake

Okeechobee and associated canals (Glades, Hendry, Martin, Palm Beach, and St. Lucie Counties), the Caloosahatchee River (Glades, Hendry, and Lee Counties), and the headwaters of the Upper St. Johns River (Brevard, Indian River, Okeechobee, and St. Lucie Counties). Surface water from these sources is most often diverted through canals or ditches, then pumped or gravity-fed onto fields or groves. A large percentage of the water that is flooded onto fields or groves is unused and pumped back into the canals or ditches for further use. Throughout the remainder of Florida, surface water for irrigation is obtained from local canals, ditches, lakes, ponds, small rivers, creeks, or tributaries. Many of the canals, ditches, or ponds that are used for irrigation are augmented with ground water.




Figure 7. General location of hydrologic units in Florida and fresh ground- and surface-water withdrawals within these units, 2000. (Map from U.S. Geological Survey, 1975; and Seaber and others, 1984.)

Public Supply

The public-supply category refers to water distributed by a publicly- or privately-owned community water system to the public. A total of 2,030 community water systems in Florida met this criteria in 2000 (Kenna Study, Florida Department of Environmental Protection, written commun., 2004; and Drinking Water Quick Look Report, Florida Department of Environmental Protection, written commun., June 2000). For this report, water-use data were collected for 1,156 systems that either served 400 people or more, or withdrew 10,000 gal/d (0.01 Mgal/d) or more. Water withdrawals from the inventoried systems totaled 2,437 Mgal/d and accounted for an estimated 99.7 percent of total public-supply withdrawals in 2000. Water withdrawn by the uninventoried systems was estimated to be about 8 Mgal/d (874 systems multiplied by 0.009 Mgal/d) and is accounted for under domestic self-supplied.

Water withdrawals for public supply in Florida in 2000 totaled 2,437 Mgal/d, of which 90 percent was obtained from ground water and 10 percent from surface water (table 4). Nearly 88 percent (14.03 million) of the State's 15.98 million residents obtained their drinking water from a public-supply water system (table 4). Of those 14.03 million residents using public-supply drinking water, 91 percent (12.80 million) were supplied from ground-water sources, and the remaining 9 percent (1.23 million) received water from surface-water sources.

Florida ranked second in the Nation to California in ground-water withdrawals for public supply (Hutson and others, 2004). The Floridan aquifer system supplied nearly 53 percent of the total public-supply withdrawals (fig. 8) and served an estimated 6.9 million people. The Biscayne aquifer supplied




Figure 8. Public-supply ground-water withdrawals in Florida by principal aquifer, 2000.

32 percent of the total public-supply withdrawals and served nearly 4 million people.

The Hillsborough River and Tampa Bypass Canal in Hillsborough County supplied 34 percent of the total surface water for public supply and Deer Point Lake in Bay County supplied 19 percent. Other significant surface-water sources of public supply water include the Braden and Manatee Rivers in Manatee County, Clear Lake in Palm Beach County, Lake Washington in Brevard County, and the Peace River and Shell Creek in Charlotte and De Soto Counties. Several of the public-supply water systems that use surface water also augment their water supply with ground water or stored surface water in an aquifer (Aquifer Storage and Recovery Systems) for later use.

Public-supply withdrawals in 2000 were smallest in September and largest in May (fig. 9). Seasonal differences in residential demands are caused by variations in temperature, precipitation, and tourism. Extreme dry conditions occurred between February and June of 2000 and the result was higher water demands from public supply primarily for lawn irrigation during these months. Lawn irrigation in central and south Florida occurs throughout the entire year.

Public suppliers provide water (deliveries) for domestic (residential), commercial, industrial, public, and other uses (see Glossary). The public-use category includes firefighting, system maintenance, and water that is lost to leakage or processing (for desalination or lime-softening). Domestic water use, which includes indoor and outdoor residential uses, accounted for




Figure 9. Average daily public-supply freshwater withdrawals by month in Florida, 2000.

14 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Table 4. Public-supplied population, water use, withdrawals, transfers, and treated water in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; water values in million gallons per day; per capita in gallons per day]

County	Population		Use		Withdrawals			Transfers			Treated
	Total ¹	Public supply	Total	Per capita	Total	Ground	Surface	Imports	Exports	Losses	Water
Alachua	217,955	179,118	28.26	158	28.26	28.26	0.00	0.00	0.00	0.00	0.00
Baker	22,259	4,326	0.88	203	0.88	0.88	0.00	0.00	0.00	0.00	0.00
Bay	148,217	129,300	51.17	396	51.17	6.28	44.89	0.00	0.00	0.00	0.00
Bradford	26,088	8,338	1.38	166	1.38	1.38	0.00	0.00	0.00	0.00	0.00
Brevard	476,230	458,282	53.35	116	27.74	13.66	14.08	25.61	0.00	0.00	0.00
Broward	1,623,018	1,603,081	258.06	161	258.06	258.06	0.00	0.00	0.00	0.00	0.09
Calhoun	13,017	4,224	0.75	178	0.75	0.75	0.00	0.00	0.00	0.00	0.00
Charlotte	141,627	115,564	14.21	123	7.28	3.29	3.99	7.72	0.00	0.79	3.29
Citrus	118,085	66,234	13.97	211	13.97	13.97	0.00	0.00	0.00	0.00	0.00
Clay	140,814	100,785	14.77	147	14.77	14.77	0.00	0.00	0.00	0.00	0.00
Collier	251,377	226,175	52.40	232	52.40	47.17	5.23	0.00	0.00	0.00	22.46
Columbia	56,513	21,235	3.67	173	3.67	3.67	0.00	0.00	0.00	0.00	0.00
DeSoto	32,209	11,233	1.90	169	10.59	4.49	6.10	0.00	8.63	0.06	0.00
Dixie	13,827	4,622	0.67	145	0.67	0.67	0.00	0.00	0.00	0.00	0.00
Duval	778,879	736,838	119.12	162	119.12	119.12	0.00	0.00	0.00	0.00	0.00
Escambia	294,410	279,294	43.56	156	44.63	44.63	0.00	0.00	1.07	0.00	0.00
Flagler	49,832	43,953	6.22	142	6.22	6.22	0.00	0.00	0.00	0.00	0.00
Franklin	11,057	9,258	1.92	207	1.92	1.92	0.00	0.00	0.00	0.00	0.00
Gadsden	45,087	27,632	4.34	157	4.34	3.06	1.28	0.00	0.00	0.00	0.00
Gilchrist	14,437	1,850	0.27	146	0.27	0.27	0.00	0.00	0.00	0.00	0.00
Glades	10,576	4,782	0.55	115	0.55	0.55	0.00	0.00	0.00	0.00	0.00
Gulf	13,332	10,338	1.47	142	1.47	1.47	0.00	0.00	0.00	0.00	0.00
Hamilton	13,327	6,366	0.95	149	0.95	0.95	0.00	0.00	0.00	0.00	0.00
Hardee	26,938	13,571	1.78	131	1.78	1.78	0.00	0.00	0.00	0.00	0.00
Hendry	36,210	20,457	4.72	231	4.72	0.95	3.77	0.00	0.00	0.00	0.00
Hernando	130,802	116,025	20.27	175	20.27	20.26	0.01	0.00	0.00	0.00	0.00
Highlands	87,366	69,820	9.14	131	9.14	9.14	0.00	0.00	0.00	0.00	0.00
Hillsborough	998,948	854,750	155.07	181	166.39	85.51	80.88	0.02	11.30	0.04	0.00
Holmes	18,564	5,860	1.38	235	1.38	1.38	0.00	0.00	0.00	0.00	0.00
Indian River	112,947	95,337	13.93	146	13.93	13.93	0.00	0.00	0.00	0.00	6.23
Jackson	46,755	16,348	2.46	150	2.46	2.46	0.00	0.00	0.00	0.00	0.00
Jefferson	12,902	5,010	0.72	144	0.72	0.72	0.00	0.00	0.00	0.00	0.00
Lafayette	7,022	1,264	0.20	158	0.20	0.20	0.00	0.00	0.00	0.00	0.00
Lake	210,528	171,137	39.92	233	39.92	39.92	0.00	0.00	0.00	0.00	0.00
Lee	440,888	357,289	52.37	147	52.37	49.09	3.28	0.00	0.00	0.00	24.37
Leon	239,452	198,937	35.70	179	35.70	35.70	0.00	0.00	0.00	0.00	0.00
Levy	34,450	11,066	2.16	195	2.16	2.16	0.00	0.00	0.00	0.00	0.00
Liberty	7,021	2,764	0.39	141	0.39	0.39	0.00	0.00	0.00	0.00	0.00
Madison	18,733	7,166	1.65	230	1.65	1.65	0.00	0.00	0.00	0.00	0.00
Manatee	264,002	250,270	39.81	159	49.92	13.87	36.05	0.00	9.71	0.40	0.00
Marion	258,916	136,842	27.99	205	27.99	27.99	0.00	0.00	0.00	0.00	0.00
Martin	126,731	87,100	18.45	212	18.45	18.45	0.00	0.00	0.00	0.00	2.82
Miami-Dade	2,253,362	2,207,800	377.27	171	394.29	394.29	0.00	0.00	17.02	0.00	0.00
Monroe	79,589	78,885	17.02	216	0.00	0.00	0.00	17.02	0.00	0.00	0.00
Nassau	57,663	24,875	6.81	274	6.81	6.81	0.00	0.00	0.00	0.00	0.00
Okaloosa	170,498	158,504	22.97	145	22.97	22.97	0.00	0.00	0.00	0.00	0.00
Okeechobee	35,910	21,600	2.23	103	2.23	0.54	1.69	0.00	0.00	0.00	0.00
Orange	896,344	813,152	186.15	229	211.76	211.76	0.00	0.00	25.61	0.00	0.00
Osceola	172,493	128,932	30.00	233	30.00	30.00	0.00	0.00	0.00	0.00	0.00
Palm Beach	1,131,184	1,035,732	229.84	222	229.84	194.57	35.27	0.00	0.00	0.00	18.27
Pasco	344,765	275,800	35.23	128	102.67	102.67	0.00	0.00	67.44	0.00	0.00
Pinellas	921,482	914,110	116.02	127	39.88	39.88	0.00	78.74	0.00	2.60	0.00
Polk	483,924	396,543	75.49	190	75.49	75.43	0.06	0.00	0.00	0.00	0.00
Putnam	70,423	23,311	3.20	137	3.20	3.20	0.00	0.00	0.00	0.00	0.00
St. Johns	123,135	105,104	16.49	157	16.49	16.49	0.00	0.00	0.00	0.00	0.00
St. Lucie	192,695	122,960	17.95	146	17.95	17.95	0.00	0.00	0.00	0.00	5.65
Santa Rosa	117,743	110,108	14.54	132	13.47	13.47	0.00	1.07	0.00	0.00	0.00
Sarasota	325,957	277,065	36.09	130	28.71	27.86	0.85	10.62	0.00	3.24	12.11
Seminole	365,196	339,403	66.90	197	66.90	66.90	0.00	0.00	0.00	0.00	0.00
Sumter	53,345	28,243	4.44	157	4.44	4.44	0.00	0.00	0.00	0.00	0.00
Suwannee	34,844	9,393	1.40	149	1.40	1.40	0.00	0.00	0.00	0.00	0.00
Taylor	19,256	10,289	1.73	168	1.73	1.73	0.00	0.00	0.00	0.00	0.00
Union	13,442	3,155	0.36	114	0.36	0.36	0.00	0.00	0.00	0.00	0.00
Volusia	443,343	414,851	54.90	132	54.90	54.90	0.00	0.00	0.00	0.00	0.00
Wakulla	22,863	9,285	2.19	236	2.19	2.19	0.00	0.00	0.00	0.00	0.00
Walton	40,601	39,024	7.35	188	7.35	7.35	0.00	0.00	0.00	0.00	0.00
Washington	20,973	7,565	1.16	153	1.16	1.16	0.00	0.00	0.00	0.00	0.00
State totals	15,982,378	14,029,530	2,429.68	174	2,436.79	2,199.36	237.43	140.80	140.78	7.11	95.29

¹University of Florida, 2001.

61 percent of the public-supply withdrawals (fig. 10 and table 5) in 2000. Domestic use was derived from the residual of the total public-supply net water use in each county minus the commercial, industrial, public, and other uses. Commercial (19 percent) and industrial (4 percent) water deliveries were estimated by multiplying county employment totals (U.S. Census Bureau, 2002) by a water-use coefficient based on average water use per employee (Davis and others, 1988) for various commercial and industrial employment sectors (Bucca and Marella, 1992). Deliveries to commercial and industrial users increased from 162 Mgal/d (Pride, 1973) in 1970, to 558 Mgal/d in 2000 (table 5), or more than 240 percent. The percentage of public-supplied water provided for public uses (including losses) in the southeastern United States (14 percent) and the percent of water provided for other uses (irrigation and power generation) in the southeastern United States (1 percent) were obtained from the American Water Works Association (1992). Some specific data for water losses were provided directly from the individual users, estimated from data provided by the water management districts, or obtained from the FDEP's monthly operating reports. Estimates of the use or deliveries of public-supply water for each county in Florida are presented in table 5.

The statewide public-supply per capita use for Florida was 174 gal/d. This value is the total public-supply water withdrawn (2,437 Mgal/d) divided by the total population served by public supply (14.03 million). Per capita use computed in this manner includes water delivered for residential, commercial, industrial, public and other uses. The public-supply per capita for 2000 is higher than previous years with the exception of 1980 (fig. 11). Both 1980 and 2000 are years that had dry conditions statewide that affected public-supply water demands primarily for residential lawn irrigation. The statewide public-supply per capita use was slightly lower than the national average for 2000 of 180 gal/d (Hutson and others, 2004). The national average is heavily influenced by the arid western states.




Figure 10. Public-supply water use (deliveries) in Florida, 2000.

Florida's average for domestic (residential) per capita use of 106 gal/d (table 5) in 2000 is slightly higher than the 103 gal/d in 1995 (Marella, 1999). The per capita use for domestic (residential) deliveries in Florida has been decreasing since the average of 144 gal/d in 1980 (Solley and others, 1983), primarily as a result of (1) conservation efforts and the use of more efficient water fixtures; (2) the use of reclaimed wastewater as a source of water for lawn irrigation; (3) the use of xeriscape landscaping; and (4) a change in the way domestic water use was derived between 1980 and 2000. The increase between 1995 and 2000 probably can be attributed to the dry conditions during 2000. It is estimated that between 25 and 75 percent of the domestic use in Florida is for outdoor purposes during certain times of the year.

The largest water withdrawals for public supply in 2000 were in Miami-Dade and Broward Counties (table 4). Miami-Dade Water and Sewer is the single largest water supplier in the State followed by Tampa Bay Water. Miami-Dade Water and Sewer withdrew 346 Mgal/d of water and served 2.0 million people; Tampa Bay Water withdrew 270 Mgal/d and served 1.7 million people (Southwest Florida Water Management District, 2002). Miami-Dade Water and Sewer obtains all of its water from the Biscayne aquifer (<http://www.miamidade.gov>) through several large wellfields within the county. Tampa Bay Water obtains about 70 percent of its water from 12 Floridan aquifer system wellfields (<http://www.tampabaywater.org>) located in three counties (Hillsborough, Pasco, and Pinellas)




Figure 11. Historical public supply average daily per capita use in Florida, 1950-2000. (Modified from Marella, 1999.)

16 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Table 5. Estimated public supply water use (deliveries), and per capita use in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; water values in million gallons per day; per capita in gallons per day]

County	Population served	Public supply water use (deliveries), in million gallons per day					Total use	Per capita use, in gallons per day	
		Domestic (residential)	Commercial	Industrial	Other uses	Public uses and losses		Public supply	Domestic (residential)
Alachua	179,118	15.18	8.03	0.81	0.28	3.96	28.26	158	85
Baker	4,326	0.37	0.33	0.05	0.01	0.12	0.88	203	86
Bay	129,300	13.73	4.74	25.03	0.51	7.16	51.17	396	106
Bradford	8,338	0.76	0.34	0.08	0.01	0.19	1.38	166	91
Brevard	458,282	31.82	11.64	1.89	0.53	7.47	53.35	116	69
Broward	1,603,081	167.93	45.18	6.24	2.58	36.13	258.06	161	105
Calhoun	4,224	0.43	0.16	0.04	0.01	0.11	0.75	178	102
Charlotte	115,564	8.59	2.45	0.19	0.14	2.84	14.21	123	74
Citrus	66,234	9.62	2.03	0.22	0.14	1.96	13.97	211	145
Clay	100,785	9.86	2.38	0.31	0.15	2.07	14.77	147	98
Collier	226,175	30.33	7.61	0.84	0.52	13.10	52.40	232	134
Columbia	21,235	1.74	1.20	0.18	0.04	0.51	3.67	173	82
DeSoto	11,233	1.15	0.42	0.04	0.02	0.27	1.90	169	102
Dixie	4,622	0.40	0.12	0.05	0.01	0.09	0.67	145	87
Duval	736,838	69.88	26.13	5.24	1.19	16.68	119.12	162	95
Escambia	279,294	28.13	8.17	0.72	0.44	6.10	43.56	156	101
Flagler	43,953	4.25	0.81	0.23	0.06	0.87	6.22	142	97
Franklin	9,258	1.40	0.21	0.02	0.02	0.27	1.92	207	151
Gadsden	27,632	2.78	0.74	0.17	0.04	0.61	4.34	157	101
Gilchrist	1,850	0.11	0.10	0.02	0.00	0.04	0.27	146	59
Glades	4,782	0.33	0.10	0.03	0.01	0.08	0.55	115	69
Gulf	10,338	1.05	0.17	0.03	0.01	0.21	1.47	142	102
Hamilton	6,366	0.55	0.25	0.01	0.01	0.13	0.95	149	86
Hardee	13,571	1.19	0.29	0.03	0.02	0.25	1.78	131	88
Hendry	20,457	2.06	0.40	1.55	0.05	0.66	4.72	231	101
Hernando	116,025	15.26	1.79	0.18	0.20	2.84	20.27	175	132
Highlands	69,820	5.04	2.59	0.14	0.09	1.28	9.14	131	72
Hillsborough	854,750	89.64	36.11	6.06	1.55	21.71	155.07	181	105
Holmes	5,860	0.80	0.33	0.05	0.01	0.19	1.38	235	137
Indian River	95,337	6.92	2.93	0.46	0.14	3.48	13.93	146	73
Jackson	16,348	1.10	0.87	0.13	0.02	0.34	2.46	150	67
Jefferson	5,010	0.43	0.16	0.02	0.01	0.10	0.72	144	86
Lafayette	1,264	0.09	0.07	0.01	0.00	0.03	0.20	158	71
Lake	171,137	29.36	3.96	0.62	0.39	5.59	39.92	233	172
Lee	357,289	26.04	11.27	1.45	0.52	13.09	52.37	147	73
Leon	198,937	20.34	9.36	0.64	0.36	5.00	35.70	179	102
Levy	11,066	1.35	0.40	0.09	0.02	0.30	2.16	195	122
Liberty	2,764	0.27	0.06	0.01	0.00	0.05	0.39	141	98
Madison	7,166	0.66	0.35	0.39	0.02	0.23	1.65	230	92
Manatee	250,270	27.40	5.39	1.05	0.40	5.57	39.81	159	109
Marion	136,842	17.25	5.03	1.52	0.27	3.92	27.99	205	126
Martin	87,100	10.77	3.69	0.49	0.18	3.32	18.45	212	124
Miami-Dade	2,207,800	245.18	66.16	9.34	3.77	52.82	377.27	171	111
Monroe	78,885	10.62	3.67	0.18	0.17	2.38	17.02	216	135
Nassau	24,875	4.03	1.39	0.37	0.07	0.95	6.81	274	162
Okaloosa	158,504	14.44	4.55	0.53	0.23	3.22	22.97	145	91
Okeechobee	21,600	1.32	0.54	0.04	0.02	0.31	2.23	103	61
Orange	813,152	103.95	48.53	5.75	1.86	26.06	186.15	229	128
Osceola	128,932	20.01	5.15	0.34	0.30	4.20	30.00	233	155
Palm Beach	1,035,732	151.64	34.78	4.35	2.30	36.77	229.84	222	146
Pasco	275,800	24.09	5.25	0.61	0.35	4.93	35.23	128	87
Pinellas	914,110	61.45	31.22	5.95	1.16	16.24	116.02	127	67
Polk	396,543	50.28	11.24	2.65	0.75	10.57	75.49	190	127
Putnam	23,311	1.63	0.93	0.16	0.03	0.45	3.20	137	70
St. Johns	105,104	10.07	3.56	0.39	0.17	2.30	16.49	157	96
St. Lucie	122,960	10.29	3.24	0.47	0.18	3.77	17.95	146	84
Santa Rosa	110,108	10.77	1.41	0.18	0.15	2.03	14.54	132	98
Sarasota	277,065	14.95	11.44	1.40	0.36	7.94	36.09	130	54
Seminole	339,403	46.38	9.09	1.39	0.67	9.37	66.90	197	137
Sumter	28,243	3.22	0.42	0.14	0.04	0.62	4.44	157	114
Suwannee	9,393	0.68	0.43	0.08	0.01	0.20	1.40	149	72
Taylor	10,289	0.98	0.36	0.13	0.02	0.24	1.73	168	95
Union	3,155	0.17	0.12	0.02	0.00	0.05	0.36	114	54
Volusia	414,851	34.21	10.90	1.55	0.55	7.69	54.90	132	82
Wakulla	9,285	1.59	0.25	0.02	0.02	0.31	2.19	236	171
Walton	39,024	5.20	1.00	0.05	0.07	1.03	7.35	188	133
Washington	7,565	0.54	0.37	0.08	0.01	0.16	1.16	153	71
State totals	14,029,530	1,484.05	464.36	93.50	24.24	363.53	2,429.68¹	174	106

¹Does not include 7.11 million gallons per day of transfer losses as reported in table 4.

and the remaining 30 percent from surface water sources in Hillsborough County. Other counties with significant withdrawals (more than 100 Mgal/d) of public-supply water are Palm Beach, Orange, Hillsborough, Duval, and Pasco (table 4). Several counties rely on water imported from other counties. Most of these transfers occur in Hillsborough, Pasco, and Pinellas Counties as Tampa Bay Water moves water from wellfields to suppliers (table 4). Miami-Dade County's water withdrawals also included water supplied (exported) to neighboring Monroe County (17 Mgal/d) for public supply throughout the Florida Keys.

Water withdrawals for public supply in Florida have increased steadily since water-use data were first collected. Total public-supply withdrawals increased 1,330 percent between 1950 and 2000 (fig. 12). Over this same period, the population of Florida increased approximately 475 percent, from 2.77 million in 1950 (Dietrich, 1978) to 15.98 million in 2000 (University of Florida, 2001). In addition to the increase in the State's population, the percentage of the population that relies on public supply for drinking water increased. In 1950, about 60 percent of the State's population was served by a public supplier; this percentage increased to 86 percent in 1985 (Marella, 1988), and has increased to 88 percent in 2000. Public-supply withdrawals increased 175 percent between 1970 and 2000, and 17 percent between 1995 and 2000. A significant part of the recent increase is a result of higher demands primarily for lawn watering caused by dry conditions during the Spring of 2000.




Figure 12. Historical public-supply freshwater withdrawals in Florida by source, 1950-2000. (Modified from Marella, 1999.)

The use of nonpotable water (treated water) for public supply has increased steadily in Florida since 1985 (Dykes and Conlon, 1989; Marella, 1995). In 2000, about 95 Mgal/d of nonpotable ground water was treated using desalination techniques or diluted with freshwater to meet the FDEP drinking-water standards for potable water (see glossary). The use of desalination or dilution of ground water for public supply was documented in nine counties (Broward, Charlotte, Collier, Indian River, Lee, Martin, Palm Beach, St. Lucie, and Sarasota) (table 4). This nonpotable water (sometimes referred to as brackish water) is included in the freshwater category for this report, thus water withdrawn for public supply in 2000 was freshwater. Future use of treated water will increase because several large desalination plants are currently being constructed that will utilize surface water.

Domestic Self-Supplied

The domestic self-supplied category comprises users that withdraw small quantities of potable water without need of a permit from the water management districts. Domestic self-supplied use includes water withdrawn by individual households (domestic wells) and by the 874 small public-supply systems (with a daily average pumpage of less than 0.01 Mgal/d) not inventoried under public supply. For this report, it was assumed that all people not served by the 1,156 inventoried public suppliers are self-supplied. Of the 199 Mgal/d withdrawn for domestic self-supplied use in 2000, an estimated 96 percent (191 Mgal/d) was from domestic wells and 4 percent (8 Mgal/d) was from the 874 small public supply systems not inventoried. Withdrawals of more than 10 Mgal/d for domestic self-supplied use in 2000 occurred in Marion, Polk, and Palm Beach Counties (table 6). It is assumed that water withdrawals for this category were derived exclusively from ground water because of its good quality and its widespread availability. About 62 percent of the domestic self-supplied water was obtained from the Floridan aquifer system (fig. 13); the remaining 38 percent was obtained from local water table or shallow aquifers (Biscayne, intermediate, sand-and-gravel, and the unnamed surficial aquifers). In many areas of Florida, the shallow or water table aquifers yield sufficient water for domestic purposes, especially where the Floridan aquifer system is relatively deep or has poor water quality.

In 2000, an estimated 1.95 million people in Florida were included in the domestic self-supplied category (table 6). According to the U.S. Census Bureau (1993), 795,560 households in Florida used an individual well as their primary source of drinking water in 1990, however the Census did not compile this data for 2000. Based on a statewide population growth rate of 23.5 percent between 1990 and 2000 (University of Florida, 2001), the number of households in 2000 with domestic wells would be nearly 982,500 (table 6). Estimated water used per well would be slightly less than 200 gal/d in 2000. Many additional households in Florida have individual wells or pump from canals for irrigation purposes only, and these withdrawals are not included in this category.

18 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Table 6. Domestic self-supplied population and water withdrawals in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee]

County	Population			People per household ²	Households with wells	Withdrawals, in million gallons per day		
	Total ¹	Public supply	Self-supplied			Total	Ground	Surface
Alachua	217,955	179,118	38,837	2.34	18,503	4.12	4.12	0.00
Baker	22,259	4,326	17,933	2.86	5,472	1.90	1.90	0.00
Bay	148,217	129,300	18,917	2.43	14,599	2.01	2.01	0.00
Bradford	26,088	8,338	17,750	2.58	6,321	1.89	1.89	0.00
Brevard	476,230	458,282	17,948	2.35	22,458	1.90	1.90	0.00
Broward	1,623,018	1,603,081	19,937	2.45	7,778	2.11	2.11	0.00
Calhoun	13,017	4,224	8,793	2.53	3,465	0.93	0.93	0.00
Charlotte	141,627	115,564	26,063	2.18	6,670	3.55	3.55	0.00
Citrus	118,085	66,234	51,851	2.20	33,424	7.20	7.20	0.00
Clay	140,814	100,785	40,029	2.77	19,055	4.24	4.24	0.00
Collier	251,377	226,175	25,202	2.39	11,382	2.67	2.67	0.00
Columbia	56,513	21,235	35,278	2.56	12,812	3.74	3.74	0.00
DeSoto	32,209	11,233	20,976	2.70	7,235	2.16	2.16	0.00
Dixie	13,827	4,622	9,205	2.44	4,881	0.98	0.98	0.00
Duval	778,879	736,838	42,041	2.51	45,073	4.46	4.46	0.00
Escambia	294,410	279,294	15,116	2.45	1,460	1.60	1.60	0.00
Flagler	49,832	43,953	5,879	2.32	2,606	0.62	0.62	0.00
Franklin	11,057	9,258	1,799	2.28	1,647	0.19	0.19	0.00
Gadsden	45,087	27,632	17,455	2.69	6,226	1.85	1.85	0.00
Gilchrist	14,437	1,850	12,587	2.61	4,200	1.33	1.33	0.00
Glades	10,576	4,782	5,794	2.51	1,252	0.61	0.61	0.00
Gulf	13,332	10,338	2,994	2.42	2,927	0.32	0.32	0.00
Hamilton	13,327	6,366	6,961	2.60	2,566	0.74	0.74	0.00
Hardee	26,938	13,571	13,367	3.06	5,394	0.64	0.64	0.00
Hendry	36,210	20,457	15,753	3.09	3,717	1.67	1.67	0.00
Hernando	130,802	116,025	14,777	2.32	13,499	1.41	1.41	0.00
Highlands	87,366	69,820	17,546	2.30	14,284	1.68	1.68	0.00
Hillsborough	998,948	854,750	144,198	2.51	70,947	4.71	4.71	0.00
Holmes	18,564	5,860	12,704	2.43	5,528	1.35	1.35	0.00
Indian River	112,947	95,337	17,610	2.25	18,773	1.87	1.87	0.00
Jackson	46,755	16,348	30,407	2.44	12,129	3.22	3.22	0.00
Jefferson	12,902	5,010	7,892	2.53	3,722	0.84	0.84	0.00
Lafayette	7,022	1,264	5,758	2.66	2,228	0.61	0.61	0.00
Lake	210,528	171,137	39,391	2.34	29,839	4.27	4.27	0.00
Lee	440,888	357,289	83,599	2.31	29,384	8.86	8.86	0.00
Leon	239,452	198,937	40,515	2.34	12,576	4.29	4.29	0.00
Levy	34,450	11,066	23,384	2.44	9,829	3.95	3.95	0.00
Liberty	7,021	2,764	4,257	2.51	1,657	0.45	0.45	0.00
Madison	18,733	7,166	11,567	2.57	4,593	1.23	1.23	0.00
Manatee	264,002	250,270	13,732	2.29	7,388	0.17	0.17	0.00
Marion	258,916	136,842	122,074	2.36	59,723	16.42	16.42	0.00
Martin	126,731	87,100	39,631	2.23	19,247	4.20	4.20	0.00
Miami-Dade	2,253,362	2,207,800	45,562	2.84	19,193	4.83	4.83	0.00
Monroe	79,589	78,885	704	2.23	1,255	0.08	0.08	0.00
Nassau	57,663	24,875	32,788	2.59	13,059	3.48	3.48	0.00
Okaloosa	170,498	158,504	11,994	2.49	3,360	1.27	1.27	0.00
Okeechobee	35,910	21,600	14,310	2.69	6,915	1.52	1.52	0.00
Orange	896,344	813,152	83,192	2.61	26,115	8.82	8.82	0.00
Osceola	172,493	128,932	43,561	2.79	13,375	4.61	4.61	0.00
Palm Beach	1,131,184	1,035,732	95,452	2.34	43,021	10.12	10.12	0.00
Pasco	344,765	275,800	68,965	2.30	43,376	4.50	4.50	0.00
Pinellas	921,482	914,110	7,372	2.17	4,385	0.41	0.41	0.00
Polk	483,924	396,543	87,381	2.52	45,174	12.45	12.45	0.00
Putnam	70,423	23,311	47,112	2.48	27,959	4.99	4.99	0.00
St. Johns	123,135	105,104	18,031	2.44	14,286	1.91	1.91	0.00
St. Lucie	192,695	122,960	69,735	2.47	29,577	7.39	7.39	0.00
Santa Rosa	117,743	110,108	7,635	2.63	3,062	0.81	0.81	0.00
Sarasota	325,957	277,065	48,892	2.13	28,710	0.43	0.43	0.00
Seminole	365,196	339,403	25,793	2.59	14,963	2.73	2.73	0.00
Sumter	53,345	28,243	25,102	2.27	11,287	4.57	4.57	0.00
Suwannee	34,844	9,393	25,451	2.54	9,970	2.70	2.70	0.00
Taylor	19,256	10,289	8,967	2.51	5,150	0.95	0.95	0.00
Union	13,442	3,155	10,287	2.76	2,275	1.10	1.10	0.00
Volusia	443,343	414,851	28,492	2.32	28,813	3.02	3.02	0.00
Wakulla	22,863	9,285	13,578	2.57	4,584	1.44	1.44	0.00
Walton	40,601	39,024	1,577	2.35	9,308	0.17	0.17	0.00
Washington	20,973	7,565	13,408	2.46	6,870	1.42	1.42	0.00
State totals	15,982,378	14,029,530	1,952,848	2.46	982,511	198.68	198.68	0.00

¹ University of Florida (2001).

² Smith and Cody (2001).





Figure 13. Domestic self-supplied ground-water withdrawals in Florida by principal aquifer, 2000.

Withdrawals for domestic self-supplied use in Florida were about 200 Mgal/d in 1970 and 2000 (fig. 14). For these two years, the same methodology was used to calculate domestic self-supplied withdrawals. In 1970, a statewide domestic per capita of 120 gal/d was used to estimate water withdrawals for all counties (Pride, 1973), while for 2000, a statewide domestic per capita of 106 gal/d was used for all counties (excluding those within the Southwest Florida Water Management District). The 2000 domestic per capita was derived from public supply withdrawals minus deliveries to commercial and industrial users and system losses as shown in table 5. This differs from 1975, 1980, 1985, 1990, and 1995 when the public supply per capita for each county was used to estimate domestic self-supplied withdrawals. After 1970, the method was changed because the public supply per capita was easily calculated and



* Methodology used for 2000 to estimate withdrawals was modified from previous years


Figure 14. Historical domestic self-supplied freshwater withdrawals in Florida, 1950-2000. (Modified from Marella, 1999.)

the higher per capita helped account for small nonpublic supply users in rural areas (such as campgrounds, churches, convenience stores, restaurants, and others). However, because the water use at these unaccounted-for systems is far less than believed, the higher per capita overestimated the water withdrawals for this category. The domestic per capita includes only what is used for households, and therefore provides a better estimate of water withdrawals for this category.

Commercial-Industrial Self-Supplied

Commercial-industrial self-supplied use is water withdrawn at commercial and industrial facilities. Commercial users include government and military facilities, schools, prisons, hospitals, recreational (non-irrigation), and nonmanufacturing facilities. Industrial users include mining, processing, and manufacturing facilities. Data were obtained for those systems having withdrawals of more than 0.01 Mgal/d; water used by the smaller commercial-industrial (non-community and non-transient non-community) water systems is insignificant and therefore is not accounted for in this report. There were 248 self-supplied commercial users and 200 self-supplied industrial users (including mines) inventoried in Florida in 2000.

Total freshwater withdrawals by commercial-industrial self-supplied systems in 2000 was 563 Mgal/d, of which 76 percent was ground water and 24 percent was surface water (table 7). The Floridan aquifer system supplied 75 percent of ground water withdrawals for this category (fig. 15). Major surface-water sources for commercial-industrial self-supplied use include Etonia and Simms Creeks in Putnam County (23 percent) and the Escambia River in Escambia County (19 percent). However, a large amount of surface water used for this category was withdrawn from unnamed mining pits or ponds for dewatering and mining operations.



Note: Intermediate aquifer equaled less than 1 percent

Figure 15. Commercial-industrial self-supplied ground-water withdrawals in Florida by principal aquifer, 2000.

20 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Table 7. Commercial-industrial self-supplied water withdrawals in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day; includes mining withdrawals]

County	Ground water			Surface water			Total water		
	Fresh	Saline	Total	Fresh	Saline	Total	Fresh	Saline	Total
Alachua	2.50	0.00	2.50	0.00	0.00	0.00	2.50	0.00	2.50
Baker	0.43	0.00	0.43	0.00	0.00	0.00	0.43	0.00	0.43
Bay	0.30	0.00	0.30	0.00	0.00	0.00	0.30	0.00	0.30
Bradford	1.25	0.00	1.25	0.00	0.00	0.00	1.25	0.00	1.25
Brevard	1.04	0.00	1.04	0.01	0.00	0.01	1.05	0.00	1.05
Broward	0.54	0.00	0.54	0.00	0.00	0.00	0.54	0.00	0.54
Calhoun	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Charlotte	0.15	0.00	0.15	2.34	0.00	2.34	2.49	0.00	2.49
Citrus	0.80	0.00	0.80	0.29	0.00	0.29	1.09	0.00	1.09
Clay	6.87	0.00	6.87	0.00	0.00	0.00	6.87	0.00	6.87
Collier	0.19	0.00	0.19	5.62	0.00	5.62	5.81	0.00	5.81
Columbia	0.34	0.00	0.34	0.00	0.00	0.00	0.34	0.00	0.34
DeSoto	0.06	0.00	0.06	1.33	0.00	1.33	1.39	0.00	1.39
Dixie	0.26	0.00	0.26	0.00	0.00	0.00	0.26	0.00	0.26
Duval	12.51	0.00	12.51	0.00	0.00	0.00	12.51	0.00	12.51
Escambia	35.77	0.00	35.77	24.65	0.00	24.65	60.42	0.00	60.42
Flagler	0.20	0.00	0.20	0.07	0.00	0.07	0.27	0.00	0.27
Franklin	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Gadsden	0.92	0.00	0.92	0.00	0.00	0.00	0.92	0.00	0.92
Gilchrist	0.26	0.00	0.26	0.00	0.00	0.00	0.26	0.00	0.26
Glades	0.14	0.00	0.14	3.93	0.00	3.93	4.07	0.00	4.07
Gulf	0.90	0.00	0.90	1.65	0.00	1.65	2.55	0.00	2.55
Hamilton	34.39	0.00	34.39	0.00	0.00	0.00	34.39	0.00	34.39
Hardee	5.93	0.00	5.93	0.00	0.00	0.00	5.93	0.00	5.93
Hendry	0.72	0.00	0.72	0.00	0.00	0.00	0.72	0.00	0.72
Hernando	19.70	0.00	19.70	0.07	0.00	0.07	19.77	0.00	19.77
Highlands	0.55	0.00	0.55	0.03	0.00	0.03	0.58	0.00	0.58
Hillsborough	14.17	0.00	14.17	5.50	0.00	5.50	19.67	0.00	19.67
Holmes	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Indian River	0.12	0.00	0.12	0.00	0.00	0.00	0.12	0.00	0.12
Jackson	1.50	0.00	1.50	0.00	0.00	0.00	1.50	0.00	1.50
Jefferson	0.20	0.00	0.20	0.00	0.00	0.00	0.20	0.00	0.20
Lafayette	0.20	0.00	0.20	0.00	0.00	0.00	0.20	0.00	0.20
Lake	10.44	0.00	10.44	0.60	0.00	0.60	11.04	0.00	11.04
Lee	8.76	0.00	8.76	7.19	0.00	7.19	15.95	0.00	15.95
Leon	0.12	0.00	0.12	0.00	0.00	0.00	0.12	0.00	0.12
Levy	0.06	0.00	0.06	1.96	0.00	1.96	2.02	0.00	2.02
Liberty	0.82	0.00	0.82	0.00	0.00	0.00	0.82	0.00	0.82
Madison	0.15	0.00	0.15	0.00	0.00	0.00	0.15	0.00	0.15
Manatee	0.47	0.00	0.47	0.53	0.00	0.53	1.00	0.00	1.00
Marion	2.08	0.00	2.08	0.00	0.00	0.00	2.08	0.00	2.08
Martin	1.20	0.00	1.20	1.95	0.00	1.95	3.15	0.00	3.15
Miami-Dade	41.65	0.00	41.65	0.00	0.00	0.00	41.65	0.00	41.65
Monroe	0.10	0.00	0.10	0.00	0.00	0.00	0.10	0.00	0.10
Nassau	32.46	0.00	32.46	0.00	1.18	1.18	32.46	1.18	33.64
Okaloosa	4.14	0.00	4.14	0.00	0.00	0.00	4.14	0.00	4.14
Okeechobee	0.36	0.00	0.36	0.00	0.00	0.00	0.36	0.00	0.36
Orange	24.21	0.00	24.21	0.00	0.00	0.00	24.21	0.00	24.21
Osceola	0.84	0.00	0.84	0.00	0.00	0.00	0.84	0.00	0.84
Palm Beach	5.76	0.00	5.76	13.21	0.00	13.21	18.97	0.00	18.97
Pasco	4.72	0.00	4.72	0.81	0.00	0.81	5.53	0.00	5.53
Pinellas	0.09	0.00	0.09	0.55	0.00	0.55	0.64	0.00	0.64
Polk	71.20	0.00	71.20	6.39	0.00	6.39	77.59	0.00	77.59
Putnam	19.14	0.00	19.14	31.12	0.00	31.12	50.26	0.00	50.26
St. Johns	0.01	0.00	0.01	0.00	0.00	0.00	0.01	0.00	0.01
St. Lucie	6.35	0.00	6.35	2.53	0.00	2.53	8.88	0.00	8.88
Santa Rosa	5.57	0.00	5.57	0.00	0.00	0.00	5.57	0.00	5.57
Sarasota	0.31	0.00	0.31	0.33	0.00	0.33	0.64	0.00	0.64
Seminole	0.08	0.00	0.08	0.00	0.00	0.00	0.08	0.00	0.08
Sumter	0.36	0.00	0.36	16.98	0.00	16.98	17.34	0.00	17.34
Suwannee	1.54	0.00	1.54	0.00	0.00	0.00	1.54	0.00	1.54
Taylor	42.15	0.00	42.15	2.96	0.00	2.96	45.11	0.00	45.11
Union	0.40	0.00	0.40	0.00	0.00	0.00	0.40	0.00	0.40
Volusia	0.49	0.00	0.49	0.00	0.00	0.00	0.49	0.00	0.49
Wakulla	0.62	0.00	0.62	0.00	0.00	0.00	0.62	0.00	0.62
Walton	0.92	0.00	0.92	0.00	0.00	0.00	0.92	0.00	0.92
Washington	0.22	0.00	0.22	0.00	0.00	0.00	0.22	0.00	0.22
State totals	430.70	0.00	430.70	132.60	1.18	133.78	563.30	1.18	564.48

Monthly withdrawals for commercial-industrial self-supplied systems fluctuated less than 200 Mgal/d during 2000 (fig. 16). Withdrawals were highest in February, May, and October and lowest in January, March, and November. Fluctuation in withdrawals occur because of increased water demand when citrus and vegetables are harvested and processed, whereas withdrawals for mining and pulp and paper usually remain fairly steady throughout the year.

Withdrawals by industrial facilities accounted for 90 percent and commercial accounted for 10 percent of the freshwater withdrawals. Mining accounted for the largest amount of water used in this category (33 percent) followed by pulp and paper manufacturing (27 percent) (fig. 17). Of the water withdrawn from the 49 inventoried mining facilities, 56 percent (103 Mgal/d) was for mining limestone and sand, 41 percent (75 Mgal/d) for phosphate mining, and 3 percent (7 Mgal/d) for mineral mining. Mining operations are located throughout Florida, but are mostly concentrated in the central part of the State. Water withdrawn in the mining industry is used primarily for material washing and conveyance, but includes water pumped to dewater the area being mined. The pulp and paper industry is located in the heavily forested areas of northern and western Florida. A small amount (8 Mgal/d) of the water withdrawn for food production and manufacturing was for the bottled water industry.

The largest amount of freshwater withdrawn for commercial-industrial self-supplied purposes was in Polk County, followed by Escambia, Putnam, Taylor, and Miami-Dade Counties (table 7). These five counties accounted for nearly one-half (49 percent) of the water withdrawn in this category during 2000.




Figure 17. Commercial-industrial self-supplied freshwater use in Florida by major industrial type, 2000.

Freshwater withdrawn for commercial-industrial self-supplied use in Florida decreased 37 percent between 1970 and 2000 (fig. 18). The increase in deliveries mentioned previously and the decrease in withdrawals indicate that most new commercial or industrial users obtained water from a public-supply water system, and many self-supplied users have either converted to public supply as their water source or have become




Figure 16. Average daily commercial-industrial self-supplied freshwater withdrawals by month in Florida, 2000.





Figure 18. Historical commercial-industrial self-supplied freshwater withdrawals in Florida by source, 1950-2000. (Modified from Marella, 1999.)

more water efficient. The conversion to public supply occurred because of the increasing cost of producing potable water and treating and discharging wastewater. Many self-supplied industries have become more water efficient in an effort to reduce discharge volumes. Between 1995 and 2000, freshwater withdrawals for this category decreased 18 percent. A large portion of this recent decrease is the result of several large industrial facilities closing in Duval and Gulf Counties.

Agricultural Self-Supplied (Irrigation and Nonirrigation)

Agricultural self-supplied use consists of water withdrawn for the irrigation of crops and for nonirrigation uses associated with farming operations. Irrigation includes the application of water on lands to assist in the growing of crops or to prevent damage to crops due to harsh climatic conditions. Nonirrigation uses include water withdrawn for livestock watering, washing dairy and farm equipment, fish farming (augmenting ponds), and other farm uses. The agricultural self-supplied category no longer includes water withdrawals for irrigation of turf grasses (golf courses and urban landscape), which is now included in the category of recreational irrigation.

Agricultural self-supplied was the largest user of freshwater in Florida, accounting for 48 percent of the State's total freshwater withdrawn in 2000. Water withdrawals totaled 3,923 Mgal/d, of which 51 percent was ground water and 49 percent was surface water (table 8). Florida ranked eleventh in the Nation for irrigation and nonirrigation withdrawals in 2000, but accounted for the largest amount of irrigation withdrawals in the eastern United States (Hutson and others, 2004).



Note: Sand-and-gravel aquifer equaled less than 1 percent

Figure 19. Agricultural self-supplied ground-water withdrawals in Florida by principal aquifer, 2000.

All water withdrawn for irrigation in Florida was assumed to be freshwater.

The Floridan aquifer system supplied 68 percent of the ground water withdrawn for agricultural self-supplied use in 2000 (fig. 19). Major sources of surface water for irrigation purposes include Lake Okeechobee and canals associated the Everglades Agricultural Area (Glades, Hendry, Martin, Palm Beach, and St. Lucie Counties), and the canals associated with the headwaters of the Upper St. Johns River (Brevard, Indian River, Okeechobee, and St. Lucie Counties). South Florida is intensively irrigated for sugarcane, citrus, and vegetables, and accounted for about 63 percent of the State's surface-water withdrawals.

Monthly withdrawals for agricultural self-supplied use had the largest seasonal variation of any water-use category. Irrigation withdrawals followed a normal seasonal pattern: they were greatest in March, April, May, and June (accounting for 54 percent of the water used) and were at a seasonal low in July, August, and September (fig. 20). A seasonal fluctuation of more than 5,000 Mgal/d in 2000 was the result of intense crop production and extreme dry conditions during the spring months, and above normal rainfall during the summer months.




Figure 20. Average daily agricultural self-supplied freshwater withdrawals by month in Florida, 2000.

Table 8. Agricultural self-supplied water withdrawals in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day]

County	Ground water			Surface water			Total water		
	Fresh	Saline	Total	Fresh	Saline	Total	Fresh	Saline	Total
Alachua	17.91	0.00	17.91	0.27	0.00	0.27	18.18	0.00	18.18
Baker	2.80	0.00	2.80	1.64	0.00	1.64	4.44	0.00	4.44
Bay	1.28	0.00	1.28	0.00	0.00	0.00	1.28	0.00	1.28
Bradford	1.00	0.00	1.00	0.02	0.00	0.02	1.02	0.00	1.02
Brevard	118.79	0.00	118.79	18.03	0.00	18.03	136.82	0.00	136.82
Broward	0.89	0.00	0.89	3.21	0.00	3.21	4.10	0.00	4.10
Calhoun	2.66	0.00	2.66	1.79	0.00	1.79	4.45	0.00	4.45
Charlotte	28.17	0.00	28.17	19.02	0.00	19.02	47.19	0.00	47.19
Citrus	2.76	0.00	2.76	0.59	0.00	0.59	3.35	0.00	3.35
Clay	5.92	0.00	5.92	0.04	0.00	0.04	5.96	0.00	5.96
Collier	137.18	0.00	137.18	5.81	0.00	5.81	142.99	0.00	142.99
Columbia	5.79	0.00	5.79	0.10	0.00	0.10	5.89	0.00	5.89
DeSoto	116.95	0.00	116.95	1.92	0.00	1.92	118.87	0.00	118.87
Dixie	1.59	0.00	1.59	0.03	0.00	0.03	1.62	0.00	1.62
Duval	3.74	0.00	3.74	0.33	0.00	0.33	4.07	0.00	4.07
Escambia	2.73	0.00	2.73	0.00	0.00	0.00	2.73	0.00	2.73
Flagler	15.67	0.00	15.67	0.03	0.00	0.03	15.70	0.00	15.70
Franklin	0.01	0.00	0.01	0.00	0.00	0.00	0.01	0.00	0.01
Gadsden	2.95	0.00	2.95	5.97	0.00	5.97	8.92	0.00	8.92
Gilchrist	13.97	0.00	13.97	0.32	0.00	0.32	14.29	0.00	14.29
Glades	19.71	0.00	19.71	49.31	0.00	49.31	69.02	0.00	69.02
Gulf	0.21	0.00	0.21	0.00	0.00	0.00	0.21	0.00	0.21
Hamilton	5.53	0.00	5.53	0.11	0.00	0.11	5.64	0.00	5.64
Hardee	81.16	0.00	81.16	0.57	0.00	0.57	81.73	0.00	81.73
Hendry	193.37	0.00	193.37	310.54	0.00	310.54	503.91	0.00	503.91
Hernando	4.26	0.00	4.26	0.03	0.00	0.03	4.29	0.00	4.29
Highlands	143.24	0.00	143.24	17.07	0.00	17.07	160.31	0.00	160.31
Hillsborough	83.84	0.00	83.84	3.31	0.00	3.31	87.15	0.00	87.15
Holmes	0.78	0.00	0.78	0.19	0.00	0.19	0.97	0.00	0.97
Indian River	64.63	0.00	64.63	157.86	0.00	157.86	222.49	0.00	222.49
Jackson	14.00	0.00	14.00	1.70	0.00	1.70	15.70	0.00	15.70
Jefferson	6.43	0.00	6.43	0.10	0.00	0.10	6.53	0.00	6.53
Lafayette	5.77	0.00	5.77	0.16	0.00	0.16	5.93	0.00	5.93
Lake	30.85	0.00	30.85	5.30	0.00	5.30	36.15	0.00	36.15
Lee	42.51	0.00	42.51	18.00	0.00	18.00	60.51	0.00	60.51
Leon	1.27	0.00	1.27	0.00	0.00	0.00	1.27	0.00	1.27
Levy	21.82	0.00	21.82	0.63	0.00	0.63	22.45	0.00	22.45
Liberty	0.16	0.00	0.16	0.00	0.00	0.00	0.16	0.00	0.16
Madison	5.83	0.00	5.83	0.11	0.00	0.11	5.94	0.00	5.94
Manatee	104.37	0.00	104.37	1.35	0.00	1.35	105.72	0.00	105.72
Marion	16.31	0.00	16.31	0.90	0.00	0.90	17.21	0.00	17.21
Martin	15.02	0.00	15.02	125.00	0.00	125.00	140.02	0.00	140.02
Miami-Dade	86.55	0.00	86.55	23.80	0.00	23.80	110.35	0.00	110.35
Monroe	0.03	0.00	0.03	0.00	0.00	0.00	0.03	0.00	0.03
Nassau	0.68	0.00	0.68	0.02	0.00	0.02	0.70	0.00	0.70
Okaloosa	0.75	0.00	0.75	0.00	0.00	0.00	0.75	0.00	0.75
Okeechobee	57.29	0.00	57.29	9.75	0.00	9.75	67.04	0.00	67.04
Orange	19.67	0.00	19.67	7.32	0.00	7.32	26.99	0.00	26.99
Osceola	80.47	0.00	80.47	25.49	0.00	25.49	105.96	0.00	105.96
Palm Beach	42.07	0.00	42.07	904.09	0.00	904.09	946.16	0.00	946.16
Pasco	22.37	0.00	22.37	0.36	0.00	0.36	22.73	0.00	22.73
Pinellas	0.43	0.00	0.43	0.03	0.00	0.03	0.46	0.00	0.46
Polk	157.53	0.00	157.53	9.33	0.00	9.33	166.86	0.00	166.86
Putnam	11.69	0.00	11.69	3.50	0.00	3.50	15.19	0.00	15.19
St. Johns	28.59	0.00	28.59	0.00	0.00	0.00	28.59	0.00	28.59
St. Lucie	45.16	0.00	45.16	192.18	0.00	192.18	237.34	0.00	237.34
Santa Rosa	5.00	0.00	5.00	0.00	0.00	0.00	5.00	0.00	5.00
Sarasota	6.88	0.00	6.88	0.66	0.00	0.66	7.54	0.00	7.54
Seminole	11.06	0.00	11.06	0.02	0.00	0.02	11.08	0.00	11.08
Sumter	14.08	0.00	14.08	0.54	0.00	0.54	14.62	0.00	14.62
Suwannee	20.64	0.00	20.64	0.38	0.00	0.38	21.02	0.00	21.02
Taylor	1.89	0.00	1.89	0.04	0.00	0.04	1.93	0.00	1.93
Union	1.05	0.00	1.05	0.02	0.00	0.02	1.07	0.00	1.07
Volusia	29.03	0.00	29.03	3.88	0.00	3.88	32.91	0.00	32.91
Wakulla	0.28	0.00	0.28	0.00	0.00	0.00	0.28	0.00	0.28
Walton	1.80	0.00	1.80	0.29	0.00	0.29	2.09	0.00	2.09
Washington	1.13	0.00	1.13	0.00	0.00	0.00	1.13	0.00	1.13
State totals	1,989.95	0.00	1,989.95	1,933.06	0.00	1,933.06	3,923.01	0.00	3,923.01

24 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

About 4.618 million acres of agricultural lands were farmed in Florida in 2000, of which 1.866 million acres (40 percent) were irrigated (table 9). Improved pasture accounted for more than one-half of the agricultural acres (54 percent); however, only 5 percent of the improved pasture was irrigated in

2000. Excluding improved pasture acreage, more than 80 percent of the acreage farmed in Florida during 2000 was irrigated. Most of the acreage was irrigated by flood systems (45 percent), followed by micro systems (38 percent), and sprinkler systems (17 percent) (table 9).

Table 9. Acres irrigated, irrigation system type, and water use in Florida by crop type, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; --- no or partial data were available]

Crop and acreage		Irrigation system type				Water use, in million gallons per day				Water withdrawn
Crop type	Total	Irrigated	Micro	Sprinkler	Flood	Ground	Surface	Reclaimed	Total	
Vegetable crops	239,674	209,925	21,025	69,951	118,949	282.35	118.69	3.30	404.34	401.04
Cabbage	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
Carrots	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
Cucumbers	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
Peppers	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
Tomatoes	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
Sweet corn	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
Melons	---	---	---	---	---	0.00	0.00	0.00	0.00	0.00
All vegetables ¹	239,674	209,925	21,025	69,951	118,949	282.35	118.69	3.30	404.34	401.04
Fruit crops	863,757	843,703	658,267	77,690	107,746	1,091.67	773.16	19.02	1,883.83	1,864.83
Blueberries	1,010	913	163	720	30	1.20	0.03	0.00	1.23	1.23
Citrus	834,802	824,602	650,726	66,548	107,328	1,056.16	769.02	19.02	1,844.18	1,825.18
Grapes	310	214	131	83	0	0.32	0.05	0.00	0.37	0.37
Peaches	5	5	5	0	0	0.01	0.00	0.00	0.01	0.01
Pecans	2,264	569	416	153	0	0.83	0.10	0.00	0.93	0.93
Strawberries	6,601	6,588	6,124	400	64	13.10	0.50	0.00	13.60	13.60
Miscellaneous	11,509	9,523	429	8,770	324	18.50	3.34	0.00	21.84	21.84
Fruit crops nonspecific	7,256	1,289	273	1,016	0	1.55	0.12	0.00	1.67	1.67
Field crops	882,313	533,683	1,031	80,648	452,004	132.93	871.91	18.08	1,022.92	1,004.84
Cotton	93,895	14,018	0	14,018	0	8.26	0.79	0.00	9.05	9.05
Field corn	89,204	28,967	72	23,484	5,411	33.87	1.87	0.63	36.37	35.74
Peanuts	84,646	25,036	0	25,036	0	20.10	1.26	0.00	21.36	21.36
Potatoes	34,132	33,196	949	2,326	29,921	25.58	6.59	0.00	32.17	32.17
Rice	12,335	12,335	0	0	12,335	0.00	33.24	0.00	33.24	33.24
Sorghum	1,658	1,480	0	1,280	200	0.91	0.13	0.00	1.04	1.04
Soybeans	39,235	2,246	0	2,246	0	0.99	0.08	0.00	1.07	1.07
Sugarcane	436,452	404,123	0	72	404,051	31.05	825.81	0.00	856.86	856.86
Tobacco	5,886	5,049	0	5,049	0	5.31	0.10	0.00	5.41	5.41
Wheat	5,757	244	0	244	0	0.29	0.03	0.00	0.32	0.32
Miscellaneous	79,113	6,989	10	6,893	86	6.57	2.01	17.45	26.03	8.58
Ornamentals and grasses	2,632,607	278,840	21,835	96,646	160,359	444.23	167.58	49.51	661.52	611.81
Field grown	25,579	24,190	4,484	13,946	5,760	54.49	11.41	2.99	68.89	65.90
Greenhouse grown	598	564	278	286	0	1.40	0.79	0.00	2.19	2.19
Container grown	38,449	36,387	17,062	19,023	302	104.28	26.37	1.48	132.13	130.65
Pasture hay (alfalfa/tame)	2,478,577	130,028	11	28,541	101,476	161.11	41.83	1.47	204.41	202.94
Sod	77,954	76,221	0	23,400	52,821	122.95	87.18	0.00	210.13	210.13
Other crops and grasses that utilize reclaimed water	11,450	11,450	0	11,450	0	0.00	0.00	43.57	43.77	0.00
Irrigation totals²	4,618,351	1,866,151	702,158	324,935	839,058	1,951.18	1,931.52	89.91	3,972.61	3,882.70
Livestock						30.96	1.51	0.00	32.47	32.47
Fish farming						7.81	0.21	0.00	8.02	8.02
Miscellaneous						0.00	0.00	0.00	0.00	0.00
Nonirrigation totals						38.77	1.72	0.00	40.49	40.49
State totals	4,618,351	1,866,151	702,158	324,935	839,058	1,989.95	1,933.06	89.91	4,013.10	3,923.01

¹Includes miscellaneous vegetables and those listed above.

²Includes vegetable crops, fruit crops, field crops, and ornamentals and grasses.

Agricultural irrigation data were compiled for 28 specific crop types in 4 main categories: vegetable, fruit, field, and ornamentals and grasses. Of the four major crop types, fruit was the largest user of water for irrigation, accounting for 47 percent of the agricultural water withdrawn (fig. 21). Citrus accounted for 98 percent of the irrigated acreage and water withdrawn for fruit crops. Total citrus acreage in 2000 was 834,800 (Florida Agricultural Statistics Service, 2000), with nearly 99 percent of the acres irrigated (table 9). Field crops, including sugarcane, was the second largest user of water, accounting for nearly 26 percent (fig. 21). Total sugarcane acreage in 2000 was 436,450 (Florida Agricultural Statistics Service, 2001a), with nearly 93 percent of the acres irrigated (table 9). Citrus and sugarcane combined accounted for 66 percent of the acres irrigated and 68 percent of the water withdrawn for agriculture self-supplied in 2000.




Figure 21. Agricultural self-supplied freshwater use in Florida by major crop type, 2000.

Palm Beach, Hendry, St. Lucie, and Indian River Counties accounted for nearly one-half (49 percent) of the total water withdrawn for agricultural self-supplied use (table 8). Palm Beach County was the largest user of surface water, and Hendry County used the most ground water. Palm Beach County had nearly 460,000 total acres irrigated, of which 330,000 were sugarcane. St. Lucie, Indian River, and Polk Counties each had more than 100,000 acres of irrigated citrus with Hendry County at just under 99,400 acres.

Water withdrawn for agricultural self-supplied use increased 88 percent between 1970 and 2000 and increased 21 percent between 1995 and 2000 (fig. 22). The long-term trend can be attributed to an increase in acres irrigated caused by the development of irrigation technology coupled with the availability of freshwater for irrigation, Florida's subtropical climate and mild winters, which allow for early spring crop production and the potential to harvest multiple crops per year,

and the State's close proximity to east coast markets and the capability to ship perishable products in a timely fashion (Marella, 1997). The more recent trend is a result of different climatic conditions between 1995 and 2000. Since irrigation is heavily dependent upon rainfall, above or below normal rainfall during peak growing seasons in Florida (usually between February and June) can dramatically change annual water demands (Marella, 1997). Severe dry conditions across most of Florida during the first part of 2000 caused water demands to be higher-than-normal especially compared to 1995, which had above-normal rainfall during the same period causing water demands to be lower-than-normal (fig. 22). A similar trend occurred between 1985 and 1990.




Figure 22. Historical agricultural self-supplied freshwater withdrawals in Florida by source, 1950-2000. (Modified from Marella, 1999.)

Total irrigated acreage increased by 17 percent between 1970 and 2000 despite crop loss to freezes, disease, insect infestation, wetland restoration, and encroaching urbanization in agricultural areas. Total citrus acreage has decreased by more than 108,000 acres between 1970 and 2000 (fig. 23); however, actual irrigated acreage increased by nearly 200,000 acres during this time. In addition, the density of citrus trees per acre has changed between 1970 and 2000 to help increase production while decreasing citrus acreage. About 80 orange trees were in production per acre in 1970 compared to 130 trees per acre in 2000 (Florida Agricultural Statistics Service, 2000). Sugar cane acreage increased by more than 250,000 acres between 1970 and 2000 (fig. 23); however, most of the increase occurred between 1970 and 1994 (Florida Department of Agriculture and Consumer Services, 1996). Harvested sugarcane acreage has remained between 400,000 and 440,000 acres since 1994 (Florida Department of Agriculture and Consumer Services, 2002).

26 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000




Figure 23. Historical agricultural acreage in Florida for selected crops, 1970-2000. (Modified from Florida Agricultural Statistics Service, 2000, 2001a, 2001b, and Marella, 1997.)

Vegetable acreage (including berries, potatoes, and watermelons) has decreased by less than 10 percent between 1970 and 2000, but has remained about 300,000 acres for this period (fig. 23). Large gains in acreage for ornamentals and grasses has occurred over the past 20 years. Floriculture crops associated with nurseries or greenhouses (flowers, foliage and woody ornamentals) increased from 35,700 acres in 1985 to 61,000 acres in 2000, and sod production increased from 32,600 acres in 1985 to 76,000 acres in 2000 (Marella, 1988). The demand for ornamentals and grass (sod) has increased because of the growth in housing and commercial establishments and the need for landscaping. Sod and many landscape plants require large amounts of water to grow and maintain.

Another significant irrigation trend in Florida is the move to more efficient irrigation systems. More than 60 percent of the acres irrigated statewide in 1980 used less-efficient flood systems, whereas only 16 percent of the acres irrigated used more-efficient micro systems. By 2000, 38 percent of the acres irrigated used micro systems compared to 45 percent using flood systems (fig 24). The increase in micro-irrigation is particularly significant for citrus, as nearly all of the newly replanted acreage is being irrigated by these more-efficient systems, and many of the older groves are being converted from flood or sprinkler systems. Many other crops are utilizing these more efficient micro-irrigation systems as farmers statewide are moving from flood or sprinkler systems to help conserve water.

In addition, the use of reclaimed wastewater, captured rainfall, and unused irrigation water (tailwater runoff) as alternative water sources has helped offset additional freshwater demands on ground- and surface-water sources. Nearly 108 Mgal/d of reclaimed wastewater was used for agricultural crop irrigation purposes in 2000 (Florida Department of Environmental Protection, 2001).




Figure 24. Historical agricultural acreage irrigated in Florida by irrigation system type, 1980-2000. (Modified from Marella, 1997 and 1999.)

Recreational Irrigation

Recreational irrigation includes the application of water on lands to assist in the growing of turf grass and landscape vegetation for lawns or recreation purposes, and also includes water used for aesthetic purposes. Turf grass recreation includes the irrigation of golf courses (including all grass and landscape associated with golf courses); turf grass lawns relates to the irrigation of all grass and landscape associated with athletic fields, cemeteries, common public or highway areas, parks, playgrounds, and lawns (primarily nonresidential, but does include some residential). Aesthetic uses include water used to fill or maintain nonagricultural ponds. Recreational irrigation is a relatively new category beginning with the 1995 compilation; however, these data have been collected since 1985 and were included under the agricultural irrigation category for 1985 and 1990. Water used for recreational irrigation may be obtained from a public water supplier, reclaimed wastewater, or self-supplied. Acreage and water-use values reported herein are for reclaimed wastewater or self-supplied withdrawals only.

Approximately 412 Mgal/d of freshwater was withdrawn (table 10) for recreational irrigation purposes in 2000, with use of an additional 227 Mgal/d obtained from reclaimed wastewater (Florida Department of Environmental Protection, 2001). Nearly 56 percent was from ground water and the remaining 44 percent was from surface water. The Floridan aquifer system supplied 62 percent and the surficial aquifer supplied 18 percent of the ground-water withdrawals for this purpose (fig. 25). Ponds, lakes and canals are the major source of surface water for irrigation purposes, and for most golf courses are usually on-site and designed to catch unused irrigation water and rainfall runoff, as well as to provide aesthetic value for the golf course. Often, these ponds or lakes are augmented with ground water or reclaimed wastewater to maintain water levels and provide storage for irrigation supplies. About 108 Mgal/d of reclaimed wastewater was used in 2000 for golf course irrigation directly or indirectly by augmenting irrigation ponds.

Table 10. Recreational irrigation water withdrawals in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee: all values in million gallons per day]

County	Ground water			Surface water			Total water		
	Fresh	Saline	Total	Fresh	Saline	Total	Fresh	Saline	Total
Alachua	4.16	0.00	4.16	0.30	0.00	0.30	4.46	0.00	4.46
Baker	0.14	0.00	0.14	0.09	0.00	0.09	0.23	0.00	0.23
Bay	1.46	0.00	1.46	0.79	0.00	0.79	2.25	0.00	2.25
Bradford	0.27	0.00	0.27	0.04	0.00	0.04	0.31	0.00	0.31
Brevard	7.75	0.00	7.75	7.08	0.00	7.08	14.83	0.00	14.83
Broward	11.86	0.00	11.86	25.14	0.00	25.14	37.00	0.00	37.00
Calhoun	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Charlotte	0.79	0.00	0.79	2.69	0.00	2.69	3.48	0.00	3.48
Citrus	3.77	0.00	3.77	0.41	0.00	0.41	4.18	0.00	4.18
Clay	1.77	0.00	1.77	0.48	0.00	0.48	2.25	0.00	2.25
Collier	15.28	0.00	15.28	10.89	0.00	10.89	26.17	0.00	26.17
Columbia	0.34	0.00	0.34	0.11	0.00	0.11	0.45	0.00	0.45
DeSoto	0.21	0.00	0.21	0.16	0.00	0.16	0.37	0.00	0.37
Dixie	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Duval	6.17	0.00	6.17	1.32	0.00	1.32	7.49	0.00	7.49
Escambia	4.79	0.00	4.79	0.77	0.00	0.77	5.56	0.00	5.56
Flagler	1.84	0.00	1.84	3.50	0.00	3.50	5.34	0.00	5.34
Franklin	0.13	0.00	0.13	0.00	0.00	0.00	0.13	0.00	0.13
Gadsden	0.15	0.00	0.15	0.15	0.00	0.15	0.30	0.00	0.30
Gilchrist	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Glades	0.02	0.00	0.02	0.40	0.00	0.40	0.42	0.00	0.42
Gulf	0.20	0.00	0.20	0.00	0.00	0.00	0.20	0.00	0.20
Hamilton	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hardee	0.21	0.00	0.21	0.04	0.00	0.04	0.25	0.00	0.25
Hendry	0.41	0.00	0.41	0.68	0.00	0.68	1.09	0.00	1.09
Hernando	3.83	0.00	3.83	0.88	0.00	0.88	4.71	0.00	4.71
Highlands	2.60	0.00	2.60	0.16	0.00	0.16	2.76	0.00	2.76
Hillsborough	9.21	0.00	9.21	4.59	0.00	4.59	13.80	0.00	13.80
Holmes	0.26	0.00	0.26	0.13	0.00	0.13	0.39	0.00	0.39
Indian River	6.81	0.00	6.81	3.22	0.00	3.22	10.03	0.00	10.03
Jackson	0.39	0.00	0.39	0.00	0.00	0.00	0.39	0.00	0.39
Jefferson	0.20	0.00	0.20	0.07	0.00	0.07	0.27	0.00	0.27
Lafayette	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Lake	5.36	0.00	5.36	3.87	0.00	3.87	9.23	0.00	9.23
Lee	11.36	0.00	11.36	11.30	0.00	11.30	22.66	0.00	22.66
Leon	1.24	0.00	1.24	0.30	0.00	0.30	1.54	0.00	1.54
Levy	0.45	0.00	0.45	0.04	0.00	0.04	0.49	0.00	0.49
Liberty	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Madison	0.20	0.00	0.20	0.06	0.00	0.06	0.26	0.00	0.26
Manatee	3.42	0.00	3.42	1.62	0.00	1.62	5.04	0.00	5.04
Marion	4.88	0.00	4.88	1.20	0.00	1.20	6.08	0.00	6.08
Martin	4.19	0.00	4.19	3.69	0.00	3.69	7.88	0.00	7.88
Miami-Dade	8.51	0.00	8.51	4.88	0.00	4.88	13.39	0.00	13.39
Monroe	1.23	0.00	1.23	0.62	0.00	0.62	1.85	0.00	1.85
Nassau	3.10	0.00	3.10	0.46	0.00	0.46	3.56	0.00	3.56
Okaloosa	3.02	0.00	3.02	0.46	0.00	0.46	3.48	0.00	3.48
Okeechobee	0.09	0.00	0.09	0.59	0.00	0.59	0.68	0.00	0.68
Orange	16.14	0.00	16.14	9.40	0.00	9.40	25.54	0.00	25.54
Osceola	1.57	0.00	1.57	0.54	0.00	0.54	2.11	0.00	2.11
Palm Beach	21.18	0.00	21.18	52.91	0.00	52.91	74.09	0.00	74.09
Pasco	5.29	0.00	5.29	1.07	0.00	1.07	6.36	0.00	6.36
Pinellas	2.39	0.00	2.39	3.00	0.00	3.00	5.39	0.00	5.39
Polk	9.65	0.00	9.65	2.29	0.00	2.29	11.94	0.00	11.94
Putnam	0.64	0.00	0.64	0.40	0.00	0.40	1.04	0.00	1.04
St. Johns	5.55	0.00	5.55	3.16	0.00	3.16	8.71	0.00	8.71
St. Lucie	3.99	0.00	3.99	4.21	0.00	4.21	8.20	0.00	8.20
Santa Rosa	6.00	0.00	6.00	0.35	0.00	0.35	6.35	0.00	6.35
Sarasota	4.89	0.00	4.89	4.07	0.00	4.07	8.96	0.00	8.96
Seminole	7.70	0.00	7.70	1.76	0.00	1.76	9.46	0.00	9.46
Sumter	3.35	0.00	3.35	0.17	0.00	0.17	3.52	0.00	3.52
Suwannee	0.07	0.00	0.07	0.02	0.00	0.02	0.09	0.00	0.09
Taylor	0.07	0.00	0.07	0.02	0.00	0.02	0.09	0.00	0.09
Union	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Volusia	9.04	0.00	9.04	3.21	0.00	3.21	12.25	0.00	12.25
Wakulla	0.20	0.00	0.20	0.00	0.00	0.00	0.20	0.00	0.20
Walton	0.27	0.00	0.27	1.52	0.00	1.52	1.79	0.00	1.79
Washington	0.39	0.00	0.39	0.00	0.00	0.00	0.39	0.00	0.39
State totals	230.45	0.00	230.45	181.28	0.00	181.28	411.73	0.00	411.73




Figure 25. Recreational irrigation ground-water withdrawals in Florida by principal aquifer, 2000.

Monthly withdrawals for recreational irrigation had a large seasonal variation. Irrigation withdrawals in 2000 were greatest in February through June during drier conditions, and lowest in July, August, September, when summer rains occurred (fig. 26). Nearly 40 percent of the water withdrawn for recreational irrigation was in Palm Beach and Broward Counties (table 10). Pinellas and Palm Beach Counties were the largest user of reclaimed wastewater for recreational irrigation.

Golf course irrigation was the largest use of water in this category, accounting for 72 percent of the water withdrawn and 48 percent of the reclaimed wastewater used. Of the nearly 192,000 acres irrigated for recreational purposes, 61 percent was for golf courses and the remaining 39 percent was for other turf grass or landscape uses. In 2000, about 1,100 golf courses were located in Florida (Florida Sports Foundation, 2000), covering an estimated 134,400 acres, of which nearly 88 percent (116,600 acres) was irrigated. According to a statewide inventory of golf courses by the National Golf Foundation (2001), an estimated 5.5 acres per golf course hole in Florida was irrigated.

Water withdrawals for recreational irrigation increased 126 percent between 1985 and 2000 (fig. 27), and golf course acreage increased 58 percent. Golf course irrigation is heavily dependent upon rainfall; normal rainfall deviations during the year can dramatically change annual water demands. In addition, golf courses in central and south Florida irrigate all year, and those along the coast must deal with high evaporation rates due to the sea breeze. These all contribute to water demands and due to the dry conditions across most of Florida during the first part of 2000, water demands were higher than normal for this year. Some of this demand is offset by the use of reclaimed water as the primary water source and the use of on-site irrigation ponds that depend on runoff. However, when rainfall does not provide enough water to maintain levels in the ponds, wells are often used to augment water to maintain ponds as an irrigation source.




Figure 26. Average daily recreational irrigation freshwater withdrawals by month in Florida, 2000.




Figure 27. Historical recreational irrigation freshwater withdrawals in Florida by source, 1985-2000. (Modified from Marella, 1999.)

Power Generation

Power generation use includes water withdrawn at thermoelectric power generation facilities (fossil fuel or nuclear) and water used at hydroelectric facilities. A total of 56 active thermoelectric power generation facilities were inventoried in 2000. An additional five thermoelectric power plants were on standby use only (used only during extreme peak demands times or while primary plants are down due to maintenance). Small private secondary power generating facilities are not included in these estimates along with the two hydroelectric power generating facilities located in Florida.

Nearly 12,613 Mgal/d of water was withdrawn for power generation purposes in 2000, of which 95 percent was saline and 5 percent was freshwater (table 11). Of the total freshwater withdrawn, 96 percent was surface water and 4 percent was ground water. More than 99 percent of the total water withdrawn (fresh and saline) for power generation is used for once-through cooling, and is returned to its source immediately after use. More than 170,000 gigawatts of total gross power was generated from the 56 inventoried facilities for 2000.

The Floridan aquifer system was the source of 84 percent (28 Mgal/d) of fresh ground-water withdrawals for power generation in 2000. Most of the surface water was withdrawn from bays or rivers along the coast, and most was returned to these sources. Major surface water sources included the Caloosahatchee River (Lee County), Crystal River and the Gulf of Mexico (Citrus County), Escambia River (Escambia County), the Indian River (Brevard, Indian River, and St. Lucie Counties), the St. Johns River (Duval and Putnam Counties), and Tampa Bay (Hillsborough and Pinellas Counties). Although several of these rivers are freshwater, the point of power plant withdrawal is often tidally influenced, and the water may be fresh, brackish, or saline due to tidal flows (McPherson and Hammett, 1991). The water withdrawn and reported herein is considered saline at most of these plants unless they are permitted by the water management district as freshwater. Public-supply deliveries to thermoelectric power plants used for domestic purposes or as boiler make-up water totaled nearly 9 Mgal/d. Additionally, 13 Mgal/d of reclaimed wastewater was used directly for cooling purposes; however, a much larger amount of cooling water was obtained from lakes or ponds supplemented by reclaimed wastewater.

30 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

Table 11. Power generation water withdrawals in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day]

County	Ground water			Surface water			Total water		
	Fresh	Saline	Total	Fresh	Saline	Total	Fresh	Saline	Total
Alachua	2.63	0.00	2.63	0.00	0.00	0.00	2.63	0.00	2.63
Baker	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Bay	0.89	0.00	0.89	0.00	265.05	265.05	0.89	265.05	265.94
Bradford	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Brevard	0.39	0.00	0.39	0.00	1,079.28	1,079.28	0.39	1,079.28	1,079.67
Broward	0.00	0.00	0.00	0.00	1,512.39	1,512.39	0.00	1,512.39	1,512.39
Calhoun	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Charlotte	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Citrus	1.55	0.00	1.55	0.00	393.90	393.90	1.55	393.90	395.45
Clay	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Collier	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Columbia	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
DeSoto	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Dixie	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Duval	8.33	0.00	8.33	0.00	661.15	661.15	8.33	661.15	669.48
Escambia	2.42	0.00	2.42	199.89	0.00	199.89	202.31	0.00	202.31
Flagler	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Franklin	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Gadsden	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Gilchrist	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Glades	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Gulf	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hamilton	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hardee	0.81	0.00	0.81	0.00	0.00	0.00	0.81	0.00	0.81
Hendry	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Hernando	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Highlands	0.08	0.00	0.08	0.00	0.00	0.00	0.08	0.00	0.08
Hillsborough	0.00	0.00	0.00	0.00	3,188.00	3,188.00	0.00	3,188.00	3,188.00
Holmes	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Indian River	0.00	0.00	0.00	0.00	39.01	39.01	0.00	39.01	39.01
Jackson	0.30	0.00	0.30	89.53	0.00	89.53	89.83	0.00	89.83
Jefferson	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Lafayette	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Lake	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Lee	0.22	0.00	0.22	0.00	492.27	492.27	0.22	492.27	492.49
Leon	2.74	0.00	2.74	0.00	0.00	0.00	2.74	0.00	2.74
Levy	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Liberty	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Madison	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Manatee	0.00	0.00	0.00	25.10	0.00	25.10	25.10	0.00	25.10
Marion	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Martin	0.54	0.00	0.54	24.63	0.00	24.63	25.17	0.00	25.17
Miami-Dade	2.08	3.68	5.76	0.00	33.19	33.19	2.08	36.87	38.95
Monroe	0.00	0.14	0.14	0.00	0.00	0.00	0.00	0.14	0.14
Nassau	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Okaloosa	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Okeechobee	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Orange	0.76	0.00	0.76	0.00	0.00	0.00	0.76	0.00	0.76
Osceola	0.03	0.00	0.03	0.00	0.00	0.00	0.03	0.00	0.03
Palm Beach	0.00	0.00	0.00	0.00	433.02	433.02	0.00	433.02	433.02
Pasco	0.14	0.00	0.14	0.00	1,956.50	1,956.50	0.14	1,956.50	1,956.64
Pinellas	0.00	0.00	0.00	0.00	419.11	419.11	0.00	419.11	419.11
Polk	4.27	0.00	4.27	15.66	0.00	15.66	19.93	0.00	19.93
Putnam	0.69	0.00	0.69	13.90	0.00	13.90	14.59	0.00	14.59
St. Johns	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
St. Lucie	0.00	0.00	0.00	0.00	1,477.95	1,477.95	0.00	1,477.95	1,477.95
Santa Rosa	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sarasota	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Seminole	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sumter	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Suwannee	0.06	0.00	0.06	101.01	0.00	101.01	101.07	0.00	101.07
Taylor	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Union	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Volusia	0.54	0.00	0.54	130.63	0.00	130.63	131.17	0.00	131.17
Wakulla	0.06	0.00	0.06	28.38	0.00	28.38	28.44	0.00	28.44
Walton	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Washington	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
State totals	29.53	3.82	33.35	628.73	11,950.82	12,579.55	658.26	11,954.64	12,612.90

Monthly freshwater withdrawals for power generation fluctuated seasonally during 2000—the highest withdrawals were from May through August, as power demands increase due to hotter weather (fig. 28). The largest amount of freshwater withdrawn was in Escambia County, and the largest amount of saline water withdrawn was in Hillsborough County (table 11). Several power plants in Duval, Hillsborough, Indian River, Orange, Osceola and Polk Counties used reclaimed wastewater to augment cooling water.




Figure 28. Average daily power generation freshwater withdrawals by month in Florida, 2000.

Total water withdrawals for power generation increased steadily between 1955 and 1980, when withdrawals peaked at 15,500 Mgal/d (fig. 29). Between 1970 and 2000, saline water withdrawals for power generation increased 25 percent, whereas freshwater withdrawals decreased 57 percent. Between 1995 and 2000, saline water withdrawals increased 9 percent, whereas freshwater withdrawals increased 3 percent. The reduction in freshwater withdrawals over the past 20 years is due to the increase in water efficiency of new or modernized facilities, and in part to increased recycling of cooling water after routing the water to cooling ponds or cooling towers, or the reclassification of a plant’s water source from freshwater to saline water. Water withdrawals at many plants are now made only to augment or replenish the water in the cooling ponds. Because of these factors, water withdrawals increased only 15 percent between 1970 and 2000, whereas total gross power generated in Florida increased 200 percent. Furthermore, differences between years could have occurred due to facility downtime caused by plant maintenance or modernization. Significant




Figure 29. Historical power generation water withdrawals in Florida by source, 1950-2000. (Modified from Marella, 1999.)

downtime can substantially reduce the annual average water withdrawn at power plant, and is particularly evident when data are collected only every 5 years.

Water Management Districts

The Florida Water Resource Act of 1972 established authority for management of the State’s water resources through five water management districts that operate under the general supervision of the Florida Department of Environmental Protection (formerly the Florida Department of Natural Resources) (Fernald and Patton, 1984). These five water management districts, which encompass the entire State, are the Northwest Florida (NFWMD), the St. Johns River (SJRWMD), the South Florida (SFWMD), the Southwest Florida (SWFWMD), and the Suwannee River (SRWMD) (fig. 2).

Of the 2000 total population in Florida, 6.60 million people (41 percent) resided in the SFWMD, followed by the SWFWMD with 3.99 million people (25 percent), the SJRWMD with 3.89 million people (24 percent), the NFWMD with 1.22 million people (8 percent), and the SRWMD with 0.25 million people (2 percent) (fig. 30). The SFWMD included the largest number of residents (6.11 million people) served by public-supply water systems (fig. 30).




Figure 30. Population and population served by public supply in Florida by water management district, 2000.

The largest amount of freshwater withdrawn was from the SFWMD, which was nearly one-half (49 percent) of the State’s total freshwater withdrawn (fig. 31 and table 12). The percentage of the total freshwater withdrawn in the other water management districts were: SJRWMD, 20 percent; SWFWMD, 19 percent; NFWMD, 8 percent; and SRWMD, 4 percent. The SFWMD accounted for the largest amount of freshwater withdrawn for public supply use (46 percent), agricultural self-supplied use (63 percent), and recreational irrigation use (51 percent). The SWFWMD accounted for the largest amount of freshwater withdrawn for commercial-industrial self-supplied use (26 percent); and the NFWMD accounted for the largest amount of freshwater withdrawn for power generation use (49 percent) (table 12). The largest amount (50 percent) of saline water withdrawn was from the SWFWMD and was used for cooling purposes at several power generation facilities located along Tampa Bay or the Gulf of Mexico (fig. 32 and table 12).




Figure 31. Freshwater withdrawals in Florida by water management district, 2000.

Table 12. Water withdrawals by category in Florida by water management district, 2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day; district totals may not be identical to data reported or published by the water management districts due to differences in data-collection procedures and categories of use or revisions in reported values; general locations of water management districts are shown in figure 2]

	Fresh ground water	Fresh surface water	Total freshwater	Total saline water	Total water
Northwest Florida					
Public Supply	145.90	46.17	192.07	0.00	192.07
Domestic self-supplied	21.75	0.00	21.75	0.00	21.75
Commercial-industrial self-supplied	51.86	26.30	78.16	0.00	78.16
Agricultural self-supplied	36.32	9.94	46.26	0.00	46.26
Recreational irrigation	18.57	4.50	23.07	0.00	23.07
Power generation	6.41	317.80	324.21	265.05	589.26
Totals	280.81	404.71	685.52	265.05	950.57
St. Johns River¹					
Public Supply	558.51	14.08	572.59	0.00	572.59
Domestic self-supplied	51.70	0.00	51.70	0.00	51.70
Commercial-industrial self-supplied	90.64	31.80	122.44	1.18	123.62
Agricultural self-supplied	392.33	214.20	606.53	0.00	606.53
Recreational irrigation	72.66	31.94	104.60	0.00	104.60
Power generation	10.86	144.53	155.39	1,779.44	1,934.83
Totals	1,176.70	436.55	1,613.25	1,780.62	3,393.87
South Florida					
Public Supply	1,070.28	49.24	1,119.52	0.00	1,119.52
Domestic self-supplied	52.43	0.00	52.43	0.00	52.43
Commercial-industrial self-supplied	88.83	35.18	124.01	0.00	124.01
Agricultural self-supplied	784.37	1,682.22	2,466.59	0.00	2,466.59
Recreational irrigation	86.21	123.68	209.89	0.00	209.89
Power generation	2.95	24.63	27.58	3,952.64	3,980.22
Totals	2,085.07	1,914.95	4,000.02	3,952.64	7,952.66
Southwest Florida²					
Public Supply	408.88	127.94	536.82	0.00	536.82
Domestic self-supplied	52.74	0.00	52.74	0.00	52.74
Commercial-industrial self-supplied	117.52	34.40	151.92	0.00	151.92
Agricultural self-supplied	683.06	24.83	707.89	0.00	707.89
Recreational irrigation	51.95	20.79	72.74	0.00	72.74
Power generation	6.77	40.76	47.53	5,957.51	6,005.04
Totals	1,320.92	248.72	1,569.64	5,957.51	7,527.15
Suwannee River					
Public Supply	15.79	0.00	15.79	0.00	15.79
Domestic self-supplied	20.06	0.00	20.06	0.00	20.06
Commercial-industrial self-supplied	81.85	4.92	86.77	0.00	86.77
Agricultural self-supplied	93.87	1.87	95.74	0.00	95.74
Recreational irrigation	1.06	0.37	1.43	0.00	1.43
Power generation	2.54	101.01	103.55	0.00	103.55
Totals	215.17	108.17	323.34	0.00	323.34
State Totals	5,078.67	3,113.10	8,191.77	11,955.82	20,147.59

¹St. Johns River Water Management District, 2004.

²Southwest Florida Water Management District, 2002.

34 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000




Figure 32. Fresh and saline water withdrawals in Florida by water management district, 2000.

Freshwater withdrawals in all districts except the SFWMD have remained about the same or decreased slightly between 1975 and 2000. However, excluding power generation, water withdrawals in all five water management districts increased between 1975 and 2000 as a result of increases in irrigated acreage, population, and tourism (fig. 33). Excluding freshwater withdrawals for power generation, the SRWMD had the largest percent increase in withdrawals between 1975 and 2000 (61 percent), followed by the SFWMD (57 percent), the SJRWMD (37 percent), and the NFWMD and SWFWMD (29 percent each). The SFWMD had the largest increase in total water withdrawals (1,440 Mgal/d) between 1975 and 2000 (fig. 33). Since 1975, fresh ground-water withdrawals increased in all five water management districts (fig.34) while fresh surface-water withdrawals decreased in all but the SFWMD (fig. 35). The decrease in fresh surface water is a result of the decrease in power generation withdrawals between 1975 and 2000.




Figure 33. Historical freshwater withdrawals in Florida by water management district, 1975-2000. (Modified from Marella, 1995 and 1999.)




Figure 34. Historical fresh ground-water withdrawals in Florida by water management district, 1975-2000. (Modified from Marella, 1995 and 1999.)




Figure 35. Historical fresh surface-water withdrawals in Florida by water management district, 1975-2000. (Modified from Marella, 1995 and 1999.)

Water Consumption and Discharges

In 2000, an estimated 52 percent (table 13) of the fresh-water withdrawn in Florida was consumed (evaporated, transpired, incorporated into products or crops, or otherwise removed from the immediate water environment); the remaining 48 percent was returned to the hydrologic system as wastewater. Less than 0.1 percent (5 Mgal/d) of the saline water was consumed. The greatest consumption of freshwater in Florida was in agricultural self-supplied (irrigation and nonirrigation) and recreational irrigation, primarily due to high evapotranspiration during hot and dry periods when irrigation demands are high (fig. 36). Water consumed ranges from 80 percent for recreational irrigation to 9 percent for power generation (freshwater only) (table 13). Nearly all (99 percent) of the total water withdrawn for power generation (fresh and saline) was returned to its source. The estimated percentage of freshwater consumed varied during the past 25 years, ranging from 33 percent in 1975 and 1980 (Leach, 1978, 1983); 43 percent in 1985 (Marella, 1988); 42 percent in 1990 (Solley and others, 1993); and 39 percent in 1995 (Marella, 1999). Estimates for 2000 are higher than past years because of prolonged drought conditions throughout the State (University of Nebraska, 2004) lead to an increase in water demands for all irrigation purposes (agriculture, recreational, and residential lawns), allowing for higher evaporation over a longer period of time. All consumption values are estimated from irrigation coefficients, industry standards, or published sources.




Figure 36. Freshwater withdrawals and estimated water consumption in Florida by category, 2000.

Table 13. Estimated freshwater consumed by category in Florida, 2000
 [Freshwater consumed in million gallons per day]

	Freshwater consumed	Percent consumed
Public supply	942.9	38.7
Domestic self-supplied	76.9	38.7
Commercial-industrial self-supplied	107.6	19.1
Agricultural self-supplied	2,720.5	69.5
Recreational irrigation	328.8	80.0
Power generation	59.3	9.0
State totals	4,236.0	51.7

Wastewater discharge includes water that was released from treatment facilities, septic tanks, or runoff from agricultural or urban lands. Discharge data for domestic (municipal) and industrial facilities are usually regulated and metered, whereas records of the amount of water released from septic tanks or that runs off agricultural or urban lands are usually not available. For this report, wastewater values were reported for domestic facilities that discharged 0.10 Mgal/d or more, and estimates of water released by septic tanks were made. No industrial wastewater discharge values were collected for 2000.

Domestic Wastewater

Domestic wastewater facilities (includes municipal and privately owned) are those systems that receive or dispose of wastewater derived principally from residential dwellings, businesses or commercial establishments, institutions and some industrial facilities (Florida Department of Environmental Regulation, 1991). According to the FDEP, there were 2,789 domestic wastewater systems in operation during 2000 (Joseph Doker, Florida Department of Environmental Protection, written commun., 2004). Total discharge from the 2,789 domestic wastewater systems for 2000 was 1,515 Mgal/d. Domestic wastewater discharge data were collected for 571 systems as reported to FDEP. Discharge from the inventoried systems totaled nearly 1,495 Mgal/d and accounted for 99 percent of the total domestic wastewater discharge. Discharge from the 2,218 uninventoried systems was estimated to be 20 Mgal/d (2,218 systems multiplied by 0.009 Mgal/d) or slightly more than 1 percent of the total discharge. Of the domestic wastewater discharged from the 571 inventoried systems, 44 percent was discharged to surface water, 34 percent was discharged to the ground through land application systems (drain fields, percolation ponds, spray




Figure 37. Treated domestic wastewater discharges in Florida by primary disposal method, 2000.

fields) and reuse systems, and 22 percent was discharged to deep aquifers through injection wells (fig. 37).

The estimated population served by the inventoried systems in 2000 was 11.07 million (table 14), and the population served by the uninventoried systems was about 0.14 million, for a total of 70 percent (11.21 million) of the State's population. The remaining 30 percent of the State's population discharged domestic wastewater to septic tanks.

Table 14. Treated domestic wastewater population served, discharge, and capacity in Florida by county, 2000

[Compiled by the U.S. Geological Survey and Florida Department of Environmental Protection; discharge and capacity values in million gallons per day]


County	Population		Wastewater discharge by source				Total number of systems	Systems inventoried	
	Total	Served	Total	Ground	Injection	Surface		Number	Capacity
Alachua	217,955	151,450	17.06	2.12	7.79	7.15	36	8	22.43
Baker	22,259	4,000	0.69	0.17	0.00	0.52	6	2	1.89
Bay	148,217	86,775	33.89	0.55	0.00	33.34	18	7	64.50
Bradford	26,088	6,850	1.93	1.60	0.00	0.33	7	2	4.15
Brevard	476,230	308,500	37.63	14.58	17.40	5.65	70	21	62.90
Broward	1,623,018	1,476,300	190.28	11.47	101.24	77.57	23	17	240.88
Calhoun	13,017	2,100	0.47	0.00	0.00	0.47	1	1	0.60
Charlotte	141,627	70,700	7.46	4.46	3.00	0.00	36	10	11.82
Citrus	118,085	18,500	2.81	2.81	0.00	0.00	80	8	5.33
Clay	140,814	73,450	7.33	1.10	0.00	6.23	29	12	14.36
Collier	251,377	164,100	28.84	26.01	1.46	1.37	36	9	34.77
Columbia	56,513	17,500	2.06	2.06	0.00	0.00	24	2	3.35
DeSoto	32,209	9,600	0.98	0.29	0.00	0.69	20	3	2.70
Dixie	13,827	3,350	0.25	0.00	0.00	0.25	3	2	0.65
Duval	778,879	592,300	84.06	0.78	0.00	83.28	67	25	123.79
Escambia	294,410	160,500	19.49	3.47	0.00	16.02	13	6	35.19
Flagler	49,832	40,750	5.79	4.83	0.00	0.96	19	8	11.46
Franklin	11,057	3,100	1.07	0.43	0.00	0.64	10	4	1.58
Gadsden	45,087	14,400	2.23	0.37	0.00	1.86	13	5	3.95
Gilchrist	14,437	1,000	0.18	0.18	0.00	0.00	6	2	0.32
Glades	10,576	0	0.12	0.12	0.00	0.00	28	1	0.14
Gulf	13,332	4,650	0.36	0.16	0.00	0.20	5	3	0.62
Hamilton	13,327	5,375	0.86	0.11	0.00	0.75	14	3	1.52
Hardee	26,938	10,525	1.17	0.38	0.00	0.79	16	4	1.73
Hendry	36,210	14,500	1.50	1.50	0.00	0.00	16	4	2.33
Hernando	130,802	38,000	4.65	4.56	0.00	0.09	46	11	9.27
Highlands	87,366	23,100	2.33	2.33	0.00	0.00	73	8	4.96
Hillsborough	998,948	720,100	92.93	29.42	0.00	63.51	144	16	144.50
Holmes	18,564	2,750	0.70	0.00	0.00	0.70	4	1	1.40
Indian River	112,947	60,325	7.37	6.99	0.00	0.38	17	6	11.56
Jackson	46,755	11,350	2.77	0.75	0.00	2.02	10	5	4.84
Jefferson	12,902	2,200	0.55	0.12	0.00	0.43	5	2	1.16
Lafayette	7,022	1,000	0.23	0.23	0.00	0.00	2	2	0.36
Lake	210,528	94,000	9.78	9.78	0.00	0.00	133	20	16.00
Lee	440,888	296,050	39.19	22.93	1.25	15.01	103	32	65.29
Leon	239,452	154,725	18.29	18.29	0.00	0.00	19	4	33.26
Levy	34,449	6,300	0.47	0.47	0.00	0.00	13	3	0.87
Liberty	7,021	0	0.13	0.13	0.00	0.00	2	1	0.20
Madison	18,733	3,400	0.92	0.92	0.00	0.00	8	2	1.12
Manatee	264,002	228,075	26.27	15.51	5.56	5.20	11	5	39.10
Marion	258,916	90,000	7.79	7.79	0.00	0.00	158	16	14.81
Martin	126,731	61,075	6.86	3.66	3.20	0.00	75	13	13.68
Miami-Dade	2,253,362	1,828,500	311.10	14.21	89.82	207.07	27	5	358.70
Monroe	79,589	35,700	7.05	0.27	0.94	5.84	306	9	12.88
Nassau	57,663	21,975	3.68	1.64	0.00	2.04	27	6	5.76
Okaloosa	170,498	110,525	16.71	16.71	0.00	0.00	17	12	27.56
Okeechobee	35,910	4,200	0.74	0.74	0.00	0.00	26	3	1.40
Orange	896,344	695,875	73.15	70.89	0.00	2.26	84	21	114.44
Osceola	172,493	120,225	16.00	15.79	0.00	0.21	48	14	23.66
Palm Beach	1,131,184	945,550	108.07	25.78	57.43	24.86	71	12	160.44
Pasco	344,765	148,925	18.04	14.94	0.00	3.10	89	15	27.27
Pinellas	921,482	874,600	106.08	48.55	33.41	24.12	28	17	179.25
Polk	483,924	213,100	28.10	17.86	0.00	10.24	198	31	50.07
Putnam	70,423	10,000	2.62	0.00	0.00	2.62	33	2	3.25
St. Johns	123,135	68,900	9.01	3.16	0.00	5.85	44	14	15.89
St. Lucie	192,695	103,400	10.84	2.04	8.63	0.17	47	16	17.58
Santa Rosa	117,743	33,100	3.93	2.26	0.00	1.67	13	5	7.43
Sarasota	325,957	219,000	21.60	12.87	1.93	6.80	71	19	33.05
Seminole	365,196	278,200	48.95	36.97	0.00	11.98	29	15	77.21
Sumter	53,345	4,000	1.84	1.84	0.00	0.00	40	4	3.20
Suwannee	34,844	6,000	0.87	0.87	0.00	0.00	13	3	1.54
Taylor	19,256	4,700	0.97	0.11	0.00	0.86	7	2	1.65
Union	13,442	4,800	0.47	0.00	0.00	0.47	1	1	0.70
Volusia	443,343	284,700	31.49	16.60	0.00	14.89	126	18	56.49
Wakulla	22,863	2,100	0.29	0.29	0.00	0.00	9	2	1.00
Walton	40,601	16,000	2.77	2.77	0.00	0.00	10	6	7.19
Washington	20,973	3,200	0.75	0.14	0.00	0.61	6	3	1.57
State totals	15,982,377	11,070,000	1,494.86	510.73	333.06	651.07	2,789	571	2,204.52

The largest domestic wastewater discharges in 2000 were in Miami-Dade and Broward Counties. Miami-Dade County discharged more than 200 Mgal/d to surface water (Atlantic Ocean) and Broward County discharged more than 100 Mgal/d to injection wells (table 14). Both Miami-Dade and Broward Counties each had more than 1 million people served by domestic wastewater systems (table 14). Miami-Dade Water and Sewer in Miami-Dade County had the single largest discharge of domestic wastewater (308 Mgal/d), which was treated and discharged at 3 facilities during 2000. Orange County discharged more than 70 Mgal/d of water to the ground through land application or reuse systems (table 14).

Domestic wastewater discharge increased 33 percent between 1985 and 2000, but decreased 3 percent between 1995 and 2000 (fig. 38). The trend between 1985 and 2000 can be attributed to increases in population served by wastewater, as many municipalities expanded service areas and treatment plant capacities. The more recent trend between 1995 and 2000 can be attributed to wet conditions in 1995, increasing infiltration into wastewater return systems directly from rainfall runoff or indirectly from ground-water seepage into underground pipes, and to dry conditions in 2000 during which little infiltration or seepage occurred. Another important trend in domestic wastewater is the decrease in the amount of water discharged to surface water bodies and the increase in the amount of water discharged to the ground through land application systems (drain fields, percolation ponds, and spray fields) and irrigation reuse systems. In 1990, about 58 percent of the domestic wastewater was discharged to surface water (Marella, 1994) compared to 44 percent in 2000 (fig. 38). This is an ongoing declining trend as restrictions on the quality and quantity of water that is discharged to surface waters has increased and the discharge of wastewater through reuse or ground-water recharge is highly encouraged. The use of wastewater discharge for irrigation and ground-water recharge has increased from 200 Mgal/d in 1986 to more than 450 Mgal/d in 2000 (Florida Department of Environmental Protection, 2001). Statewide domestic wastewater discharge values were not available prior to 1985 and the quantity of water discharged to surface water, injection wells, or ground water was not determined for 1985.

Septic Tanks

Septic tanks (also referred to as on-site disposal systems) are buried watertight receptacles constructed to promote the separation of solids, grease, and liquid components of waste-water in the absence of oxygen. The liquid fraction from the septic tank is discharged to the ground through a subsurface drain field for further treatment, and then either transpires to the atmosphere through surface vegetation or percolates into the shallow water table (Marella, 1994 and



* Discharge method for 1985 was not differentiated.

Figure 38. Historical treated domestic wastewater discharges in Florida, 1985-2000 (Modified from Marella, 1994 and 1999.)

1998). Septic tanks are most commonly used for individual households or small commercial establishments (campgrounds, churches, convenience stores, restaurants, small motels, and facilities alike) that are in rural or remote areas, or are in urban areas that are not served by a domestic wastewater system. In 2000, nearly 1.95 million septic tanks were in use for the treatment and disposal of wastewater in Florida (Dale Holcomb, Florida Department of Health, written commun., 2003). Miami-Dade and Polk Counties each had over 100,000 septic tanks (table 15).

Based on 55 gal/d of discharge per person (Tchobanoglous, 1991) and 2.46 people per household in Florida (Smith and Cody, 2001), discharge from each tank would be about 135 gal/d. Estimated discharge from the 1.95 million septic tanks in Florida in 2000 would be about 263 Mgal/d, an increase of 24 percent from the 210 Mgal/d discharged in 1990 (Marella, 1994). The number of septic tanks in Florida has increased from 0.94 million in 1970, 1.17 million in 1980, and 1.56 million in 1990 (Marella, 1994). The number of people who relied on septic tanks increased to 4.77 million in 2000 (table 15).

Table 15. Treated domestic wastewater and septic tank population served and discharge in Florida by county, 2000

[Compiled by the U.S. Geological Survey, Florida Department of Environmental Protection, and the Florida Department of Health. Discharge values in million gallons per day]

County	Total systems	Domestic wastewater systems								
		Inventoried			Non-inventoried			Septic tanks		
		Systems	Discharge	Population	Systems	Discharge	Population	Number	Discharge	Population
Alachua	36	8	17.06	151,450	28	0.25	1,852	26,362	3.56	64,653
Baker	6	2	0.69	4,000	4	0.04	296	5,925	0.80	17,963
Bay	18	7	33.89	86,775	11	0.10	741	25,169	3.40	60,701
Bradford	7	2	1.93	6,850	5	0.04	296	7,502	1.01	18,942
Brevard	70	21	37.63	308,500	49	0.44	3,259	63,068	8.51	164,471
Broward	23	17	190.28	1,476,300	6	0.05	370	59,899	8.09	146,348
Calhoun	1	1	0.47	2,100	0	0.00	0	4,503	0.61	10,917
Charlotte	36	10	7.46	70,700	26	0.23	1,704	37,230	5.03	69,223
Citrus	80	8	2.81	18,500	72	0.64	4,741	46,272	6.25	94,844
Clay	29	12	7.33	73,450	17	0.15	1,111	23,731	3.20	66,253
Collier	36	9	28.84	164,100	27	0.24	1,778	28,435	3.84	85,499
Columbia	24	2	2.06	17,500	22	0.20	1,481	19,160	2.59	37,532
DeSoto	20	3	0.98	9,600	17	0.15	1,111	7,770	1.05	21,498
Dixie	3	2	0.25	3,350	1	0.01	74	7,609	1.03	10,403
Duval	67	25	84.06	592,300	42	0.37	2,741	71,258	9.62	183,838
Escambia	13	6	19.49	160,500	7	0.06	444	52,768	7.12	133,466
Flagler	19	8	5.79	40,750	11	0.10	741	3,893	0.53	8,341
Franklin	10	4	1.07	3,100	6	0.05	370	4,191	0.57	7,587
Gadsden	13	5	2.23	14,400	8	0.07	519	13,029	1.76	30,168
Gilchrist	6	2	0.18	1,000	4	0.04	296	5,629	0.76	13,141
Glades	28	1	0.12	0	27	0.24	1,778	4,780	0.65	8,798
Gulf	5	3	0.36	4,650	2	0.02	148	4,493	0.61	8,534
Hamilton	14	3	0.86	5,375	11	0.10	741	3,209	0.43	7,211
Hardee	16	4	1.17	10,525	12	0.11	815	5,363	0.72	15,598
Hendry	16	4	1.50	14,500	12	0.11	815	7,029	0.95	20,895
Hernando	46	11	4.65	38,000	35	0.31	2,296	37,752	5.10	90,506
Highlands	73	8	2.33	23,100	65	0.58	4,296	28,407	3.83	59,970
Hillsborough	144	16	92.93	720,100	128	1.14	8,444	88,675	11.97	270,404
Holmes	4	1	0.70	2,750	3	0.03	222	7,210	0.97	15,592
Indian River	17	6	7.37	60,325	11	0.10	741	26,292	3.55	51,881
Jackson	10	5	2.77	11,350	5	0.04	296	14,772	1.99	35,109
Jefferson	5	2	0.55	2,200	3	0.03	222	4,442	0.60	10,480
Lafayette	2	2	0.23	1,000	0	0.00	0	2,498	0.34	6,022
Lake	133	20	9.78	94,000	113	1.01	7,481	54,556	7.37	109,047
Lee	103	32	39.19	296,050	71	0.63	4,667	72,815	9.83	140,171
Leon	19	4	18.29	154,725	15	0.13	963	30,306	4.09	83,764
Levy	13	3	0.47	6,300	10	0.09	667	15,199	2.05	27,482
Liberty	2	1	0.13	0	1	0.01	74	2,777	0.37	6,947
Madison	8	2	0.92	3,400	6	0.05	370	6,244	0.84	14,963
Manatee	11	5	26.27	228,075	6	0.05	370	14,294	1.93	35,557
Marion	158	16	7.79	90,000	142	1.26	9,333	81,253	10.97	159,583
Martin	75	13	6.86	61,075	62	0.55	4,074	25,436	3.43	61,582
Miami-Dade	27	5	311.10	1,828,500	22	0.20	1,481	125,654	16.96	423,381
Monroe	306	9	7.05	35,700	297	2.64	19,556	25,575	3.45	24,333
Nassau	27	6	3.68	21,975	21	0.19	1,407	14,508	1.96	34,281
Okaloosa	17	12	16.71	110,525	5	0.04	296	19,373	2.62	59,677
Okeechobee	26	3	0.74	4,200	23	0.20	1,481	11,916	1.61	30,229
Orange	84	21	73.15	695,875	63	0.56	4,148	82,234	11.10	196,321
Osceola	48	14	16.00	120,225	34	0.30	2,222	20,402	2.75	50,046
Palm Beach	71	12	108.07	945,550	59	0.53	3,926	60,930	8.23	181,708
Pasco	89	15	18.04	148,925	74	0.66	4,889	64,572	8.72	190,951
Pinellas	28	17	106.08	874,600	11	0.10	741	13,820	1.87	46,141
Polk	198	31	28.10	213,100	167	1.49	11,037	100,359	13.55	259,787
Putnam	33	2	2.62	10,000	31	0.28	2,074	30,050	4.06	58,349
St. Johns	44	14	9.01	68,900	30	0.27	2,000	24,376	3.29	52,235
St. Lucie	47	16	10.84	103,400	31	0.28	2,074	39,357	5.31	87,221
Santa Rosa	13	5	3.93	33,100	8	0.07	519	33,971	4.59	84,124
Sarasota	71	19	21.60	219,000	52	0.46	3,407	48,942	6.61	103,550
Seminole	29	15	48.95	278,200	14	0.12	889	32,268	4.36	86,107
Sumter	40	4	1.84	4,000	36	0.32	2,370	14,701	1.98	46,975
Suwannee	13	3	0.87	6,000	10	0.09	667	12,862	1.74	28,177
Taylor	7	2	0.97	4,700	5	0.04	296	7,231	0.98	14,260
Union	1	1	0.47	4,800	0	0.00	0	2,960	0.40	8,642
Volusia	126	18	31.49	284,700	108	0.96	7,111	74,430	10.05	151,532
Wakulla	9	2	0.29	2,100	7	0.06	444	7,812	1.05	20,319
Walton	10	6	2.77	16,000	4	0.04	296	17,244	2.33	24,305
Washington	6	3	0.75	3,200	3	0.03	222	7,952	1.07	17,551
State totals	2,789	571	1,494.86	11,070,000	2,218	19.75	146,291	1,944,704	262.56	4,766,086

Water Withdrawal Trends, 1970-2000

Statewide withdrawal and water-use estimates have been compiled for Florida every 5 years since 1950; however, variations in historical water-use values are sometimes difficult to assess because of differences in data-collection techniques and sources of information through the years. Since 1970, statewide water-use data for all withdrawal categories have been collected, tabulated, and published seven times by many agencies (five water management districts, the Florida Geological Survey, the Florida Department of Environmental Protection, and the U.S. Geological Survey).

Total water (fresh and saline) withdrawn in Florida increased by 4,990 Mgal/d or 33 percent between 1970 and 2000 (fig. 39 and table 16), during which time the population increased by 9.2 million people (135 percent) (fig. 1 and table 16). During this period, freshwater withdrawals increased nearly 46 percent (2,580 Mgal/d), and saline water withdrawals increased 25 percent (2,410 Mgal/d) (fig. 39 and table 16). Overall, water withdrawals were highest in 1975 and 1980 as power generation demands peaked during this time. Since then,

power generation demands have decreased due to the increase in water efficiency of new or modernized facilities and the recycling of cooling water after routing the water to cooling ponds or cooling towers. Between 1990 and 1995, freshwater withdrawals decreased by about 4 percent and saline withdrawals increased by 6 percent (Marella, 1999). In contrast, freshwater withdrawals increased 13 percent and saline withdrawals increased 9 percent between 1995 and 2000. This decrease in freshwater withdrawals between 1990 and 1995 as well as the increase between 1995 and 2000, can be attributed to significantly drier conditions (lower rainfall) in 1990 and 2000 during key times of the year coupled with significant wetter conditions (higher rainfall) during 1995. Since 1990, water-conservation and the use of alternative water sources such as reclaimed wastewater have helped offset some of the increase in water demands. It is difficult to determine how much effect these measures have had on water use nor how much rainfall changes water demands.




Figure 39. Historical fresh and saline water withdrawals in Florida, 1950-2000. (Modified from Marella, 1999.)

Table 16. Historical population and water withdrawals in Florida by source, 1950-2000

[Compiled by the U.S. Geological Survey, Tallahassee; values in million gallons per day; ----, data not available]

Year	Population, in millions		Water withdrawals, in million gallons per day				Total all water	Non-potable water ²	Reclaimed water ³
	Total	Served by public supply	Fresh Total	Fresh Ground	Fresh Surface	Saline water			
1950	2.77	1.66	1,454.0	614.0	840.0 ¹	1,200.0 ¹	2,654.0		
1955	3.86	2.30	2,167.0	1,017.0	1,150.0	645.0	2,812.0	---	---
1960	4.95	3.37	3,760.0	1,560.0	2,200.0	3,360.0	7,120.0	---	---
1965	5.87	4.81	6,852.0	2,218.5	4,633.5	6,261.0	13,113.0	---	---
1970	6.79	5.42	5,612.3	2,786.7	2,825.6	9,545.0	15,157.3	1.6	---
1975	8.48	6.81	6,772.7	3,214.6	3,558.1	11,502.0	18,274.7	1.7	---
1977	8.72	6.99	6,546.3	3,429.5	3,116.8	---	---	1.2	---
1980	9.75	7.79	6,701.2	3,677.2	3,024.0	13,897.0	20,598.2	2.5	---
1985	11.32	9.74	6,313.4	4,047.7	2,265.7	10,798.3	17,111.7	17.3	206.0 ⁴
1990	12.94	11.23	7,583.6	4,664.7	2,918.9	10,366.1	17,949.7	47.9	266.0
1995	14.15	12.21	7,229.9	4,348.8	2,881.1	10,965.7	18,195.6	57.9	402.0
2000	15.98	14.03	8,191.8	5,078.7	3,113.1	11,955.8	20,147.6	95.3	575.0

¹ Data for 1950 power generation did not show fresh or saline totals, but estimates were made using percentages from 1955.

² Nonpotable water includes water treated through desalination or diluted with fresher water to meet drinking water standards.

³ Reclaimed (waste) water includes all water reported by DEP as all types of reuse and wastewater disposal.

⁴ Value is reported for 1986, no data was compiled for 1985.

Data sources: Population

1950-1970 - University of Florida (Dietrich, 1978).

1975-2000 - Florida Statistical Abstracts and United States Statistical Abstracts.

Population served and water withdrawals

1950 - Modified from U.S. Geological Survey Circular 115 (MacKichan, 1951).

1955 - Modified from U.S. Geological Survey Circular 398 (MacKichan, 1957).

1960 - Modified from U.S. Geological Survey Circular 456 (MacKichan and Kammerer, 1961).

1965 - Modified from U.S. Geological Survey Circular 556 (Murray, 1968).

1970 - Modified from U.S. Geological Survey Circular 676 (Murray and Reeves, 1972).

1975 - USGS Water-Resources Investigations Report 78-17 (Leach, 1978).

1977 - USGS Water-Resources Investigations Report 79-112 (Leach and Healy, 1980).

1980 - USGS Water-Resources Investigations Report 82-4090 (Leach, 1983).

1985 - USGS Water-Resources Investigations Report 88-4103 (Marella, 1988).

1990 - USGS Water-Resources Investigations Report 92-4140 (Marella, 1992).

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

Reclaimed water

1985-2000 - 2000 Reuse Inventory (Florida Department of Environmental Protection, 2001)

Ground-water withdrawals increased 2,292 Mgal/d or 82 percent between 1970 and 2000 (fig. 40 and table 16). This long-term trend in ground-water withdrawals is a result of: (1) improvements in drilling techniques; (2) the ability to pump large quantities of high-quality water more economically from large, deep wells; and (3) increases in demand posed by population growth and crop irrigation. Since 1980, ground water has been the primary source of freshwater in Florida, supplying about 60 percent of the total freshwater withdrawn during this period.

Fresh surface-water withdrawals have increased 288 Mgal/d or 10 percent between 1970 and 2000 (fig. 40 and

table 16). Fresh surface-water withdrawals peaked in 1975, primarily due to power generation demands; however, since then many plants converted to more efficient closed systems and newer plants rely on saline surface water. Increases in fresh surface-water demands posed by increased irrigated agricultural acreage in South Florida during 1970 through 2000 helped offset the large decrease in withdrawals for power generation. Between 1950 and 1980, surface water was the primary source of freshwater in Florida, supplying more than half of the total freshwater withdrawn during this period. Surface water supplies about 40 percent of the State's freshwater withdrawn since 1980.




Figure 40. Historical freshwater withdrawals in Florida by source, 1950-2000. (Modified from Marella, 1999.)

Between 1970 and 2000, total freshwater withdrawals increased for public supply by 176 percent, for agricultural self-supplied by 87 percent, and domestic self-supplied remained nearly the same (table 17). However, during this period, withdrawals for commercial-industrial self-supplied decreased by 37 percent, and power generation decreased (thermoelectric) by 57 percent. Recreational irrigation withdrawals increased 127 percent between 1985 and 2000; prior to 1985 recreational irrigation water use was included in the agricultural self-supplied category. Increases in public-supply withdrawals are a result of population growth and tourism during this time period, along with higher demands posed by commercial and industrial facilities that are on public-supply water systems. Agricultural self-supplied increases during this period are a result of irrigation development and the availability of water for irrigation, which allowed farmers to increase crop production while decreasing the risk of crop damage due to prolonged dry conditions. Commercial-industrial self-supplied withdrawals decreased because self-supplied users began using public supply as a water source, many facilities began recirculating more water (primarily to lower their discharge amount), and the closure of several large paper mills and phosphate facilities. Power generation decreased between 1970 and 1990, as many of the

facilities changed to more modern and efficient cooling systems.

The use of nonpotable ground water as a source of supply, primarily for public supply, also has increased in Florida. Nonpotable implies that water does not meet drinking-water standards set by the FDEP and thus needs some form of desalination treatment (see Glossary) or dilution to meet set standards. The amount of nonpotable water treated to meet drinking-water standards increased from 17 Mgal/d in 1985 (Marella, 1995) to 95 Mgal/d in 2000 (table 16). Several public-supply water systems depend on nonpotable treated water, but most use this source to augment existing sources.

The use of reclaimed wastewater also has increased in Florida since the mid 1980's. According to the FDEP, nearly 575 Mgal/d of reclaimed wastewater was used in 2000, compared to 206 Mgal/d used in 1986 (Florida Department of Environmental Protection, 2001) (table 16). However, a large portion of this reported value is for the sole purpose of disposing treated wastewater, and not to offset freshwater demands. Only about one-third of this reclaimed water is used to replace or offset a freshwater source. The use of reclaimed wastewater reduces the demand for freshwater, and is expected to continue to increase statewide.

Table 17. Historical freshwater withdrawals in Florida by category, 1970-2000

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day, ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial self-supplied		Agricultural self-supplied		Recreational irrigation		Power Generation		Total freshwater withdrawals		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1970	753.10	130.30	1.6	209.20	0.00	683.60	215.90	1,136.35	964.35	----	4.50	1,515.00	2,786.75	2,825.55	5,612.30
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	962.80	161.30	1.7	225.75	2.05	721.85	160.70	1,289.90	1,640.70	----	14.30	1,593.40	3,214.60	3,558.15	6,772.75
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	1,059.10	172.80	1.2	213.00	1.01	703.68	153.54	1,437.29	1,479.50	----	16.43	1,309.93	3,429.50	3,116.78	6,546.28
1978	1,052.60	154.10	2.0	239.30	1.00	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	1,225.95	180.45	2.5	243.40	0.10	615.24	85.08	1,572.80	1,452.60	----	19.80	1,305.80	3,677.19	3,024.03	6,701.22
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	1,491.80	193.64	17.3	259.29	0.00	631.53	77.28	1,526.66	1,271.15	119.65	18.74	661.76	4,047.67	2,265.67	6,313.34
1986	1,542.77	191.28	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	1,634.68	199.73	37.4	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	1,693.21	211.29	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	1,754.10	217.80	42.4	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	1,698.82	226.33	47.9	299.38	0.00	630.88	139.06	1,800.19	1,695.03	212.31	23.14	760.72	4,664.72	2,918.86	7,583.58
1991	1,701.31	245.04	46.3	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	1,780.05	242.61	47.5	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	1,810.38	218.47	53.4	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1,800.06	210.64	58.4	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1,868.77	210.47	57.9	296.74	0.00	438.12	253.71	1,527.52	1,716.58	196.38	21.25	615.88	4,348.78	2,881.14	7,229.92
1996	1,922.40	208.98	65.5	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	1,929.16	221.04	66.0	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	2,057.04	218.38	75.1	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	2,087.59	239.07	88.5	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	2,199.36	237.43	95.3	198.68	0.00	430.70	132.60	1,989.95	1,933.06	230.45	29.53	628.73	5,078.67	3,113.10	8,191.77

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or diluted with fresher water to meet standards.

Domestic self-supplied withdrawals differ for 2000 as a revised procedure was used to calculate withdrawals compared to the previous years (1975-1995).

Commercial-industrial self-supplied includes water withdrawals for all mining purpose, as well as golf course irrigation from 1970-1984.

Agricultural self-supplied includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational irrigation includes water used for all turf grass irrigation (golf, commercial, industrial, and public) purposes. This category was accounted for under Agricultural self-supplied from 1970 through 1984.

Data sources: 1970-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-2000 - USGS unpublished water-use data files, Tallahassee, Florida.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - St. Johns River Water Management District, 2004; Southwest Florida Water Management District, 2002; and USGS Scientific Investigations Report 2004-5151.

Selected References

- American Water Works Association, 1992, Water industry data base: utility profiles: Denver, Colorado, American Water Works Association, 80 p.
- Bucca, Jane, and Marella, R.L., 1992, An improved method for determining the nonresidential water use component of total public water supply estimates, *in* Jones, M.E., and Laenen, Antonius (eds.), *Interdisciplinary approaches in hydrology and hydrogeology*: American Institute of Hydrology, p. 511-523.
- Buros, O.K., 1989, Desalting practices in the United States: American Water Works Association Journal, v. 81, no. 11, November, p. 38-42.
- Campbell, K.M., 1986. The industrial minerals of Florida: Tallahassee, Florida Bureau of Geology Information Circular No. 102, 94 p.
- Davis, W.Y., Rodrigo, D.M., Opitz, E.M., Dziegielewski, Bbenedykt, Baumann, D.D., and Boland, J.J., 1988, IWR-Main water use forecasting system, version 5.1: Users manual and system description: Fort Belvoir, Va., U.S. Army Corps of Engineers, Institute for Water Resources, IWR-Main Report 88-R-6, 273 p.
- Dietrich, T.S., 1978, The urbanization of Florida's population: An historical perspective of county growth 1830-1970: Gainesville, University of Florida, College of Business Administration, Bureau of Economic and Business Research, 211 p.
- Duerr, A.D., Hunn, J.D., Lewelling, B.R., and Trommer, J.T., 1988, Geohydrology and 1985 water withdrawals of the aquifer systems in southwest Florida, with emphasis on the intermediate aquifer system: U.S. Geological Survey Water-Resources Investigations 87-4259, 115 p.
- Dykes, G.M., and Conlon, W.J., 1989. Use of membrane technology in Florida: American Water Works Association Journal, v. 81, no. 11, November, p. 43-46.
- Fernald, E.A., and Patton, D.J., eds., 1984, Water resources atlas of Florida: Tallahassee, Florida State University, Institute of Science and Public Affairs, 291 p.
- Fernald, E.A., and Purdum, E.D., eds., 1992, Atlas of Florida: Tallahassee, Florida State University, Institute of Science and Public Affairs, 280 p.
- Fernald, E.A., and Purdum, E.D., eds., 1998, Water resource atlas of Florida: Tallahassee, Florida State University, Institute of Science and Public Affairs, 312 p.
- Florida Agricultural Statistics Service, 2000, Commercial citrus inventory 2000: Orlando, Florida Agricultural Statistics Service, Florida Department of Agriculture and Consumer Services, 115 p.
- 2001a, Field crop summary 2000: Orlando, Florida Agricultural Statistics Service, Florida Department of Agriculture and Consumer Services, 19 p.
- 2001b, Vegetable summary 1999-2000: Orlando, Florida Agricultural Statistics Service, Florida Department of Agriculture and Consumer Services, 62 p.
- 2001c, Livestock, dairy, and poultry summary 2000: Orlando, Florida Agricultural Statistics Service, Florida Department of Agriculture and Consumer Services, 79 p.
- 2002, Catfish - February, 2002: Orlando, Florida Agricultural Statistics Service, Florida Department of Agriculture and Consumer Services, 2 p.
- Florida Department of Agriculture and Consumer Services, 1996, Florida Agricultural Fast Facts, 1996 Directory: Tallahassee, Florida Department of Agricultural and Consumer Services, Division of Marketing and Development, 72 p.
- 2000, 2000-2001 Annual report: Gainesville, Bureau of Plant and Apiary Inspection, Division of Plant Industry, 63 p.
- 2002, Florida Agricultural Fast Facts, 2002 Directory: Tallahassee, Florida Department of Agricultural and Consumer Services, Division of Marketing and Development, 168 p.
- ¹Florida Department of Environmental Regulation, 1990, Drinking water standards, monitoring, and reporting: Tallahassee, chap. 17-550, 48 p.
- 1991, Domestic wastewater facilities: Tallahassee, chap. 17-600, 70 p.
- Florida Department of Environmental Protection, 2001, 2000 Reuse inventory: Tallahassee, Florida Department of Environmental Protection, Division of Resource Management, 113 p.
- Florida Sports Foundation, [2000], Florida destination golf - Florida's official golf guide: Tallahassee, Florida Sports Foundation, 56 p.
- Heath, R.C., and Conover, C.S., 1981, Hydrologic almanac of Florida: U.S. Geological Survey Open-File Report 81-1107, 239 p.
- Hodges, A.W., Haydu, J.J., van Blokland, P.J., and Bell, A.P., 1994, Contribution of the turf grass industry to Florida's economy, 1991/92: A value added approach: Gainesville, University of Florida, Institute of Food and Agricultural Sciences, Food and Resource Economics Department, Economic Report ER94-1, 83 p.
- ¹Florida Department of Environmental Regulation is now Florida Department of Environmental Protection.

- Hughes, G.H., 1975, Perspective on use of freshwater for cooling systems of thermoelectric power plants in Florida: U.S. Geological Survey Water-Resources Investigations 43-75, 30 p.
- Hutson, S.S., Barber, N.L., Kenny, J.F., Linsey, K.S., Lumia, D.S., and Maupin, M.A., 2004, Estimated use of water in the United States in 2000: Reston, Va., U.S. Geological Survey Circular 1268, 46 p.
- Izuno, F.T., and Haman, D.Z., 1987, Basic irrigation terminology: Gainesville, University of Florida, Institute of Food and Agricultural Sciences, Agricultural Engineering Fact Sheet AE-66, 4 p.
- Jones, J.W., Allen, L.H., Shih, S.F., Rogers, J.S., Hammond, L.C., Smajstrla, A.G., and Martsof, J.D., 1984, Estimated and measured evapotranspiration for Florida climate, crops, and soils: Gainesville, University of Florida, Institute of Food and Agricultural Sciences, Bulletin 840 (technical), 65 p.
- Knowles, Leel, Jr., O'Reilly, A.M., and Adamski, J.C., 2002, Hydrogeology and simulated effects of ground-water withdrawals from the Floridan aquifer system in Lake County and in the Ocala National Forest and vicinity, North-Central Florida: U.S. Geological Survey Water-Resources Investigations 02-4207, 140 p.
- Leach, S.D., 1978, Source, use, and disposition of water in Florida, 1975: U.S. Geological Survey Water-Resources Investigations 78-17, 90 p.
- 1983, Source, use, and disposition of water in Florida, 1980: U.S. Geological Survey Water-Resources Investigations 82-4090, 337 p.
- Leach, S.D., and Healy, H.G., 1980, Estimated water use in Florida, 1977: U.S. Geological Survey Water-Resources Investigations Report 79-112, 76 p.
- MacKichan, K.A., 1951, Estimated use of water in the United States, 1950: U.S. Geological Survey Circular 115, 13 p.
- 1957, Estimated use of water in the United States, 1955: U.S. Geological Survey Circular 398, 18 p.
- MacKichan, K.A., and Kammerer, J.C., 1961, Estimated use of water in the United States, 1960: U.S. Geological Survey Circular 456, 44 p.
- Marella, R.L., 1988, Water withdrawals, use, and trends in Florida, 1985: U.S. Geological Survey Water-Resources Investigations Report 88-4103, 43 p.
- 1992, Water withdrawals, use, and trends in Florida, 1990: U.S. Geological Survey Water-Resources Investigations Report 92-4140, 38 p.
- 1994, Estimated discharge of treated wastewater in Florida, 1990: U.S. Geological Survey Open-File Report 93-364, 53 p.
- 1995, Water-use data by category, county, and water management district in Florida, 1950-90: U.S. Geological Survey Open-File Report 94-521, 114 p.
- 1997, Irrigated crop acreage and water withdrawals in Florida, 1990: Florida Department of Environmental Protection, Tallahassee, Florida Geological Survey, Map Series 143, 1 sheet.
- 1998, Water-Quality assessment of Southern Florida-Wastewater discharge and runoff: U.S. Geological Survey Fact Sheet FS-032-98, 6 p.
- 1999, Water withdrawals, use, discharge, and trends in Florida, 1995: U.S. Geological Survey, Water-Resources Investigations Report 99-4002, 90 p.
- Marella, R.L. and Berndt, M.P., 2005, Water withdrawals and trends from the Floridan aquifer system in the southeastern United States, 2000: U.S. Geological Survey Circular 1278.
- McPherson, B.F., and Hammett, K.M., 1991, Tidal Rivers of Florida, *in* Livingston, R.J., ed., *The rivers of Florida: New York, Springer-Verlag Incorporated, Ecological Studies* 83, p. 31-46.
- Murray, C.R., 1968, Estimated use of water in the United States, 1965: U.S. Geological Survey Circular 556, 53 p.
- Murray, C.R., and Reeves, E.B., 1972, Estimated use of water in the United States, 1970: U.S. Geological Survey Circular 676, 37 p.
- 1977, Estimated use of water in the United States, 1975: U.S. Geological Survey Circular 765, 37 p.
- National Golf Foundation, 2001, Golf course directory - 2001 Edition: Jupiter, Florida, v. I, Alabama-Illinois, 480 p.
- Pride, R.W., 1973, Estimated use of water in Florida, 1970: Tallahassee, Florida Bureau of Geology Information Circular 83, 31 p.
- 1975, Estimated water use in Florida, 1965 (2d ed.): Tallahassee, Florida Bureau of Geology Map Series 36, 1 sheet.
- Purdum, E.D., 1994, Florida County Atlas and Municipal Fact Book: Tallahassee, Florida State University, Institute of Science and Public Affairs, 146 p.
- St. Johns River Water Management District, 1984, Consumptive uses of water; applicant's handbook: Palatka, Resource Management Department, June 1, 1984, 49 p.
- 2004, Water use in the St. Johns River Water Management District, 2000: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2004-FS1, 6 p.
- Seaber, P.R., Kapinos, F.P., and Knapp, G.L., 1984, State hydrologic unit maps: U.S. Geological Survey Open-File Report 84-708, 63 p.
- Sepulveda, Nicasio, 2002, Simulation of ground-water flow in the intermediate and floridan aquifer systems in peninsular Florida: U.S. Geological Survey Water-Resources Investigations 02-4009, 130 p.
- Shoemaker, W.B., and Edwards, K.M., 2003, Potential for salt-water intrusion into the lower Tamiami aquifer near Bonita Springs, southwestern Florida: U.S. Geological Survey Water-Resources Investigations 03-4262, 74 p.
- Smajstrla, A.G., 1986, Agricultural field scale irrigation requirements simulation model: Technical manual: Gainesville, University of Florida, Agricultural Engineering Department, 52 p.
- Smajstrla, A.G., Boman, B.J., Clark, G.A., Haman, D.Z., Harrison, D.S., Izuno, F.T., and Zazueta, F.S., 1988, Efficiencies of Florida agricultural irrigation systems:

- Gainesville, University of Florida, Institute of Food and Agricultural Sciences, Bulletin 247, 15 p.
- Smajstrla, A.G., and others, 1993, Microirrigation in Florida: Systems, acreage and cost: Gainesville, University of Florida, Institute of Food and Agricultural Sciences, Bulletin 276, 12 p.
- Smith, S.K., and Cody, Scott, 2001, Number of households and average household size in Florida: 1980, 1990 and 2000: Gainesville, University of Florida, Bureau of Economic and Business Research, v. 34, no. 2, Bulletin no. 129, 4 p.
- Smith, S.K., and Nogle, J.M., 2001, Projections of Florida population by county 2000-2030: Gainesville, University of Florida, Bureau of Economic and Business Research, v. 34, no. 1, Bulletin no. 128, 8 p.
- Solley, W.B., Chase, E.B., and Mann, W.B., 1983, Estimated use of water in the United States in 1980: U.S. Geological Survey Circular 1001, 56 p.
- Solley, W.B., Merk, C.F., and Pierce, R.R., 1988, Estimated use of water in the United States in 1985: U.S. Geological Survey Circular 1004, 82 p.
- Solley, W.B., Pierce, R.R., and Perlman, H.A., 1993, Estimated use of water in the United States in 1990: U.S. Geological Survey Circular 1081, 76 p.
- 1998, Estimated use of water in the United States in 1995: U.S. Geological Survey Circular 1200, 78 p.
- South Florida Water Management District, 1990, Desalination: An additional water source for south Florida: West Palm Beach, South Florida Water Management District, Office of Communications, (PIO 276 290 5M), 4 p.
- Southwest Florida Water Management District, 2002, 2000 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 151 p.
- Tchobanoglous, George, 1991, Wastewater engineering, treatment, disposal, and reuse: Metcalf and Eddy, Inc., 3rd edition, 1334 p.
- U.S. Census Bureau, 1993, 1990 Census of housing, detailed housing characteristics: Florida: Washington D.C., U.S. Department of Commerce, Bureau of the Census, Ch-2-11, 793 p.
- 2001, Statistical abstract of the United States 2002 (122nd ed.): Washington D.C., U.S. Department of Commerce, Bureau of the Census, 999 p.
- 2002, County business patterns, 2000, Florida: Washington D.C., U.S. Department of Commerce, Bureau of the Census, CBP-00-11, 327 p.
- U.S. Department of Agriculture, 1999, 1997 Census of Agriculture—Florida State and County data: Washington D.C., U.S. Department of Agriculture, National Agricultural Statistics Service, Geographic Area Series, Part 9 (AC 97-A-9), vol. 1, 444 p.
- U.S. Geological Survey, 1975, State of Florida, Hydrologic unit map-1974: U.S. Geological Survey, 1 sheet, scale 1:500,000.
- U.S. Soil Conservation Service, 1970, Irrigation water requirement (revised), U.S. Department of Agriculture Technical Release no. 21, 88 p.
- 1982, Florida irrigation guide: Gainesville, U.S. Department of Agriculture, Soil Conservation Service, 300 p.
- University of Florida, 2001, Florida Population: Census summary 2000: Gainesville, University of Florida, Bureau of Economic and Business Research, 59 p.
- 2002, Florida statistical abstract 2002 (36th ed.): Gainesville, University of Florida, Bureau of Economic and Business Research, 846 p.
- University of Nebraska, 2004, National Drought Summaries January 4-December 26, 2000: accessed February 19, 2004, <http://www.drought.unl.edu/dm/archive/2000>.
- Vecchioli, John, and Foose, D.W., 1985, Florida ground-water resources, in the National water summary, 1984—Hydrologic events, selected water quality trends and ground-water resources: U.S. Geological Survey Water-Supply Paper 2275, p. 173-178.

Water-Use Bibliography for Florida

Florida

- Conover, C.S., Vecchioli, John, and Foose, D.W., 1989, Ground-water sources and withdrawals for 1985: Tallahassee, Florida Bureau of Geology Map Series 124, 1 sheet.
- Healy, H.G., 1972, Public water supplies of selected municipalities in Florida, 1970: Tallahassee, Florida Bureau of Geology Information Circular no. 81, 213 p.
- 1977, Public water supplies of selected municipalities in Florida, 1975: U.S. Geological Survey Water-Resources Investigations Report 77-53, 309 p.
- Leach, S.D., 1978, Source, use, and disposition of water in Florida, 1975: U.S. Geological Survey Water-Resources Investigations 78-17, 90 p.
- 1978, Freshwater use in Florida, 1975: Tallahassee, Florida Bureau of Geology Map Series 87, 1 sheet.
- 1982, Estimated water use in Florida, 1980: Tallahassee, Florida Bureau of Geology Map Series 103, 1 sheet.
- 1982, Consumptive use of freshwater in Florida, 1980: Tallahassee, Florida Bureau of Geology Map Series 105, 1 sheet.
- 1983, Source, use, and disposition of water in Florida, 1980: U.S. Geological Survey Water-Resources Investigations 82-4090, 337 p.
- 1984, Projected public supply and rural (self-supplied) water use in Florida through the year 2020: Tallahassee, Florida Bureau of Geology Map Series 108, 1 sheet.

- Leach, S.D., and Healy, H.G., 1980, Estimated water use in Florida, 1977, U.S. Geological Survey Water-Resources Investigations Report 79-112, 76 p.
- Marella, R.L., 1988, Water withdrawals, use and trends in Florida, 1985: U.S. Geological Survey Water-Resources Investigations Report 88-4103, 43 p.
- 1989, Freshwater withdrawals, water-use trends in Florida, 1985: Tallahassee, Florida Bureau of Geology Map Series 123, 1 sheet.
- 1990, Florida water supply and use, *in* the National water summary 1987--Hydrologic events and water supply and use: U.S. Geological Survey Water-Supply Paper 2350, p. 207-214.
- 1990, Public supply water use in Florida, 1987: U.S. Geological Survey Open-File Report 90-596, 39 p.
- 1992, Factors that affect public-supply water use in Florida, with a section on projected water use to the year 2020: U.S. Geological Survey Water-Resources Investigations Report 91-4123, 35 p.
- 1992, Water withdrawals, use, and trends in Florida, 1990: U.S. Geological Survey Water-Resources Investigations Report 92-4140, 38 p.
- 1993, Public supply water use in Florida, 1990: U.S. Geological Survey Open-File Report 93-134, 46 p.
- 1994, Estimated discharge of treated wastewater in Florida, 1990: U.S. Geological Survey Open-File Report 93-364, 53 p.
- 1995, Water-use data by category, county, and water management district in Florida, 1950-90: U.S. Geological Survey Open-File Report 94-521, 114 p.
- 1997, Freshwater withdrawals, water-use trends in Florida, 1990: Tallahassee, Florida Department of Environmental Protection, Florida Geological Survey, Map Series 141, 1 sheet.
- 1997 Irrigated crop acreage and water withdrawals in Florida, 1990: Tallahassee, Florida Department of Environmental Protection, Florida Geological Survey, Map Series 143, 1 sheet.
- 1998, Water-Quality assessment of Southern Florida-Wastewater discharge and runoff: U.S. Geological Survey Fact Sheet FS-032-98, 6 p.
- 1999, Water withdrawals, use, discharge, and trends in Florida, 1995: U.S. Geological Survey, Water-Resources Investigations Report 99-4002, 90 p.
- Marella, R.L. and Berndt, M.P., 2005, Water withdrawals and trends from the Floridan aquifer system in the southeastern United States, 2000: U.S. Geological Survey Circular 1278.
- Pride, R.W., 1973, Estimated use of water in Florida, 1970: Tallahassee, Florida Bureau of Geology Information Circular 83, 31 p.
- 1975, Estimated water use in Florida, 1965 (2d ed.): Tallahassee, Florida Bureau of Geology Map Series 36, 1 sheet.
- Spechler, R.M., 1983, Estimated irrigation water use in Florida, 1980: Tallahassee, Florida Bureau of Geology Map Series 106, 1 sheet.

Northwest Florida Water Management District

- Bielby, Camille, 1987, 1985 Annual water use survey: Havana, Northwest Florida Water Management District Program Development Series 87-1, 107 p.
- Kranzer, B.S., 1983, Water use in the Northwest Florida Water Management District, an examination of current and past use: Havana, Northwest Florida Water Management District Water Resources Special Report 83-3, 47 p.
- Marella, R.L., Mokray, M.F., and Hallock-Solomon, M.J., 1998, Water use trends and demand projections in the Northwest Florida Water Management District: U.S. Geological Survey Open-File Report 98-269, 37 p.
- Northwest Florida Water Management District, 1981, Public water supply systems in the coastal areas of Escambia, Santa Rosa, Okaloosa, Bay, and Walton Counties: Havana, Northwest Florida Water Management District Water Resources Special Report 83-1, 153 p.
- 1996, 1995 Annual reuse report: Havana, Northwest Florida Water Management District, 29 p.
- 1998, District Water Supply Assessment: Havana, Northwest Florida Water Management District Water Resources Assessment 98-2, 143 p.
- Richards, C.J., 1993, Historical water use and projected demands for southeastern Santa Rosa County: Havana, Northwest Florida Water Management District Water Resources Technical File Report 93-1, 33 p.

St. Johns River Water Management District

- Florence, B.L., 1990, Annual water use survey: 1988: Palatka, St. Johns River Water Management District Technical Publication SJ 90-12, 128 p.
- 1991, Annual water use survey: 1989: Palatka, St. Johns River Water Management District Technical Publication SJ 91-6, 126 p.
- 1992, Annual water use survey: 1990: Palatka, St. Johns River Water Management District Technical Publication SJ 92-4, 122 p.
- 1994, Annual water use survey; 1991: Palatka, St. Johns River Water Management District Technical Publication SJ 94-4, 130 p.
- 1995, Annual water use survey: 1992: Palatka, St. Johns River Water Management District Technical Publication SJ 95-2, 128 p.
- 1996, Annual water use survey: 1993: Palatka, St. Johns River Water Management District Technical Publication SJ 96-1, 128 p.
- 1996, Annual water use survey: 1994: Palatka, St. Johns River Water Management District Technical Publication SJ 96-3, 126 p.

- Florence, B.L., and Moore, Cynthia, 1997, Annual water use survey; 1995: Palatka, St. Johns River Water Management District Technical Publication SJ 97-4, 128 p.
- 1999, Annual water use survey; 1996: Palatka, St. Johns River Water Management District Technical Publication SJ 99-3, 128 p.
- Lewis, Kathryn, Carriker, Roy, and Marella, R.L., 1981, Analysis of residential demand for water in the St. Johns River Water Management District: Palatka, St. Johns River Water Management District Technical Publication SJ 81-2, 107 p.
- Marella, R.L., 1981, Annual water use survey; 1979: Palatka, St. Johns River Water Management District Technical Publication SJ 81-3, 127.
- 1982, Annual water use survey; 1980: Palatka, St. Johns River Water Management District Technical Publication SJ 82-5, 107 p.
- 1983, Annual water use survey; 1981: Palatka, St. Johns River Water Management District Technical Publication SJ 83-9, 107 p.
- 1984, Annual water use survey; 1982: Palatka, St. Johns River Water Management District Technical Publication SJ 84-2, 97 p.
- 1984, Annual water use survey; 1983: Palatka, St. Johns River Water Management District Technical Publication SJ 84-5, 103 p.
- 1985, Annual water use survey; 1984: Palatka, St. Johns River Water Management District Technical Publication SJ 85-7, 113 p.
- 1986, Annual water use survey; 1985: Palatka, St. Johns River Water Management District Technical Publication SJ 86-5, 117 p.
- 1987, Annual water use survey; 1980 Revised Edition: Palatka, St. Johns River Water Management District Technical Publication SJ 82-5, 55 p.
- 1988, Annual water use survey; 1986: Palatka, St. Johns River Water Management District Technical Publication SJ 88-7, 128 p.
- 1990, Annual water use survey; 1987: Palatka, St. Johns River Water Management District Technical Publication SJ 90-4, 126 p.
- Scott, Elaine, 1980, Annual water use survey; 1978: Palatka, St. Johns River Water Management District Technical Publication SJ 80-5, 107 p.
- St. Johns River Water Management District, 1996, 1995 Annual reuse report: Palatka, St. Johns River Water Management District, 29 p.
- 1998, Water supply assessment, 1998: Palatka, St. Johns River Water Management District Technical Publication SJ 98-2, 154 p.
- 2001, Water use in the St. Johns River Water Management District, 1997: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2000-FS1, 6 p.
- 2001, Water use in the St. Johns River Water Management District, 1998: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2001-FS1, 6 p.
- 2002, Water use in the St. Johns River Water Management District, 1999: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2002-FS1, 6 p.
- 2004, Water use in the St. Johns River Water Management District, 2000: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2004-FS1, 6 p.

South Florida Water Management District

- Alvarez, J.A., and Bacon, D.D., 1988, Production zones of major public water supply wellfields for the counties of the South Florida Water Management District: West Palm Beach, South Florida Water Management District, Resource Planning Department, Technical Publication 88-4, app. II, 17 p.
- South Florida Water Management District, 1992, Water supply needs and sources 1990-2010: West Palm Beach, South Florida Water Management District, Planning Department, 204 p.
- 1996, 1995 Annual reuse report: West Palm Beach, South Florida Water Management District, Planning Department, 35 p.
- 1998, Districtwide Water Supply Assessment: West Palm Beach, South Florida Water Management District, 249 p.
- Woehlcke, L.C., Bucca, Jane, and Loving, D.R., 1982, A potable water use data base for South Florida, 1980: West Palm Beach, South Florida Water Management District, Planning Department, Technical Memorandum, 23 p.

Southwest Florida Water Management District

- Duerr, A.D., and Sohm, J.E., 1983, Estimated water use in Southwest Florida, 1981, and summary of annual water use, 1970, 1975, and 1977-81: U.S. Geological Survey Open-File Report 83-45, 75 p.
- Duerr, A.D., and Trommer, J.T., 1981, Estimated water use in the Southwest Florida Water Management District and adjacent areas, 1979: U.S. Geological Survey Open-File Report 81-56, 58 p.
- 1981, Estimated water use in the Southwest Florida Water Management District and adjacent areas, 1980: U.S. Geological Survey Open-File Report 81-1060, 60 p.
- 1982, The benchmark farm program--a method for estimated irrigation water use in Southwest Florida: U.S. Geological Survey Water-Resources Investigations Report 82-17, 49 p.
- Sorensen, L.A., 1992, 1989 and 1990 Estimated water use in the Southwest Florida Water Management District: Brooksville, Southwest Florida Water Management District, Planning Department, 132 p.

- Sorensen, L.A., Burns, L.L., Miles, D.M., Norris, Henry, and Tuttell, Maryellen, 1990, 1988 Estimated water use in the Southwest Florida Water Management District: Brooksville, Southwest Florida Water Management District, Planning Department, 96 p.
- Southwest Florida Water Management District, 1984, Estimated water use in the Southwest Florida Water Management District, 1982: Brooksville, Southwest Florida Water Management District Planning and Performance Evaluation Section, 20 p.
- 1994, Estimated water use in the Southwest Florida Water Management District, 1991 and 1992: Brooksville, Resource Projects Department, Conservation Projects Section, 109 p.
- 1995, 1993 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Projects Department, Conservation Projects Section, 111 p.
- 1996, 1994 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Projects Department, Conservation Projects Section, 111 p.
- 1996, 1995 Annual reuse report: Brooksville, Resource Projects Department, Conservation Projects Section, 31 p.
- 1997, 1995 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Projects Department, Conservation Projects Section, 54 p.
- 1997, 1996 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 54 p.
- 1998, Southwest Florida Water Management District water supply assessment 1995-2020: Brooksville, Resource Projects Department, 48 p.
- 1999, 1997 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Water Resource Projects Section, 39 p.
- 1999, 1998 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 39 p.
- 2001, 1999 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 39 p.
- 2002, 2000 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 151 p.
- Stieglitz, E.H., 1985, Estimated water use in the Southwest Florida Water Management District, 1983: Brooksville, Southwest Florida Water Management District, Planning Department, 33 p.
- 1985, Estimated water use in the Southwest Florida Water Management District, 1984: Brooksville, Southwest Florida Water Management District, Planning Department, 44 p.
- 1986, Estimated water use in the Southwest Florida Water Management District, 1985: Brooksville, Southwest Florida Water Management District, Planning Department, 53 p.
- Stieglitz, E.H., and Tomik, K.E., 1987, Estimated water use in the Southwest Florida Water Management District, 1986: Brooksville, Southwest Florida Water Management District, Planning Department, 57 p.
- Tuttell, Maryellen, and Sorensen, L.A., 1989, 1987 Estimated water use in the Southwest Florida Water Management District: Brooksville, Southwest Florida Water Management District, Planning Department, 99 p.

Suwannee River Water Management District

- Suwannee River Water Management District, 1979, Water use, 1977: Live Oak, Suwannee River Water Management District Information Circular #7, 47 p.
- 1998, Water supply assessment: Live Oak, Suwannee River Water Management District, 35.

Additional Water-Use Reports or Papers

- Barlow, P.M., 2003, Ground water in freshwater-saltwater environments of the Atlantic Coast: Reston, Va., U.S. Geological Survey Circular 1262, 113 p.
- Betz, J.V., 1984, Chapter 8, Water Use, in Fernald, E.A., and Purdum, E.D., eds., Water resource atlas of Florida: Tallahassee, Florida State University, Institute of Science and Public Affairs, p. 108-115.
- Bucca, Jane, and Marella, R.L., 1992, An improved method for determining the nonresidential water use component of total public water supply estimates, in Jones, M.E., and Laenen, Antonius (eds.), Interdisciplinary approaches in hydrology and hydrogeology: American Institute of Hydrology, p. 511-523.
- Duerr, A.D., and Trommer, J.T., 1982, The benchmark farm program--a method for estimated irrigation water use in southwest Florida: U.S. Geological Survey Water-Resources Investigations Report 82-17, 49 p.
- Dykes, G.M., and Conlon, W.J., 1989, Use of membrane technology in Florida: American Water Works Association Journal, v. 81, no. 11, November, p. 43-46.
- Fanning, J.L., 1997, Water use in Georgia by county for 1995: Atlanta, Georgia Department of Natural Resources, Environmental Protection Division, Georgia Geological Survey, Information Circular 101, 95 p.
- , 2003, Water use in Georgia by county for 2000 and water-use trends for 1980-2000, Atlanta, Georgia Department of Natural Resources, Environmental

50 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

- Protection Division, Georgia Geological Survey, Information Circular 106, 176 p
- Franks, B.J., 1981, Land application of domestic wastewater in Florida--Statewide assessment of impact on ground-water quality: U.S. Geological Survey Water-Resources Investigations Report 81-3, 37 p.
- Harrison, D.S., Smajstrla, A.G., Choate, R.E., and Isaacs, G.W., 1983, Irrigation in Florida Agriculture in the 80's: Gainesville, University of Florida, Institute of Food and Agricultural Sciences, Bulletin 196, 10 p.
- Holland, T.W., 1992, Water-use data collection techniques in the southeastern United States, Puerto Rico, and the U.S. Virgin Islands: U.S. Geological Survey Water-Resources Investigations Report 92-4028, 76 p.
- Hughes, G.H., 1975, Perspective on use of freshwater for cooling systems of thermoelectric powerplants in Florida: U.S. Geological Survey Water-Resources Investigations Report 43-75, 30 p.
- Marella, R.L., 1983, Ground-water withdrawals from the Floridan aquifer in Duval County--1980: Palatka, St. Johns River Water Management District Technical Publication SJ 83-7, 1 sheet.
- 1984, Ground-water withdrawals from the Floridan aquifer in the Nassau County Area: Palatka, St. Johns River Water Management District Map Series 84-4, 1 sheet.
- 1984, Ground-water withdrawals from the Floridan aquifer in the Clay and portions of Bradford Counties: Palatka, St. Johns River Water Management District Map Series 84-14, 1 sheet.
- 1987, Water use, *in* the Indian River Lagoon Joint Reconnaissance Report, St. Johns River Water Management District and the South Florida Water Management District, November 1987, chap. 8, p. 8-1 to 8-9.
- 1998, Water Quality Assessment of Southern Florida--Wastewater discharges and runoff: U.S. Geological Survey Fact Sheet 032-98, 6 p.
- Marella, R.L. and Berndt, M.P., 2005, Water withdrawals and trends from the Floridan aquifer system in the southeastern United States, 2000: U.S. Geological Survey Circular 1278.
- Marella, R.L., Fanning, J.L., and Mooty, W.S., 1993, Estimated use of water in the Apalachicola-Chattahoochee-Flint River basin during 1990, with State summaries for 1970 to 1990: U.S. Geological Survey Water-Resources Investigations Report 93-4084, 45 p.
- Marella, R.L., and Fanning, J.L., 1996, National Water Quality Assessment of the Georgia-Florida Coastal Plain study unit--Water withdrawals and treated wastewater discharges, 1990: U.S. Geological Survey Water-Resources Investigations Report 95-4084, 76 p.
- Marella, R.L., and York, D.W., 1998, Chapter 6, Water Use, *in* Fernald, E.A., and Purdum, E.D., eds., Water resource atlas of Florida: Tallahassee, Florida State University, Institute of Science and Public Affairs, p. 114-135.
- Miller, M.L., and Alvarez, J.A., 1984, Public-supply water use, Palm Beach County, Florida, 1978-82: U.S. Geological Survey Open-File Report 84-240, 14 p.
- Mooty, W.S., and Richardson, J.R., 1998, Water use in Alabama, 1995: U.S. Geological Survey Water Resources Investigations Report 98-4154, 92 p.
- O'Donnell, T.H., 1977, Municipal water suppliers in Lee County, 1974: U.S. Geological Survey Open-File Report 77-277, 96 p.
- Vickers, Amy, 2001, Hand book of water use and conservation, Water Plow Press, Amherst, MA, ISBN 1-931579-07-05, 446 p.

Appendixes

Appendix 1 and 2

Purpose, Data Sources, and Limitations

Appendix 1 provides water-use data for each Florida county for the period 1965-2000; appendix 2 provides water-use data for each water management district for the period 1975-2000. These tables provide a summary of all water-use data collected since 1965 (published or not published) for each category by the many different agencies involved in water-use compilations throughout the State. Since 1965, water-use data have been collected and compiled on various levels by the Florida Bureau of Geology, Florida Department of Environmental Protection, Northwest Florida Water Management District, St. Johns River Water Management District, South Florida Water Management District, Southwest Florida Water Management District, Suwannee River Water Management District, and U.S. Geological Survey. Prior to 1965, water-use data were only collected and published on a Statewide level. Some county data are available prior to 1965 and are published in the Florida Bureau of Geology regional or county-wide water-resource investigations. However, these data are often not consistent from one county or category to another. All totaled, data presented in appendix 1 and 2 were compiled from nearly 60 publications and countless unpublished data files.

The appendix in this report is the second effort to present all available water-use data by county and water management district. Historical data were summarized and published in 1995

for the period between 1965 and 1990 by the USGS (Marella, 1995). The tables presented in this appendix will provide the most recent data for the years between 1990 and 2000 and also will include any data modifications for the period between 1965 and 1990. Some additional data have been added to these tables that depict the amount of public-supply water treated and the amount of water used for recreational irrigation (including golf courses). These tables are available to download from the USGS web site: http://fl.water.usgs.gov/Water-Use/Water_Use_web/index.htm.

Water withdrawals and use data presented herein represent the average daily quantities used, as derived from annual data, and are expressed in million gallons per day (Mgal/d). Water-use values are reported to two places to the right of the decimal or to the nearest 10,000 gallons per day (gal/d). Water-use data herein may not be identical to the water-use data published by the water management districts or FDEP because of differences in data-collection procedures, categories, methodology, or changes made after publishing.

Data collected between 1975 and 2000 by the water management districts for individual counties are for the portion of the county within that district. If a county is within two water management districts and only one district compiled data, then the county totals are not complete, and therefore, they are only published or presented for that district. No partial county data are presented in these tables, only complete county data sets unless otherwise noted.

Selected References for Appendix 1 and 2

- Bielby, Camille, 1987, 1985 Annual water use survey: Havana, Northwest Florida Water Management District Program Development Series 87-1, 107 p.
- Duerr, A.D., and Sohm, J.E., 1983, Estimated water use in Southwest Florida, 1981, and summary of annual water use, 1970, 1975, and 1977-81: U.S. Geological Survey Open-File Report 83-45, 75 p.
- Duerr, A.D., and Trommer, J.T., 1981, Estimated water use in the Southwest Florida Water Management District and adjacent areas, 1979: U.S. Geological Survey Open-File Report 81-56, 58 p.
- 1981, Estimated water use in the Southwest Florida Water Management District and adjacent areas, 1980: U.S. Geological Survey Open-File Report 81-1060, 60 p.
- Florence, B.L., 1990, Annual water use survey: 1988: Palatka, St. Johns River Water Management District Technical Publication SJ 90-12, 128 p.
- 1991, Annual water use survey: 1989: Palatka, St. Johns River Water Management District Technical Publication SJ 91-6, 126 p.
- 1992, Annual water use survey: 1990: Palatka, St. Johns River Water Management District Technical Publication SJ 92-4, 122 p.
- 1994, Annual water use survey; 1991: Palatka, St. Johns River Water Management District Technical Publication SJ 94-4, 130 p.
- 1995, Annual water use survey: 1992: Palatka, St. Johns River Water Management District Technical Publication SJ 95-2, 128 p.

- 1996, Annual water use survey; 1993: Palatka, St. Johns River Water Management District Technical Publication SJ 96-1, 128 p.
- 1996, Annual water use survey; 1994: Palatka, St. Johns River Water Management District Technical Publication SJ 96-3, 126 p.
- Florence, B.L., and Moore, Cynthia, 1997, Annual water use survey; 1995: Palatka, St. Johns River Water Management District Technical Publication SJ 97-4, 128 p.
- 1999, Annual water use survey; 1996: Palatka, St. Johns River Water Management District Technical Publication SJ 99-3, 128 p.
- Kranzer, B.S., 1983, Water use in the Northwest Florida Water Management District, an examination of current and past use: Havana, Northwest Florida Water Management District Water Resources Special Report 83-3, 47 p.
- Leach, S.D., 1978, Source, use, and disposition of water in Florida, 1975: U.S. Geological Survey Water-Resources Investigations 78-17, 90 p.
- 1983, Source, use, and disposition of water in Florida, 1980: U.S. Geological Survey Water-Resources Investigations 82-4090, 337 p.
- Leach, S.D., and Healy, H.G., 1980, Estimated water use in Florida, 1977, U.S. Geological Survey Water-Resources Investigations Report 79-112, 76 p.
- Marella, R.L., 1981, Annual water use survey; 1979: Palatka, St. Johns River Water Management District Technical Publication SJ 81-3, 127
- 1982, Annual water use survey; 1980: Palatka, St. Johns River Water Management District Technical Publication SJ 82-5, 107 p
- 1983, Annual water use survey; 1981: Palatka, St. Johns River Water Management District Technical Publication SJ 83-9, 107 p.
- 1984, Annual water use survey; 1982: Palatka, St. Johns River Water Management District Technical Publication SJ 84-2, 97 p.
- 1984, Annual water use survey; 1983: Palatka, St. Johns River Water Management District Technical Publication SJ 84-5, 103 p.
- 1985, Annual water use survey; 1984: Palatka, St. Johns River Water Management District Technical Publication SJ 85-7, 113 p.
- 1986, Annual water use survey; 1985: Palatka, St. Johns River Water Management District Technical Publication SJ 86-5, 117 p.
- 1987, Annual water use survey; 1980 Revised Edition: Palatka, St. Johns River Water Management District Technical Publication SJ 82-5, 55 p.
- 1988, Water withdrawals, use and trends in Florida, 1985: U.S. Geological Survey Water-Resources Investigations Report 88-4103, 43 p.
- 1988, Annual water use survey; 1986: Palatka, St. Johns River Water Management District Technical Publication SJ 88-7, 128 p.
- 1990, Annual water use survey; 1987: Palatka, St. Johns River Water Management District Technical Publication SJ 90-4, 126 p.
- 1992, Water withdrawals, use, and trends in Florida, 1990: U.S. Geological Survey Water-Resources Investigations Report 92-4140, 38 p.
- 1995, Water-use data by category, county, and water management district in Florida, 1950-90: U.S. Geological Survey Open-File Report 94-521, 114 p.
- 1999, Water withdrawals, use, discharge, and trends in Florida, 1995: U.S. Geological Survey, Water-Resources Investigations Report 99-4002, 90 p.
- 2004, Water withdrawals, use, discharge, and trends in Florida, 2000: U.S. Geological Survey Scientific Investigations Report 2004-5151, 136 p.
- Pride, R.W., 1973, Estimated use of water in Florida, 1970: Tallahassee, Florida Bureau of Geology Information Circular 83, 31 p.
- 1975, Estimated water use in Florida, 1965 (2d ed.): Tallahassee, Florida Bureau of Geology Map Series 36, 1 sheet.
- Scott, Elaine, 1980, Annual water use survey; 1978: Palatka, St. Johns River Water Management District Technical Publication SJ 80-5, 107 p.
- St. Johns River Water Management District, 2001, Water use in the St. Johns River Water Management District, 1997: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2000-FS1, 6 p.
- 2001, Water use in the St. Johns River Water Management District, 1998: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2001-FS1, 6 p.
- 2002, Water use in the St. Johns River Water Management District, 1999: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2002-FS1, 6 p.
- 2004, Water use in the St. Johns River Water Management District, 2000: Palatka, St. Johns River Water Management District Technical Fact Sheet SJ2004-FS1, 6 p.
- Sorensen, L.A., 1992, 1989 and 1990 Estimated water use in the Southwest Florida Water Management District: Brooksville, Southwest Florida Water Management District, Planning Department, 132 p.
- Sorensen, L.A., Burns, L.L., Miles, D.M., Norris, Henry, and Tuttell, Maryellen, 1990, 1988 Estimated water use in the Southwest Florida Water Management District: Brooksville, Southwest Florida Water Management District, Planning Department, 96 p.
- Southwest Florida Water Management District, 1984, Estimated water use in the Southwest Florida Water Management District, 1982: Brooksville, Southwest Florida Water Management District Planning and Performance Evaluation Section, 20 p.
- 1994, Estimated water use in the Southwest Florida Water Management District, 1991 and 1992: Brooksville,


54 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000

- Resource Projects Department, Conservation Projects Section, 109 p.
- 1995, 1993 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Projects Department, Conservation Projects Section, 111 p.
- 1996, 1994 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Projects Department, Conservation Projects Section, 111 p.
- 1997, 1995 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Projects Department, Conservation Projects Section, 54 p.
- 1997, 1996 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 54 p
- 1999, 1997 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Water Resource Projects Section, 39 p
- 1999, 1998 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 39 p
- 2001, 1999 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 39 p
- 2002, 2000 Estimated water use in the Southwest Florida Water Management District: Brooksville, Resource Conservation and Development Department, Conservation Projects Section, 151 p
- Stieglitz, E.H., 1985, Estimated water use in the Southwest Florida Water Management District, 1983: Brooksville, Southwest Florida Water Management District, Planning Department, 33 p.
- 1985, Estimated water use in the Southwest Florida Water Management District, 1984: Brooksville, Southwest Florida Water Management District, Planning Department, 44 p.
- 1986, Estimated water use in the Southwest Florida Water Management District, 1985: Brooksville, Southwest Florida Water Management District, Planning Department, 53 p.
- Stieglitz, E.H., and Tomik, K.E., 1987, Estimated water use in the Southwest Florida Water Management District, 1986: Brooksville, Southwest Florida Water Management District, Planning Department, 57 p.
- Tuttell, Maryellen, and Sorensen, L.A., 1989, 1987 Estimated water use in the Southwest Florida Water Management District: Brooksville, Southwest Florida Water Management District, Planning Department, 99 p.
- Suwannee River Water Management District, 1979, Water use, 1977: Live Oak, Suwannee River Water Management District Information Circular #7, 47 p.

Appendix 1. Freshwater withdrawals by category for each county in Florida, 1965-2000

County	Page	County	Page
Alachua	57	Lake	90
Baker	58	Lee	91
Bay	59	Leon	92
Bradford	60	Levy	93
Brevard	61	Liberty	94
Broward	62	Madison	95
Calhoun	63	Manatee	96
Charlotte	64	Marion	97
Citrus	65	Martin	98
Clay	66	Miami-Dade	99
Collier	67	Monroe	100
Columbia	68	Nassau	101
DeSoto	69	Okaloosa	102
Dixie	70	Okeechobee	103
Duval	71	Orange	104
Escambia	72	Osceola	105
Flagler	73	Palm Beach	106
Franklin	74	Pasco	107
Gadsden	75	Pinellas	108
Gilchrist	76	Polk	109
Glades	77	Putnam	110
Gulf	78	St. Johns	111
Hamilton	79	St. Lucie	112
Hardee	80	Santa Rosa	113
Hendry	81	Sarasota	114
Hernando	82	Seminole	115
Highlands	83	Sumter	116
Hillsborough	84	Suwannee	117
Holmes	85	Taylor	118
Indian River	86	Union	119
Jackson	87	Volusia	120
Jefferson	88	Wakulla	121
Lafayette	89	Walton	122
		Washington	123

56 Water Withdrawals, Use, Discharge, and Trends in Florida, 2000



Appendix 1. Freshwater withdrawals by category in Alachua County, 1965-2000

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	8.60	0.00	0.00	0.00	13.00	0.00	2.41	0.75	----	----	1.00	0.00	25.67	0.75	26.42
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	22.30	0.00	0.00	0.00	1.40	0.00	2.70	0.70	----	----	0.51	0.00	34.10	0.70	34.80
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	14.90	0.00	0.00	0.00	6.53	0.00	5.07	2.09	----	----	0.50	0.00	30.81	2.09	32.90
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	16.52	0.00	0.00	0.00	7.18	0.00	7.44	1.92	----	----	1.30	0.00	36.52	1.92	38.44
1978	18.14	0.00	0.00	0.00	4.10	0.00	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	18.18	0.00	0.00	0.00	1.53	0.00	15.97	4.32	----	----	0.93	0.00	44.06	4.32	48.38
1981	19.06	0.00	0.00	0.00	2.08	0.00	----	----	----	----	----	----	N/A	N/A	N/A
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	21.25	0.00	0.00	0.00	3.61	0.00	13.96	0.04	1.56	0.00	3.26	0.00	51.42	0.04	51.46
1986	21.71	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	21.88	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	22.20	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	23.63	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	22.95	0.00	0.00	0.00	2.29	0.00	16.06	0.25	2.03	0.11	2.41	0.00	52.12	0.36	52.48
1991	21.54	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	22.32	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	23.41	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	23.55	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	24.09	0.00	0.00	0.00	2.53	0.00	10.36	0.45	1.30	0.06	2.53	0.24	47.56	0.75	48.31
1996	24.68	0.00	0.00	0.00	2.59	0.00	----	----	----	----	2.59	0.00	N/A	N/A	N/A
1997	24.79	0.00	0.00	0.00	2.24	0.00	----	----	----	----	2.42	0.00	N/A	N/A	N/A
1998	27.47	0.00	0.00	0.00	2.21	0.00	----	----	----	----	2.50	0.00	N/A	N/A	N/A
1999	27.77	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	28.26	0.00	0.00	0.00	2.50	0.00	17.91	0.27	4.16	0.30	2.63	0.00	59.58	0.57	60.15

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Baker County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.20	0.00	0.25	0.00	0.15	0.00	0.11	0.01	---	---	0.00	0.00	0.71	0.01	0.72
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.50	0.00	0.79	0.00	0.00	0.00	0.00	0.00	---	---	0.00	0.00	1.29	0.00	1.29
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.54	0.00	0.82	0.00	0.32	0.00	0.88	0.72	---	---	0.00	0.00	2.56	0.72	3.28
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.56	0.00	0.86	0.00	0.32	0.00	0.96	1.16	---	---	0.00	0.00	2.70	1.16	3.86
1978	0.55	0.00	1.01	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	0.61	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1980	0.60	0.00	1.55	0.00	0.30	0.00	1.32	1.51	---	---	0.00	0.00	3.77	1.51	5.28
1981	0.54	0.00	---	---	0.50	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.60	0.00	1.86	0.00	0.66	0.00	2.95	2.19	0.00	0.00	0.00	0.00	6.07	2.19	8.26
1986	0.63	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.63	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.65	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.73	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.81	0.00	2.84	0.00	0.92	0.00	3.30	2.20	0.00	0.00	0.00	0.00	7.87	2.20	10.07
1991	0.76	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.74	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.76	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.71	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.68	0.00	2.50	0.00	0.54	0.00	0.93	0.63	0.09	0.00	0.00	0.00	4.74	0.63	5.37
1996	0.75	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.76	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.77	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.82	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.88	0.00	1.90	0.00	0.43	0.00	2.80	1.64	0.14	0.09	0.00	0.00	6.15	1.73	7.88

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.
 Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.
 Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.
 Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).
 1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.
 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).
 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Bay County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	5.10	0.00	1.80	0.00	2.40	31.70	0.10	0.02	---	---	0.29	0.00	9.69	31.72	41.41
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.70	37.40	4.38	0.00	2.00	0.50	0.30	0.00	---	---	0.12	0.00	7.50	37.90	45.40
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	1.95	32.59	0.89	0.00	1.35	0.00	0.07	0.16	---	---	0.68	0.00	4.94	32.75	37.69
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	2.91	36.05	1.76	0.00	1.32	0.00	0.40	0.01	---	---	0.70	0.00	7.09	36.06	43.15
1978	2.78	36.85	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	3.16	36.34	1.57	0.00	1.48	0.00	0.38	0.28	---	---	0.70	0.00	7.29	36.62	43.91
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	3.85	36.40	5.56	0.00	0.61	0.92	0.01	0.07	1.27	0.82	0.74	0.00	12.04	38.21	50.25
1986	6.17	41.78	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	6.47	39.40	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	6.43	40.10	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	6.14	38.81	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	7.07	40.11	5.47	0.00	0.65	1.46	0.26	0.00	1.43	0.59	1.14	0.00	16.02	42.16	58.18
1991	5.88	40.02	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	6.26	40.97	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	6.43	36.82	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	5.61	39.02	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	5.09	44.23	6.48	0.00	0.51	0.00	0.29	0.00	1.09	0.37	0.70	0.00	14.16	44.60	58.76
1996	4.78	41.62	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	5.16	44.92	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	5.60	36.99	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	5.60	45.39	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	6.28	44.89	2.01	0.00	0.30	0.00	1.28	0.00	1.46	0.79	0.89	0.00	12.22	45.68	57.90

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Bradford County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.30	0.00	0.30	0.00	0.00	0.00	0.14	0.07	---	---	0.07	0.00	0.81	0.07	0.88
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.70	0.00	1.06	0.00	1.40	0.00	0.07	0.07	---	---	0.02	0.00	3.25	0.07	3.32
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.83	0.00	0.96	0.00	3.96	0.00	0.09	0.09	---	---	0.00	0.00	5.84	0.09	5.93
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.08	0.00	1.28	0.00	3.37	0.00	0.91	0.10	---	---	0.10	0.00	6.74	0.10	6.84
1978	0.92	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	1.05	0.00	2.08	0.00	4.77	0.00	0.84	0.05	---	---	0.00	0.00	8.74	0.05	8.79
1981	1.17	0.00	---	---	4.47	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	1.34	0.00	2.60	0.00	4.34	0.00	0.89	0.00	0.08	0.00	0.00	0.00	9.25	0.00	9.25
1986	1.35	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	1.49	0.00	2.67	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	1.65	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	1.49	0.00	2.75	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	1.61	0.00	3.33	0.00	2.99	0.00	0.82	0.06	0.21	0.00	0.00	0.00	8.96	0.06	9.02
1991	1.58	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	1.47	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	1.52	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	1.40	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	1.33	0.00	2.38	0.00	2.98	0.00	0.69	0.05	0.08	0.00	0.00	0.00	7.46	0.05	7.51
1996	1.36	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	1.25	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	1.36	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	1.37	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	1.38	0.00	1.89	0.00	1.25	0.00	1.00	0.02	0.27	0.04	0.00	0.00	5.79	0.06	5.85

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Brevard County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	6.97	3.60	13.93	4.00	0.00	0.00	56.84	5.18	---	---	0.20	0.00	69.01	8.78	77.79
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	3.48	7.72	15.30	3.38	0.00	0.00	47.90	0.00	---	---	0.03	0.00	55.19	7.72	62.91
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	3.82	8.90	14.40	2.70	0.00	0.00	40.17	21.72	---	---	0.50	0.00	47.64	30.62	78.26
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	4.60	10.80	14.72	2.84	0.00	0.00	39.06	2.88	---	---	0.29	0.00	47.24	13.68	60.92
1978	5.38	9.46	14.97	4.42	0.00	0.24	123.13	5.75	---	---	0.10	0.00	133.27	15.21	148.48
1979	6.54	11.88	17.51	3.30	0.00	0.23	249.22	32.17	---	---	0.26	0.00	259.55	44.05	303.60
1980	7.25	12.97	17.73	4.10	0.00	0.24	158.99	7.37	---	---	0.25	0.00	170.83	20.34	191.17
1981	7.13	12.41	17.03	1.56	0.00	0.15	356.14	54.99	---	---	28.00	0.00	392.98	67.40	460.38
1982	6.87	11.33	16.44	4.55	0.00	0.15	76.69	15.27	---	---	0.24	0.00	88.50	26.60	115.10
1983	7.22	11.91	16.70	4.45	0.00	0.16	76.99	15.64	---	---	0.31	0.00	89.13	27.55	116.68
1984	8.46	13.21	20.34	5.09	0.00	0.14	85.43	16.45	---	---	0.28	0.00	99.40	29.66	129.06
1985	9.21	14.09	22.10	5.62	0.00	0.24	98.99	23.54	1.33	2.16	0.31	0.00	115.70	39.79	155.49
1986	9.59	15.47	22.47	5.92	0.00	0.57	104.86	8.40	2.27	2.47	0.28	0.00	123.49	26.34	149.83
1987	10.08	15.61	23.49	6.13	0.00	0.21	91.13	9.82	1.85	1.78	0.27	0.00	109.67	27.21	136.88
1988	10.79	16.15	25.02	6.89	0.00	0.19	98.23	12.37	2.26	2.28	0.23	0.00	118.59	30.80	149.39
1989	11.29	15.49	24.86	6.98	0.00	0.19	89.13	8.22	2.50	2.64	0.25	0.00	110.34	26.35	136.69
1990	11.55	16.24	23.52	5.08	0.00	0.19	99.41	9.07	1.37	1.33	0.25	0.00	117.85	26.64	144.49
1991	10.70	15.24	22.46	4.93	0.00	0.13	109.95	9.10	2.81	2.49	0.28	0.00	128.80	26.83	155.63
1992	11.26	14.83	24.85	0.87	0.00	0.13	83.69	8.49	1.48	1.34	0.25	0.00	97.68	24.66	122.34
1993	11.40	14.92	25.06	0.97	0.00	0.15	139.42	11.18	1.74	2.88	0.28	0.00	153.96	28.98	182.94
1994	10.69	15.71	23.54	0.83	0.00	0.87	136.51	11.41	1.77	2.94	0.32	0.00	150.99	30.06	181.05
1995	14.97	12.15	23.97	5.23	0.00	1.80	88.43	8.56	2.29	1.79	0.31	0.00	113.03	22.50	135.53
1996	17.16	11.07	25.34	5.36	0.00	1.75	96.61	9.33	3.29	2.79	0.33	0.00	124.50	23.19	147.69
1997	14.12	11.85	24.61	5.23	0.00	1.67	50.67	5.01	2.38	2.22	0.34	0.00	74.41	19.08	93.49
1998	15.06	12.15	25.73	5.21	0.00	3.46	165.68	13.29	3.17	2.73	0.49	0.00	193.07	28.17	221.24
1999	14.91	12.11	24.50	5.10	0.00	2.66	50.25	5.28	2.22	1.92	0.41	0.00	75.55	19.31	94.86
2000	13.66	14.08	25.61	1.90	0.00	1.04	118.79	18.03	7.75	7.08	0.39	0.00	143.53	39.20	182.73

Public supply transfers represent water withdrawn in Orange County and imported for use in Brevard County.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Broward County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	74.90	0.00	10.20	0.00	3.00	0.00	21.37	63.57	---	---	0.09	0.00	109.56	63.57	173.13
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	102.00	1.00	8.41	0.00	2.00	1.00	30.00	39.80	---	---	0.14	0.00	142.55	41.80	184.35
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	139.78	0.00	8.16	0.00	2.50	1.00	17.91	59.96	---	---	0.15	0.00	168.50	60.96	229.46
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	155.85	0.00	3.12	0.00	2.98	1.00	19.54	60.58	---	---	0.11	0.00	181.60	61.58	243.18
1978	154.90	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	184.39	0.00	5.89	0.00	1.19	0.00	32.90	11.11	---	---	0.10	0.00	224.47	11.11	235.58
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	187.95	0.00	2.96	0.00	1.36	0.00	25.33	0.79	12.51	4.31	0.07	0.00	230.18	5.10	235.28
1986	198.60	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	207.64	0.00	4.76	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	218.30	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	220.92	0.00	3.40	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	192.53	0.00	9.60	0.00	1.63	0.00	2.18	13.74	38.79	8.01	0.05	0.00	244.78	21.75	266.53
1991	206.28	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	213.68	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	217.62	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	218.81	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	222.30	0.00	2.16	0.00	0.34	0.00	3.33	6.70	39.03	13.05	0.42	0.00	267.58	19.75	287.33
1996	235.35	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	228.94	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	237.22	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	246.38	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	258.06	0.00	2.11	0.00	0.54	0.00	0.89	3.21	11.86	25.14	0.00	0.00	273.46	28.35	301.81

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Calhoun County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.20	0.00	0.80	0.00	1.10	0.50	0.41	0.01	---	---	0.00	0.00	2.51	0.51	3.02
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.20	0.00	0.53	0.00	0.00	0.00	0.13	0.03	---	---	0.00	0.00	0.86	0.03	0.89
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.28	0.00	0.50	0.01	0.36	0.00	1.21	0.15	---	---	0.00	0.00	2.35	0.16	2.51
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.32	0.00	0.54	0.01	0.36	0.00	1.25	1.29	---	---	0.00	0.00	2.47	1.30	3.77
1978	0.35	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	0.33	0.00	0.69	0.02	0.72	0.00	1.58	0.12	---	---	0.00	0.00	3.32	0.14	3.46
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.39	0.00	0.92	0.00	0.00	0.00	0.34	0.03	0.00	0.00	0.00	0.00	1.65	0.03	1.68
1986	0.41	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.40	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.40	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.43	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.53	0.00	1.24	0.00	0.00	0.00	0.11	0.50	0.00	0.00	0.00	0.00	1.88	0.50	2.38
1991	0.50	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.52	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.55	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.57	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.68	0.00	1.27	0.00	0.00	0.00	1.43	0.81	0.00	0.00	0.00	0.00	3.38	0.81	4.19
1996	0.70	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.82	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.76	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.71	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.75	0.00	0.93	0.00	0.00	0.00	2.66	1.79	0.00	0.00	0.00	0.00	4.34	1.79	6.13

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Charlotte County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.40	0.90	0.40	0.00	0.00	0.00	5.98	0.32	---	---	0.00	0.00	6.57	1.22	7.79
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.50	0.96	1.34	0.00	0.00	0.00	28.20	0.70	---	---	0.00	0.00	30.09	1.66	31.75
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.18	1.80	2.09	0.00	0.10	0.00	34.65	0.00	---	---	0.00	0.00	36.10	1.80	37.90
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.08	1.83	2.06	0.00	0.10	0.00	21.28	0.00	---	---	0.00	0.00	22.69	1.83	24.52
1978	0.00	2.03	2.07	0.00	0.00	0.00	26.90	0.80	---	---	0.00	0.00	27.08	2.83	29.91
1979	0.00	2.06	1.97	0.00	0.00	0.00	35.53	1.09	---	---	0.00	0.00	36.22	3.15	39.37
1980	0.00	2.41	2.49	0.00	0.00	0.00	22.84	2.49	---	---	0.00	0.00	23.73	4.90	28.63
1981	0.00	2.42	3.41	0.00	0.00	0.00	41.67	4.59	---	---	0.00	0.00	42.66	7.01	49.67
1982	2.32	2.70	3.35	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1983	2.35	2.48	3.43	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1984	2.79	2.74	3.60	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1985	2.64	2.72	5.28	0.00	0.00	0.00	41.99	3.47	1.13	0.30	0.00	0.00	48.74	6.49	55.23
1986	2.72	2.75	4.03	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	3.03	3.17	4.26	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	3.10	3.02	5.91	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	2.02	3.71	9.84	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	2.62	3.44	9.73	0.00	4.25	0.00	35.07	1.79	0.39	1.49	0.00	0.00	43.95	6.72	50.67
1991	3.07	3.07	8.08	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	2.84	3.28	10.28	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	2.93	3.19	8.53	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	2.66	3.41	6.02	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	2.44	5.12	6.49	5.85	0.33	6.19	27.46	0.96	0.09	1.44	0.00	0.00	36.17	13.71	49.88
1996	2.73	3.37	6.92	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	2.82	3.38	6.47	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	3.42	3.28	7.07	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	3.34	3.57	7.19	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	3.29	3.99	7.72	3.55	0.00	0.15	28.17	19.02	0.79	2.69	0.00	0.00	35.95	28.04	63.99

Charlotte County is located within the South Florida and the Southwest Florida Water Management Districts. The values shown represent the County totals as reported by the Districts.

Public supply transfers represent water withdrawn in DeSoto County and imported for use in Charlotte County.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Citrus County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.70	0.00	0.48	0.00	0.20	0.00	0.51	1.21	---	---	0.00	0.00	1.89	1.21	3.10
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.20	0.00	2.00	0.00	0.20	0.00	4.90	0.04	---	---	0.02	0.00	7.32	0.04	7.36
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.59	0.00	3.24	0.00	1.32	0.00	0.38	0.24	---	---	0.63	0.00	6.16	0.24	6.40
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.66	0.00	3.60	0.00	1.32	0.00	1.23	0.37	---	---	0.63	0.00	7.44	0.37	7.81
1978	0.50	0.00	4.01	0.00	2.50	0.00	2.44	0.60	---	---	0.65	0.00	10.10	0.60	10.70
1979	0.90	0.00	3.95	0.00	2.51	0.00	4.90	0.76	---	---	0.57	0.00	12.83	0.76	13.59
1980	0.90	0.00	5.20	0.00	0.34	0.21	4.58	0.23	---	---	0.57	0.00	11.59	0.44	12.03
1981	1.00	0.00	5.60	0.00	0.20	0.00	7.31	0.38	---	---	0.71	0.00	14.82	0.38	15.20
1982	4.98	0.00	1.71	0.00	0.05	0.00	7.04	0.42	---	---	0.71	0.00	14.49	0.42	14.91
1983	5.54	0.00	3.70	0.00	0.09	0.00	6.79	0.41	---	---	0.75	0.00	16.87	0.41	17.28
1984	6.03	0.00	5.46	0.00	0.41	0.00	8.67	0.53	---	---	1.14	0.00	21.71	0.53	22.24
1985	6.55	0.00	7.83	0.00	0.85	0.00	7.50	0.70	1.10	0.00	1.37	0.00	25.20	0.70	25.90
1986	7.30	0.00	5.08	0.00	2.26	0.00	9.70	0.59	---	---	1.51	0.00	25.85	0.59	26.44
1987	7.79	0.00	5.48	0.00	2.57	0.00	6.71	0.46	---	---	1.55	0.00	24.10	0.46	24.56
1988	8.78	0.00	8.18	0.00	3.08	0.00	2.41	2.22	---	---	1.72	0.00	24.17	2.22	26.39
1989	8.74	0.00	8.59	0.00	2.36	0.00	4.17	0.26	---	---	1.63	0.00	25.49	0.26	25.75
1990	8.84	0.00	9.20	0.00	2.13	0.00	1.51	0.14	1.83	0.13	1.50	0.00	25.01	0.27	25.28
1991	9.92	0.00	6.78	0.00	0.38	4.59	2.39	0.88	2.35	1.97	0.71	0.00	22.53	7.44	29.97
1992	11.73	0.00	6.94	0.00	3.00	1.46	2.52	0.72	4.04	1.92	0.63	0.00	28.86	4.10	32.96
1993	10.53	0.00	9.38	0.00	3.75	0.66	1.76	0.14	4.85	1.19	0.72	0.00	30.99	1.99	32.98
1994	9.40	0.00	7.34	0.00	4.85	0.49	1.38	0.10	4.47	1.17	0.71	0.00	28.15	1.76	29.91
1995	10.08	0.00	6.74	0.00	5.31	0.43	1.24	0.05	3.39	1.25	1.52	0.00	28.28	1.73	30.01
1996	11.03	0.00	8.66	0.00	5.07	2.73	1.36	0.10	3.41	2.27	0.75	0.00	30.28	5.10	35.38
1997	11.36	0.00	7.02	0.00	6.04	3.38	1.33	0.08	4.53	1.62	0.64	0.00	30.92	5.08	36.00
1998	14.99	0.00	9.01	0.00	5.05	1.08	1.60	0.11	3.67	1.75	0.78	0.00	35.10	2.94	38.04
1999	13.25	0.00	6.49	0.00	0.89	0.21	1.51	0.09	4.82	0.63	0.72	0.00	27.68	0.93	28.61
2000	13.97	0.00	7.20	0.00	0.80	0.29	2.76	0.59	3.77	0.41	1.55	0.00	30.05	1.29	31.34

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1990-1994, and 1996-1999 - Southwest Florida Water Management District, published water-use data, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Clay County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.00	0.00	0.55	0.00	1.50	0.00	5.32	0.00	---	---	0.00	0.00	8.37	0.00	8.37
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	1.60	0.00	2.33	0.00	1.50	0.00	4.10	0.00	---	---	0.00	0.00	9.53	0.00	9.53
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	5.01	0.00	1.81	0.00	6.62	4.30	0.48	0.04	---	---	0.00	0.00	13.92	4.34	18.26
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	4.99	0.00	1.90	0.00	4.69	4.30	1.86	1.37	---	---	0.00	0.00	13.44	5.67	19.11
1978	5.33	0.00	5.08	0.00	3.20	0.00	2.05	1.29	---	---	0.00	0.00	15.66	1.29	16.95
1979	5.24	0.00	2.49	0.00	5.14	0.00	3.24	1.27	---	---	0.00	0.00	16.11	1.27	17.38
1980	6.12	0.00	7.44	0.00	3.20	0.00	2.29	1.30	---	---	0.00	0.00	19.05	1.30	20.35
1981	7.40	0.00	4.43	0.00	5.57	0.00	2.88	1.27	---	---	0.00	0.00	20.28	1.27	21.55
1982	5.93	0.00	3.58	0.00	5.52	0.00	2.80	1.27	---	---	0.00	0.00	17.83	1.27	19.10
1983	6.39	0.00	2.34	0.00	5.58	0.00	2.12	1.27	---	---	0.00	0.00	16.43	1.27	17.70
1984	6.96	0.00	4.07	0.00	5.42	0.00	2.00	1.40	---	---	0.00	0.00	18.45	1.40	19.85
1985	8.40	0.00	2.70	0.00	7.31	4.43	2.42	0.63	0.43	0.25	0.00	0.00	21.26	5.31	26.57
1986	9.23	0.00	3.31	0.00	7.05	4.43	1.82	0.00	1.20	0.43	0.00	0.00	22.61	4.86	27.47
1987	9.92	0.00	3.64	0.00	6.98	0.00	1.78	0.00	1.09	0.38	0.00	0.00	23.41	0.38	23.79
1988	10.39	0.00	3.87	0.00	6.99	0.00	1.81	0.00	0.94	0.31	0.00	0.00	24.00	0.31	24.31
1989	10.23	0.00	4.05	0.00	7.03	0.00	1.74	0.00	0.85	0.26	0.00	0.00	23.90	0.26	24.16
1990	11.11	0.00	4.53	0.00	6.56	0.00	1.84	0.00	1.16	0.44	0.00	0.00	25.20	0.44	25.64
1991	10.61	0.00	4.99	0.00	7.28	0.00	1.56	0.00	1.00	0.38	0.00	0.00	25.44	0.38	25.82
1992	10.42	0.00	4.97	0.00	5.45	0.00	1.60	0.00	0.39	0.13	0.00	0.00	22.83	0.13	22.96
1993	9.43	0.00	4.50	0.00	6.22	0.00	0.54	0.00	0.70	0.36	0.00	0.00	21.39	0.36	21.75
1994	11.14	0.00	5.74	0.00	4.99	0.00	0.54	0.00	0.68	0.36	0.00	0.00	23.09	0.36	23.45
1995	12.04	0.00	3.59	0.00	4.46	0.00	0.47	0.00	0.72	0.24	0.00	0.00	21.28	0.24	21.52
1996	12.19	0.00	2.33	0.00	5.02	0.00	1.12	0.00	0.86	0.31	0.00	0.00	21.52	0.31	21.83
1997	11.45	0.00	2.44	0.00	4.36	0.00	1.01	0.00	0.88	0.31	0.00	0.00	20.14	0.31	20.45
1998	13.49	0.00	3.63	0.00	2.74	0.00	1.10	0.00	0.96	0.35	0.00	0.00	21.92	0.35	22.27
1999	14.54	0.00	3.99	0.00	2.68	0.00	1.10	0.00	0.86	0.45	0.00	0.00	23.17	0.45	23.62
2000	14.77	0.00	4.24	0.00	6.87	0.00	5.92	0.04	1.77	0.48	0.00	0.00	33.57	0.52	34.09

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Collier County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	2.00	1.30	0.00	0.00	0.50	0.00	48.36	4.84	----	----	0.00	0.00	52.16	6.14	58.30
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	5.00	0.90	0.00	0.00	0.50	0.00	47.10	0.40	----	----	0.00	0.00	53.62	1.30	54.92
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	11.93	0.00	0.00	0.75	0.00	0.00	64.77	5.00	----	----	0.00	0.00	77.10	5.75	82.85
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	14.12	0.00	0.00	0.45	0.00	0.00	72.93	4.82	----	----	0.00	0.00	87.66	5.27	92.93
1978	15.60	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	19.30	0.00	0.00	2.28	2.33	0.00	83.61	3.61	----	----	0.00	0.00	107.52	3.61	111.13
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	21.08	4.30	N/A	3.32	4.67	0.00	77.99	5.47	6.86	0.00	0.00	0.00	113.92	9.77	123.69
1986	23.90	4.50	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	24.69	4.61	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	27.54	4.96	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	28.32	4.96	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	31.44	5.27	N/A	8.89	0.58	3.72	143.91	5.59	10.48	3.91	0.00	0.00	195.30	18.49	213.79
1991	29.18	5.35	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	31.77	4.70	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	32.98	4.03	N/A	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	37.67	4.31	8.03	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	36.46	2.84	8.55	5.57	0.15	3.07	131.26	11.47	13.54	3.84	0.00	0.00	186.98	21.22	208.20
1996	38.19	3.50	10.59	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	39.17	4.42	9.35	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	44.27	5.09	10.21	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	44.11	4.71	11.69	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	47.17	5.23	22.46	2.67	0.19	5.62	137.18	5.81	15.28	10.89	0.00	0.00	202.49	27.55	230.04

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, Florida.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Columbia County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.20	0.00	0.60	0.00	0.00	0.00	0.40	0.13	----	----	0.00	0.00	2.20	0.13	2.33
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	1.70	0.00	1.04	0.00	0.00	0.00	0.20	0.00	----	----	0.00	0.00	2.94	0.00	2.94
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	1.70	0.00	1.33	0.00	0.12	0.00	1.29	0.28	----	----	0.00	0.00	4.44	0.28	4.72
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	1.84	0.00	1.73	0.00	0.31	0.00	1.52	0.17	----	----	0.00	0.00	5.40	0.17	5.57
1978	2.04	0.00	3.32	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	2.01	0.00	3.76	0.00	0.27	0.00	1.16	0.82	----	----	0.00	0.00	7.20	0.82	8.02
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	1.99	0.00	4.86	0.00	0.09	0.00	2.60	0.05	0.27	0.00	0.00	0.00	9.81	0.05	9.86
1986	2.25	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	2.23	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	2.35	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	2.73	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	2.93	0.00	5.21	0.00	0.03	0.00	2.78	0.23	0.20	0.00	0.00	0.00	11.15	0.23	11.38
1991	2.60	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	2.88	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	2.99	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	2.64	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	2.87	0.00	2.66	0.00	0.23	0.00	8.37	0.59	0.15	0.00	0.00	0.00	14.28	0.59	14.87
1996	3.15	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	3.17	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	3.46	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	3.55	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	3.67	0.00	3.74	0.00	0.34	0.00	5.79	0.10	0.34	0.11	0.00	0.00	13.88	0.21	14.09

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in DeSoto County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.30	0.40	0.40	0.00	1.98	1.73	20.47	2.25	---	---	0.00	0.00	23.15	4.38	27.53
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.50	1.34	1.34	0.00	0.70	0.00	64.70	0.00	---	---	0.00	0.00	66.75	1.34	68.09
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.76	2.09	2.09	0.00	0.59	0.00	64.72	2.00	---	---	0.00	0.00	67.19	4.09	71.28
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.74	2.06	2.06	0.00	1.10	0.00	63.10	1.93	---	---	0.00	0.00	66.06	3.99	70.05
1978	0.70	2.07	2.07	0.00	0.50	0.00	39.46	1.20	---	---	0.00	0.00	41.91	3.27	45.18
1979	0.70	1.97	1.97	0.00	0.72	0.00	42.29	1.28	---	---	0.00	0.00	44.92	3.25	48.17
1980	0.70	2.49	2.49	0.00	0.53	0.00	34.26	1.03	---	---	0.00	0.00	36.69	3.52	40.21
1981	0.90	3.41	3.41	0.00	0.40	0.00	59.43	1.81	---	---	0.00	0.00	61.93	5.22	67.15
1982	0.79	3.31	3.30	0.00	0.06	0.00	43.82	1.36	---	---	0.00	0.00	45.96	4.67	50.63
1983	0.80	3.43	3.43	0.00	0.14	0.00	43.38	1.34	---	---	0.00	0.00	45.48	4.77	50.25
1984	0.72	3.61	3.60	0.00	0.34	0.00	72.23	2.23	---	---	0.00	0.00	75.48	5.84	81.32
1985	0.68	5.28	5.28	0.00	0.37	0.00	71.65	2.27	0.55	0.00	0.00	0.00	74.93	7.55	82.48
1986	0.61	4.04	4.03	0.00	0.28	0.00	63.54	1.97	---	---	0.00	0.00	66.59	6.01	72.60
1987	0.79	4.28	4.26	0.00	0.25	0.00	61.96	2.92	---	---	0.00	0.00	64.53	7.20	71.73
1988	0.86	5.93	5.91	0.00	0.25	0.00	47.21	2.71	---	---	0.00	0.00	49.94	8.64	58.58
1989	1.08	10.18	10.41	0.00	0.27	0.00	67.72	7.58	---	---	0.00	0.00	70.95	17.76	88.71
1990	1.63	9.19	9.98	0.00	0.26	0.22	97.74	10.51	0.53	0.00	0.00	0.00	102.05	19.92	121.97
1991	0.96	8.26	8.26	0.00	0.01	0.34	72.07	2.36	0.04	0.29	0.00	0.00	74.83	11.25	86.08
1992	0.75	8.47	8.47	0.00	0.01	0.72	66.07	1.52	0.04	0.30	0.00	0.00	68.51	11.01	79.52
1993	0.62	8.76	8.53	0.00	0.01	0.45	56.67	0.53	0.12	0.30	0.00	0.00	59.87	10.04	69.91
1994	1.05	9.29	9.28	0.00	0.24	0.78	52.31	0.56	0.16	0.20	0.00	0.00	55.98	10.83	66.81
1995	1.97	10.51	10.54	0.00	0.02	0.39	54.15	1.28	0.17	0.17	0.00	0.00	58.59	12.35	70.94
1996	2.96	9.68	11.26	0.00	0.15	0.85	63.54	1.04	0.19	0.22	0.00	0.00	69.07	11.79	80.86
1997	2.30	10.19	11.39	0.00	0.09	0.04	63.97	0.65	0.26	0.18	0.00	0.00	68.84	11.06	79.90
1998	1.84	12.60	12.54	0.00	0.04	0.18	69.04	0.96	0.41	0.08	0.00	0.00	72.88	13.82	86.70
1999	2.54	10.63	11.88	0.00	0.04	0.19	75.13	0.19	0.22	0.14	0.00	0.00	79.69	11.15	90.84
2000	4.49	6.10	8.63	0.00	0.06	1.33	116.95	1.92	0.21	0.16	0.00	0.00	123.87	9.51	133.38

Public supply transfers represents water withdrawn in DeSoto County and exported for use in Charlotte and Sarasota Counties and includes transfer losses.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1990-1994, and 1996-1999 - published water-use data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Duval County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	60.00	0.00	0.00	0.00	47.00	1.00	0.04	0.09	----	----	2.80	0.00	127.24	1.09	128.33
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	67.80	0.00	36.20	0.00	60.90	0.00	3.70	0.00	----	----	0.30	0.00	168.90	0.00	168.90
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	73.50	0.00	32.90	0.00	48.63	0.14	2.33	0.22	----	----	2.10	0.00	159.46	0.36	159.82
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	69.90	0.00	27.60	0.00	48.23	0.05	2.31	2.29	----	----	1.87	0.00	149.91	2.34	152.25
1978	59.56	0.00	23.44	0.00	42.98	0.00	0.42	1.65	----	----	2.63	0.00	129.03	1.65	130.68
1979	74.60	0.00	27.70	0.00	35.80	0.00	7.28	2.25	----	----	2.65	0.00	148.03	2.25	150.28
1980	73.25	0.00	20.26	0.00	40.46	0.00	4.79	1.39	----	----	2.65	0.00	141.41	1.39	142.80
1981	74.71	0.00	23.85	0.00	39.20	0.00	4.97	2.34	----	----	2.48	0.00	145.21	2.34	147.55
1982	70.19	0.00	17.08	0.00	35.81	0.00	4.62	2.26	----	----	2.16	0.00	129.86	2.26	132.12
1983	70.45	0.00	18.05	0.00	38.69	0.00	4.72	2.29	----	----	1.98	0.00	133.89	2.29	136.18
1984	77.55	0.00	19.17	0.00	40.07	0.00	4.62	2.37	----	----	2.79	0.00	144.20	2.37	146.57
1985	84.86	0.00	15.50	0.00	38.08	0.00	7.04	0.39	3.90	0.98	2.06	0.00	151.44	1.37	152.81
1986	89.20	0.00	16.62	0.00	22.82	0.00	4.93	0.24	4.24	0.98	3.09	0.00	140.90	1.22	142.12
1987	94.28	0.00	17.27	0.00	37.75	0.00	4.98	0.12	3.75	0.86	3.57	0.00	161.60	0.98	162.58
1988	93.44	0.00	15.55	0.00	40.11	0.00	4.95	0.10	3.74	0.86	3.81	0.00	161.60	0.96	162.56
1989	94.06	0.00	13.41	0.00	39.89	0.00	5.87	0.27	3.35	0.75	4.44	0.00	161.02	1.02	162.04
1990	96.32	0.00	8.37	0.00	33.93	0.00	5.32	0.42	4.21	0.98	4.83	0.00	152.98	1.40	154.38
1991	92.55	0.00	7.56	0.00	35.69	0.00	3.22	0.07	3.62	0.83	4.77	0.00	147.41	0.90	148.31
1992	94.80	0.00	8.35	0.00	31.17	0.00	2.93	0.05	1.02	0.22	4.42	0.00	142.69	0.27	142.96
1993	99.78	0.00	8.35	0.00	23.41	0.00	1.93	0.12	3.48	0.84	5.00	0.00	141.95	0.96	142.91
1994	99.09	0.00	8.66	0.00	24.56	0.00	1.92	0.12	3.41	0.83	4.62	0.00	142.26	0.95	143.21
1995	99.62	0.00	11.87	0.00	24.75	0.00	0.88	0.05	2.02	0.43	5.47	0.00	144.61	0.48	145.09
1996	106.81	0.00	11.50	0.00	24.15	0.00	1.09	0.08	2.97	0.65	4.54	0.00	151.06	0.73	151.79
1997	104.93	0.00	11.18	0.00	25.83	0.00	0.94	0.06	3.03	0.67	5.44	0.00	151.35	0.73	152.08
1998	107.43	0.00	11.57	0.00	22.02	0.00	1.30	0.09	3.34	0.73	5.28	0.00	150.94	0.82	151.76
1999	112.47	0.00	13.88	0.00	16.37	0.00	0.92	0.06	2.76	0.60	5.54	0.00	151.94	0.66	152.60
2000	119.12	0.00	4.46	0.00	12.51	0.00	3.74	0.33	6.17	1.32	8.33	0.00	154.33	1.65	155.98

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Escambia County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	17.40	0.00	0.00	0.00	41.20	39.60	0.10	0.10	---	---	0.50	218.00	62.20	257.70	319.90
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	20.30	0.00	0.00	0.00	47.80	42.90	0.10	0.00	---	---	0.80	205.00	74.63	247.90	322.53
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	27.46	0.00	0.00	0.00	44.75	31.70	0.44	0.44	---	---	2.52	265.40	81.66	297.54	379.20
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	30.48	0.00	0.00	0.00	47.36	25.60	0.77	0.39	---	---	2.50	249.70	85.29	275.69	360.98
1978	29.72	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	30.53	0.00	0.00	0.00	34.47	20.51	1.17	1.32	---	---	2.50	335.60	72.62	357.43	430.05
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	33.07	0.00	0.00	0.00	35.54	16.89	1.42	0.48	3.80	0.00	2.20	241.40	80.74	258.77	339.51
1986	34.45	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	35.20	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	36.22	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	35.19	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	36.88	0.00	0.00	0.00	32.39	19.34	1.53	0.14	4.70	0.00	2.20	190.43	81.36	209.91	291.27
1991	33.44	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	35.50	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	36.54	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	34.81	0.00	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	37.73	0.00	0.62	0.00	37.20	22.19	1.66	0.31	4.74	0.62	2.11	159.60	86.83	182.72	269.55
1996	39.10	0.00	1.11	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	39.18	0.00	1.15	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	42.32	0.00	1.17	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	41.09	0.00	1.05	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	44.63	0.00	1.07	1.60	35.77	24.65	2.73	0.00	4.79	0.77	2.42	199.89	91.94	225.31	317.25

Public supply transfers represent water withdrawn in Escambia County and exported for use in Santa Rosa County.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Flagler County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day: ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.20	0.00	0.00	0.00	0.00	0.00	3.70	0.00	----	----	0.00	0.00	4.03	0.00	4.03
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.30	0.00	0.00	0.00	0.00	0.00	9.00	0.20	----	----	0.00	0.00	9.50	0.20	9.70
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.62	0.00	0.00	0.00	0.00	0.00	8.68	0.00	----	----	0.00	0.00	9.39	0.00	9.39
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	0.62	0.00	0.00	0.00	0.00	0.00	3.45	0.05	----	----	0.00	0.00	4.35	0.05	4.40
1978	0.52	0.00	0.00	0.00	0.00	0.00	3.21	0.05	----	----	0.00	0.00	4.30	0.05	4.35
1979	1.37	0.00	0.00	0.00	0.00	0.00	8.44	0.23	----	----	0.00	0.00	10.14	0.23	10.37
1980	0.77	0.00	0.00	0.00	0.00	0.00	4.14	0.05	----	----	0.00	0.00	5.35	0.05	5.40
1981	1.70	0.00	0.00	0.00	0.04	0.00	6.96	0.17	----	----	0.00	0.00	9.24	0.17	9.41
1982	1.45	0.00	0.00	0.00	0.05	0.00	5.85	0.17	----	----	0.00	0.00	7.83	0.17	8.00
1983	1.55	0.00	0.00	0.00	0.00	0.00	6.45	0.17	----	----	0.00	0.00	8.27	0.17	8.44
1984	1.77	0.00	0.00	0.00	0.10	0.00	6.13	0.48	----	----	0.00	0.00	8.29	0.48	8.77
1985	2.22	0.00	0.00	0.00	0.10	0.00	6.12	0.05	0.19	0.88	0.00	0.00	8.94	0.93	9.87
1986	2.12	0.00	0.00	0.00	0.13	0.00	6.73	0.01	0.16	1.32	0.00	0.00	9.41	1.33	10.74
1987	2.56	0.00	0.00	0.00	0.16	0.00	5.59	0.01	0.14	1.08	0.00	0.00	8.70	1.09	9.79
1988	3.03	0.00	0.00	0.00	0.16	0.00	6.70	0.01	0.11	0.91	0.00	0.00	10.42	0.92	11.34
1989	3.34	0.00	0.00	0.00	0.16	0.00	6.64	0.00	0.12	0.96	0.00	0.00	11.02	0.96	11.98
1990	3.85	0.00	0.00	0.00	0.25	0.00	7.34	0.00	0.16	1.20	0.00	0.00	13.47	1.20	14.67
1991	3.79	0.00	0.00	0.00	0.16	0.00	5.15	0.00	0.14	1.08	0.00	0.00	11.39	1.08	12.47
1992	4.09	0.00	0.00	0.00	0.15	0.00	8.02	0.00	0.08	0.63	0.00	0.00	14.58	0.63	15.21
1993	4.07	0.00	0.00	0.00	0.17	0.00	7.01	0.15	0.13	0.89	0.00	0.00	13.02	1.04	14.06
1994	4.20	0.00	0.00	0.00	0.20	0.00	7.39	0.15	0.13	0.88	0.00	0.00	13.61	1.03	14.64
1995	4.51	0.00	0.00	0.00	0.18	0.00	6.65	0.00	0.11	0.85	0.00	0.00	13.30	0.85	14.15
1996	4.50	0.00	0.00	0.00	0.07	0.00	6.66	0.00	0.18	1.42	0.00	0.00	13.24	1.42	14.66
1997	4.93	0.00	0.00	0.00	0.07	0.00	6.07	0.00	0.18	1.44	0.00	0.00	13.21	1.44	14.65
1998	5.22	0.00	0.00	0.00	0.06	0.00	8.22	0.00	0.20	1.63	0.00	0.00	15.83	1.63	17.46
1999	6.08	0.00	0.00	0.00	0.05	0.00	9.46	0.00	0.15	1.23	0.00	0.00	18.28	1.23	19.51
2000	6.22	0.00	0.00	0.00	0.20	0.07	15.67	0.03	1.84	3.50	0.00	0.00	24.55	3.60	28.15

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Franklin County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.50	0.00	0.10	0.00	0.20	0.00	0.11	0.01	----	----	0.00	0.00	0.91	0.01	0.92
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.50	0.00	0.37	0.00	0.00	0.00	0.00	0.00	----	----	0.00	0.00	0.87	0.00	0.87
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.99	0.00	0.12	0.01	0.00	0.00	0.00	0.01	----	----	0.00	0.00	1.11	0.02	1.13
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	1.11	0.00	0.20	0.00	0.01	0.00	0.01	0.01	----	----	0.00	0.00	1.33	0.01	1.34
1978	1.08	0.00	0.18	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	1.00	0.00	0.29	0.00	0.01	0.00	0.00	0.00	----	----	0.00	0.00	1.30	0.00	1.30
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	1.20	0.00	0.15	0.00	0.00	0.00	0.77	0.00	0.86	0.00	0.00	0.00	2.98	0.00	2.98
1986	1.54	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	1.52	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	1.46	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	1.58	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	1.63	0.00	0.19	0.00	0.00	0.00	0.37	0.11	0.37	0.00	0.00	0.00	2.56	0.11	2.67
1991	1.46	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	1.56	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	1.76	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1.74	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1.75	0.00	0.40	0.00	0.00	0.00	0.17	0.00	0.57	0.00	0.00	0.00	2.89	0.00	2.89
1996	1.81	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	1.66	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	1.85	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	1.79	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	1.92	0.00	0.19	0.00	0.00	0.00	0.01	0.00	0.13	0.00	0.00	0.00	2.25	0.00	2.25

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Gadsden County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.60	1.20	0.00	0.00	0.20	2.10	0.80	4.90	---	---	0.00	0.00	2.70	8.20	10.90
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.80	1.20	0.00	0.00	0.10	2.10	0.50	2.10	---	---	0.00	0.00	4.06	5.40	9.46
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.96	1.18	0.00	0.10	0.09	1.94	0.03	2.73	---	---	0.00	0.00	3.05	5.95	9.00
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.08	1.48	0.00	0.10	0.09	1.95	0.38	4.63	---	---	0.00	0.00	3.63	8.16	11.79
1978	1.12	1.30	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	1.22	1.02	0.00	0.08	0.14	1.70	0.32	6.38	---	---	0.00	0.00	4.30	9.18	13.48
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	1.60	1.13	0.00	0.00	1.96	0.00	2.59	6.19	0.00	0.35	0.00	0.00	8.45	7.67	16.12
1986	1.87	1.28	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	1.93	1.43	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	2.10	1.45	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	2.05	1.25	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	2.13	1.29	0.00	0.00	2.16	0.00	2.99	7.43	0.25	0.26	0.00	0.00	9.53	8.98	18.51
1991	2.16	1.18	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	2.14	1.24	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	2.25	1.29	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	2.12	1.17	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	2.43	1.43	0.00	0.00	1.04	1.81	1.76	5.33	0.11	0.10	0.00	0.00	7.71	8.67	16.38
1996	2.47	1.39	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	2.72	1.45	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	2.82	1.47	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	2.76	1.54	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	3.06	1.28	0.00	1.85	0.00	0.92	0.00	2.95	0.15	0.15	0.00	0.00	8.93	7.40	16.33

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Glades County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.10	0.00	0.10	0.01	0.00	0.00	0.35	9.25	---	---	0.00	0.00	0.55	9.26	9.81
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.20	0.00	0.26	0.00	0.40	0.00	5.20	41.10	---	---	0.00	0.00	6.06	41.10	47.16
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.20	0.00	0.40	0.00	0.00	0.00	11.46	42.01	---	---	0.00	0.00	12.06	42.01	54.07
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.22	0.00	0.42	0.00	0.66	0.00	12.41	46.92	---	---	0.00	0.00	13.71	46.92	60.63
1978	0.21	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	0.21	0.00	0.72	0.00	0.00	0.00	17.46	107.31	---	---	0.00	0.00	18.39	107.31	125.70
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.25	0.00	0.56	0.00	1.48	0.00	13.67	66.12	0.00	0.00	0.00	0.00	15.96	66.12	82.08
1986	0.36	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.26	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.40	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.42	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.40	0.00	0.58	0.00	0.27	0.00	17.91	61.88	0.00	0.67	0.00	0.00	19.16	62.55	81.71
1991	0.37	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.34	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.36	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.36	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.38	0.00	0.56	0.00	0.36	14.53	19.65	64.04	0.00	0.07	0.00	0.00	20.95	78.64	99.59
1996	0.39	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.73	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.79	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.78	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.55	0.00	0.61	0.00	0.14	3.93	19.71	49.31	0.02	0.40	0.00	0.00	21.03	53.64	74.67

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Gulf County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.10	0.30	0.00	0.00	0.00	24.90	0.03	0.00	----	----	0.00	0.00	0.33	25.20	25.53
1966	----	----	----	----	----	----	----	----	----	----	0.00	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.10	0.40	0.00	0.00	0.20	35.73	0.00	0.00	----	----	0.00	0.00	0.79	36.13	36.92
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.11	0.64	0.00	0.00	0.52	33.20	0.26	0.08	----	----	0.00	0.00	1.32	33.92	35.24
1976	----	----	----	----	----	----	----	----	----	----	0.00	----	----	----	----
1977	0.16	0.72	0.00	0.00	0.52	33.20	0.24	0.04	----	----	0.00	0.00	1.48	33.96	35.44
1978	0.17	0.82	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	0.96	0.00	0.00	0.00	0.54	33.40	0.02	0.04	----	----	0.00	0.00	2.03	33.44	35.47
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	0.98	0.00	0.00	0.00	0.42	31.68	1.34	2.17	0.55	0.00	0.00	0.00	4.03	33.85	37.88
1986	1.10	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	1.07	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	1.15	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	1.28	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	1.26	0.00	0.00	0.34	0.58	32.89	4.72	5.17	0.21	0.00	0.00	0.00	7.11	38.06	45.17
1991	1.10	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	1.13	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	1.30	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1.20	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1.28	0.00	0.00	0.40	0.68	27.98	0.58	0.00	0.17	0.00	0.00	0.00	3.11	27.98	31.09
1996	1.35	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	1.36	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	1.41	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	1.00	0.40	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	1.47	0.00	0.00	0.32	0.90	1.65	0.21	0.00	0.20	0.00	0.00	0.00	3.10	1.65	4.75

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Hamilton County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.20	0.00	0.00	0.00	10.30	0.00	0.83	0.28	---	---	0.00	0.00	11.58	0.28	11.86
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.50	0.00	0.00	0.00	18.40	0.00	0.50	0.20	---	---	0.00	0.00	19.82	0.20	20.02
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.60	0.00	0.00	0.00	30.30	0.00	1.75	0.22	---	---	0.00	0.00	32.93	0.22	33.15
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.81	0.00	0.00	0.00	28.80	0.00	1.79	0.40	---	---	0.00	0.00	31.78	0.40	32.18
1978	0.76	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	0.68	0.00	0.00	0.00	35.82	0.00	0.71	0.22	---	---	0.00	0.00	37.78	0.22	38.00
1981	0.62	0.00	0.00	0.00	41.56	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.73	0.00	0.00	0.00	38.79	0.00	2.71	0.00	0.20	0.00	0.00	0.00	43.38	0.00	43.38
1986	0.77	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.83	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.90	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.92	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.97	0.00	0.00	0.00	44.08	0.00	4.41	0.00	0.03	0.00	0.00	0.00	50.60	0.00	50.60
1991	0.86	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.82	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.93	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.87	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.87	0.00	0.00	0.00	39.64	0.00	4.80	0.00	0.03	0.00	0.00	0.00	46.18	0.00	46.18
1996	0.89	0.00	0.00	0.00	34.58	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.82	0.00	0.00	0.00	34.05	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.84	0.00	0.00	0.00	32.68	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.87	0.00	0.00	0.00	34.90	0.00	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.95	0.00	0.00	0.00	34.39	0.00	5.53	0.11	0.00	0.00	0.00	0.00	41.61	0.11	41.72

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Hardee County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.80	0.00	0.00	0.00	0.60	0.00	42.24	2.24	---	---	0.00	0.00	43.96	2.24	46.20
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.70	0.00	1.01	0.00	0.10	0.00	62.50	0.00	---	---	0.00	0.00	64.31	0.00	64.31
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	1.20	0.00	1.16	0.00	1.45	0.00	93.30	0.00	---	---	0.00	0.00	97.11	0.00	97.11
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.02	0.00	1.05	0.00	9.94	0.00	57.35	0.00	---	---	0.00	0.00	69.36	0.00	69.36
1978	0.80	0.00	1.33	0.00	11.00	0.00	49.30	0.00	---	---	0.00	0.00	62.43	0.00	62.43
1979	1.20	0.00	1.09	0.00	16.00	0.00	35.99	0.00	---	---	0.00	0.00	54.28	0.00	54.28
1980	1.30	0.00	1.19	0.00	0.80	0.00	40.29	0.00	---	---	0.00	0.00	43.58	0.00	43.58
1981	1.50	0.00	1.11	0.00	0.25	0.00	58.77	0.00	---	---	0.00	0.00	61.63	0.00	61.63
1982	1.36	0.00	1.12	0.00	2.91	0.00	41.17	0.00	---	---	0.00	0.00	46.56	0.00	46.56
1983	1.03	0.00	1.30	0.00	0.11	0.00	44.51	0.00	---	---	0.00	0.00	46.95	0.00	46.95
1984	1.29	0.00	1.72	0.00	4.69	0.00	81.28	0.00	---	---	0.00	0.00	88.98	0.00	88.98
1985	1.32	0.00	1.82	0.00	4.83	0.00	84.78	0.00	1.43	0.00	0.00	0.00	94.18	0.00	94.18
1986	1.30	0.00	1.82	0.00	7.13	0.00	60.03	0.00	---	---	0.00	0.00	70.28	0.00	70.28
1987	1.35	0.00	1.99	0.00	7.61	0.00	43.53	1.86	---	---	0.00	0.00	54.48	1.86	56.34
1988	1.44	0.00	1.96	0.00	0.29	0.00	63.58	1.91	---	---	0.00	0.00	67.27	1.91	69.18
1989	1.48	0.00	2.04	0.00	0.10	0.00	57.24	6.36	---	---	0.00	0.00	60.86	6.36	67.22
1990	1.43	0.00	1.93	0.00	0.10	0.00	57.05	5.95	0.09	0.00	0.00	0.00	60.60	5.95	66.55
1991	1.52	0.00	1.30	0.00	1.04	0.00	55.63	2.28	0.00	0.00	0.00	0.00	59.49	2.28	61.77
1992	1.46	0.00	1.21	0.00	0.80	0.01	58.97	2.43	0.00	0.00	0.00	0.00	62.44	2.44	64.88
1993	1.60	0.00	1.30	0.00	0.07	0.41	50.74	0.00	0.02	0.00	0.00	0.00	53.73	0.41	54.14
1994	1.44	0.00	1.28	0.00	0.06	0.01	50.64	0.34	0.10	0.00	0.00	0.00	53.52	0.35	53.87
1995	1.61	0.00	1.77	0.00	2.81	0.00	43.93	0.51	0.12	0.00	0.25	0.00	50.49	0.51	51.00
1996	1.63	0.00	1.49	0.00	6.92	0.00	57.89	0.59	0.13	0.00	0.00	0.00	68.06	0.59	68.65
1997	1.57	0.00	0.63	0.00	2.32	0.00	54.92	0.42	0.16	0.00	0.00	0.00	59.60	0.42	60.02
1998	1.74	0.00	0.58	0.00	0.55	0.00	59.64	0.76	0.15	0.00	0.00	0.00	62.66	0.76	63.42
1999	1.69	0.00	0.49	0.00	4.15	0.00	62.60	0.56	0.16	0.00	0.00	0.00	69.09	0.56	69.65
2000	1.78	0.00	0.64	0.00	5.93	0.00	81.16	0.57	0.21	0.04	0.81	0.00	90.53	0.61	91.14

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Hendry County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.10	0.40	0.00	0.00	0.00	1.10	38.36	212.07	---	---	0.00	0.00	39.16	213.57	252.73
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.20	1.20	0.00	0.00	0.10	0.20	25.00	221.40	---	---	0.00	0.00	25.95	222.80	248.75
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.25	1.80	0.00	0.00	0.22	0.60	80.59	213.08	---	---	0.00	0.00	81.76	215.48	297.24
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.25	1.30	0.00	0.00	0.25	0.71	86.56	204.95	---	---	0.00	0.00	87.68	206.96	294.64
1978	0.26	1.44	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	0.38	1.62	0.00	0.00	0.23	0.00	110.42	134.77	---	---	0.00	0.00	111.87	136.39	248.26
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.51	2.38	0.00	0.00	0.50	0.00	29.77	158.06	0.00	0.00	0.00	0.00	31.56	160.44	192.00
1986	0.51	2.56	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.48	2.68	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.67	2.73	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.80	2.90	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.76	2.84	0.00	1.97	1.06	0.00	167.82	317.43	0.00	1.08	0.00	0.00	171.61	321.35	492.96
1991	0.74	3.08	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.85	2.74	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.70	2.87	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.73	2.85	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.77	3.25	0.00	1.21	0.97	0.00	149.33	400.32	0.82	0.34	0.00	0.00	153.10	403.91	557.01
1996	0.76	3.06	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.80	2.59	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.79	3.17	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.79	4.16	0.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.95	3.77	0.00	1.67	0.00	0.72	193.37	310.54	0.41	0.68	0.00	0.00	197.12	314.99	512.11

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Hernando County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.50	0.00	0.40	0.00	27.00	0.00	1.90	0.15	---	---	0.00	0.00	29.80	0.15	29.95
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.60	0.00	1.44	0.00	19.60	0.00	1.70	0.02	---	---	0.00	0.00	23.34	0.02	23.36
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.75	0.00	2.35	0.00	25.38	0.00	3.07	0.31	---	---	0.00	0.00	31.55	0.31	31.86
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.92	0.00	2.69	0.00	33.76	0.00	5.12	0.89	---	---	0.00	0.00	42.49	0.89	43.38
1978	1.00	0.00	2.88	0.00	33.80	0.00	5.03	0.80	---	---	0.00	0.00	42.71	0.80	43.51
1979	1.10	0.00	3.19	0.00	34.60	0.00	5.43	0.89	---	---	0.00	0.00	44.32	0.89	45.21
1980	1.10	0.00	3.89	0.00	31.60	0.00	4.90	0.10	---	---	0.00	0.00	41.49	0.10	41.59
1981	1.10	0.00	4.29	0.00	34.30	1.90	6.59	0.10	---	---	0.00	0.00	46.28	2.00	48.28
1982	5.59	0.00	2.23	0.00	31.36	0.00	6.96	0.00	---	---	0.00	0.00	46.14	0.00	46.14
1983	5.32	0.00	2.38	0.00	25.59	0.00	4.61	0.00	---	---	0.00	0.00	37.90	0.00	37.90
1984	6.94	0.00	3.90	0.00	28.73	0.00	5.75	0.00	---	---	0.00	0.00	45.32	0.00	45.32
1985	7.88	0.00	0.96	0.00	26.74	0.00	3.67	0.00	1.61	0.00	0.00	0.00	40.86	0.00	40.86
1986	9.08	0.00	2.00	0.00	29.02	0.00	6.76	0.00	---	---	0.00	0.00	46.86	0.00	46.86
1987	10.48	0.00	0.31	0.00	28.88	0.00	4.24	0.81	---	---	0.00	0.00	43.91	0.81	44.72
1988	12.74	0.00	0.65	0.00	24.25	0.00	1.74	2.03	---	---	0.00	0.00	39.38	2.03	41.41
1989	14.36	0.00	1.51	0.00	24.80	0.00	5.31	0.27	---	---	0.00	0.00	45.98	0.27	46.25
1990	15.01	0.00	1.71	0.00	23.31	0.00	2.36	0.35	2.40	0.00	0.00	0.00	44.79	0.35	45.14
1991	14.43	0.00	2.39	0.00	11.59	1.95	3.69	4.02	2.73	0.76	0.00	0.00	34.83	6.73	41.56
1992	16.40	0.00	2.02	0.00	13.99	1.23	3.62	3.61	4.00	0.60	0.00	0.00	40.03	5.44	45.47
1993	16.93	0.00	3.49	0.00	13.88	1.34	2.95	0.00	2.90	1.42	0.00	0.00	40.15	2.76	42.91
1994	15.85	0.01	2.93	0.00	15.10	0.36	2.61	0.00	3.33	1.42	0.00	0.00	39.82	1.79	41.61
1995	17.06	0.01	2.44	0.00	13.97	0.70	2.37	0.00	3.74	1.33	0.00	0.00	39.58	2.04	41.62
1996	16.55	0.01	1.27	0.00	11.62	1.15	2.43	0.00	4.41	1.20	0.00	0.00	36.28	2.36	38.64
1997	18.20	0.01	1.53	0.00	19.76	0.03	2.41	0.00	4.26	0.83	0.00	0.00	46.16	0.87	47.03
1998	20.54	0.01	1.22	0.00	15.12	0.00	2.65	0.02	3.53	0.50	0.00	0.00	43.06	0.53	43.59
1999	20.49	0.01	1.32	0.00	25.85	0.06	2.71	0.03	3.90	0.57	0.00	0.00	54.27	0.67	54.94
2000	20.26	0.01	1.41	0.00	19.70	0.07	4.26	0.03	3.83	0.88	0.00	0.00	49.46	0.99	50.45

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Highlands County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn	
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface
1965	4.40	0.60	0.39	0.00	0.00	0.00	31.72	21.11	----	----	0.00	123.00	36.51	144.71
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	4.10	0.60	0.73	0.00	0.10	0.00	22.40	35.40	----	----	0.02	226.00	27.35	262.00
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	4.26	0.00	1.84	0.00	0.70	0.00	86.90	58.70	----	----	0.00	95.23	93.70	153.93
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	3.73	0.00	1.66	0.00	0.70	0.00	97.53	63.36	----	----	0.00	95.23	103.62	158.59
1978	4.30	0.00	1.89	0.00	0.70	0.00	155.78	29.90	----	----	0.00	76.00	162.67	105.90
1979	4.30	0.00	2.11	0.00	0.10	0.00	42.88	9.58	----	----	0.00	76.40	49.39	85.98
1980	5.00	0.00	3.37	0.00	0.95	0.00	57.56	32.93	----	----	0.00	76.41	66.88	109.34
1981	6.84	0.00	3.46	0.00	0.30	0.00	69.88	40.28	----	----	0.00	77.00	80.48	117.28
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	7.88	0.00	2.77	0.00	0.51	0.00	87.85	14.16	1.39	0.00	0.00	9.11	100.40	23.27
1986	7.73	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1987	7.82	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1988	8.42	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1989	8.76	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1990	8.30	0.00	3.43	0.00	0.21	0.00	115.35	10.14	1.99	0.00	0.07	1.98	129.35	12.12
1991	8.61	0.18	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1992	8.42	0.21	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1993	8.75	0.21	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1994	7.72	0.08	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1995	8.27	0.06	2.86	0.00	0.50	0.85	99.06	5.62	2.14	0.08	0.00	0.03	112.83	6.64
1996	8.07	0.09	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1997	7.92	0.08	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1998	8.57	0.11	----	----	----	----	----	----	----	----	----	----	N/A	N/A
1999	8.79	0.09	----	----	----	----	----	----	----	----	----	----	N/A	N/A
2000	9.14	0.00	1.68	0.00	0.55	0.03	143.24	17.07	2.60	0.16	0.08	0.00	157.29	17.26

Highlands County is located within the South Florida and the Southwest Florida Water Management Districts. The values shown represent the County totals as reported by the Districts. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Hillsborough County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	18.80	38.20	13.20	0.00	82.00	2.00	44.73	11.21	----	----	0.00	0.00	152.03	51.41	203.44
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	35.20	44.60	28.00	0.00	40.00	11.90	64.20	5.10	----	----	0.00	0.00	153.84	61.60	215.44
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	31.44	52.70	24.27	0.00	20.80	8.10	48.50	2.27	----	----	0.00	0.00	122.00	63.07	185.07
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	41.38	56.64	27.96	0.00	26.76	7.40	52.28	2.52	----	----	0.00	0.00	129.15	66.56	195.71
1978	43.40	45.00	22.40	0.00	35.50	9.30	61.60	3.00	----	----	0.00	0.00	150.15	57.30	207.45
1979	54.21	39.00	25.11	0.00	21.40	7.90	62.54	3.02	----	----	0.00	0.00	146.29	49.92	196.21
1980	58.27	49.87	23.44	0.00	20.54	6.23	74.97	2.18	----	----	0.00	0.00	159.70	58.28	217.98
1981	63.61	58.70	22.31	0.00	15.20	7.20	91.57	3.15	----	----	0.00	0.00	176.53	69.05	245.58
1982	65.74	48.26	24.30	0.00	14.96	6.59	78.21	2.42	----	----	0.00	0.00	169.08	57.27	226.35
1983	56.77	49.48	23.07	0.00	22.57	0.07	79.77	2.47	----	----	0.07	0.00	177.49	52.02	229.51
1984	78.67	50.79	30.82	0.00	30.72	0.00	124.50	3.85	----	----	0.01	0.00	243.77	54.64	298.41
1985	81.81	64.46	32.18	0.00	34.36	0.00	103.15	2.84	4.75	0.61	0.05	0.00	230.17	67.91	298.08
1986	73.72	54.06	30.05	0.00	24.83	0.00	103.24	3.16	----	----	0.00	0.00	208.25	57.22	265.47
1987	63.64	58.96	17.44	0.00	20.47	0.00	88.52	3.07	----	----	0.00	0.00	187.30	62.03	249.33
1988	68.58	65.77	14.06	0.00	16.42	4.59	84.77	5.29	----	----	0.00	0.00	180.52	75.65	256.17
1989	68.06	68.69	7.36	0.00	22.07	4.18	96.50	12.80	----	----	0.04	0.00	201.42	85.67	287.09
1990	61.79	76.05	12.65	0.00	25.82	4.21	82.97	7.98	6.47	0.70	0.00	0.00	179.73	88.94	268.67
1991	56.78	77.28	13.20	0.00	22.46	9.33	73.74	19.99	3.75	3.58	0.00	0.00	159.59	110.18	269.77
1992	65.98	80.89	22.40	0.00	20.02	10.12	72.15	15.82	3.14	3.77	0.00	0.00	165.53	110.60	276.13
1993	63.13	68.07	17.23	0.00	36.83	5.62	64.74	3.15	6.49	4.11	0.00	0.00	185.04	80.95	265.99
1994	64.92	65.80	22.26	0.00	32.51	5.59	62.79	3.24	5.04	2.85	0.00	0.00	178.67	77.48	256.15
1995	63.75	65.75	15.67	0.00	27.60	5.54	60.39	2.90	5.11	2.93	0.00	0.00	169.22	77.12	246.34
1996	67.53	67.78	20.34	0.00	12.36	5.61	69.66	3.04	6.81	4.15	0.00	0.00	167.92	80.58	248.50
1997	77.16	74.52	19.80	0.00	12.24	4.42	63.18	3.15	9.38	4.39	0.00	0.00	172.35	86.48	258.83
1998	80.01	77.35	21.35	0.00	11.72	5.11	64.67	2.99	7.69	4.75	0.00	0.00	166.66	90.20	256.86
1999	83.91	86.67	29.57	0.00	10.28	5.60	68.25	3.28	7.48	4.25	0.00	0.00	175.57	99.80	275.37
2000	85.51	80.88	11.30	0.00	14.17	5.50	83.84	3.31	9.21	4.59	0.00	0.00	197.44	94.28	291.72

Public supply transfers represent water withdrawn in Hillsborough County and exported for use in Pinellas County and includes transfer losses.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Holmes County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.20	0.00	0.00	0.00	0.00	0.00	0.00	0.15	----	----	0.00	0.00	0.70	0.15	0.85
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.30	0.00	0.00	0.00	0.00	0.00	0.04	0.00	----	----	0.00	0.00	1.26	0.00	1.26
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.20	0.00	0.00	0.00	0.02	0.00	0.14	0.33	----	----	0.00	0.00	1.21	0.33	1.54
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	0.60	0.00	0.00	0.00	0.02	0.00	0.28	0.27	----	----	0.00	0.00	2.06	0.27	2.33
1978	0.66	0.00	0.00	0.00	0.00	0.00	0.00	0.00	----	----	0.00	0.00	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	0.64	0.00	0.00	0.00	0.06	0.00	0.41	0.41	----	----	0.00	0.00	2.38	0.41	2.79
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	0.82	0.00	0.00	0.00	0.00	0.00	2.20	0.43	0.00	0.00	0.00	0.00	5.04	0.43	5.47
1986	0.85	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1987	0.88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1988	0.90	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1989	0.92	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1990	1.10	0.00	0.00	0.00	0.00	0.00	2.21	1.16	0.00	0.00	0.00	0.00	5.60	1.16	6.76
1991	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1992	1.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1993	1.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1994	1.22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1995	1.18	0.00	0.00	0.00	0.00	0.00	2.70	0.71	0.00	0.00	0.00	0.00	6.53	0.71	7.24
1996	1.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1997	1.15	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1998	1.26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1999	1.28	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
2000	1.38	0.00	0.00	0.00	0.00	0.00	0.78	0.19	0.26	0.13	0.00	0.00	3.77	0.32	4.09

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Indian River County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.50	0.00	1.00	0.00	0.00	0.00	26.45	36.68	---	---	0.01	0.00	28.96	36.68	65.64
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	3.10	0.00	1.80	0.00	0.50	0.00	97.40	33.90	---	---	0.10	0.00	102.90	33.90	136.80
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	4.49	0.00	2.80	0.00	0.44	0.00	39.69	258.47	---	---	4.30	0.00	51.72	258.47	310.19
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	5.82	0.00	3.12	0.00	0.44	0.00	18.85	56.92	---	---	4.40	0.00	32.63	56.92	89.55
1978	6.98	0.00	3.25	0.00	1.05	0.00	49.56	165.80	---	---	1.50	0.00	62.34	165.80	228.14
1979	6.23	0.00	7.96	0.00	0.67	0.00	66.23	235.02	---	---	0.43	0.00	81.52	235.02	316.54
1980	6.21	0.00	4.04	0.00	1.07	0.00	64.15	214.55	---	---	4.30	0.00	79.77	214.55	294.32
1981	8.07	0.00	7.46	0.00	0.24	0.00	105.67	202.92	---	---	0.15	0.00	121.59	202.92	324.51
1982	7.51	0.00	11.77	0.00	0.23	0.00	88.47	162.44	---	---	0.28	0.00	108.26	162.44	270.70
1983	8.87	0.00	8.19	0.00	0.23	0.00	88.99	175.78	---	---	0.24	0.00	106.52	175.78	282.30
1984	8.08	0.00	7.34	0.00	0.26	0.00	88.63	174.40	---	---	0.31	0.00	104.62	174.40	279.02
1985	8.84	0.00	7.29	0.00	0.12	0.00	27.46	105.05	0.97	2.13	0.31	0.00	44.99	107.18	152.17
1986	10.38	0.00	7.47	0.00	0.18	0.00	41.14	112.81	3.09	1.44	0.24	0.00	62.50	114.25	176.75
1987	11.98	0.00	8.33	0.00	0.26	0.00	41.09	105.22	2.24	1.06	0.25	0.00	64.15	106.28	170.43
1988	12.28	0.00	8.96	0.00	0.27	0.00	53.69	134.77	2.88	1.37	0.08	0.00	78.16	136.14	214.30
1989	13.74	0.00	10.29	0.00	0.31	0.00	43.99	106.16	2.55	1.20	0.08	0.00	70.96	107.36	178.32
1990	13.17	0.00	8.94	0.00	0.29	0.00	47.89	164.43	2.48	1.19	0.08	0.00	72.85	165.62	238.47
1991	11.51	0.00	7.32	0.00	0.29	0.00	40.27	97.25	2.65	1.27	0.08	0.00	62.12	98.52	160.64
1992	11.63	0.00	6.90	0.00	0.22	0.00	75.30	178.00	1.62	0.77	0.08	0.00	95.75	178.77	274.52
1993	11.39	0.00	5.17	0.00	0.18	0.00	59.00	132.50	2.80	1.38	0.08	0.00	79.66	133.88	213.54
1994	9.91	0.00	4.25	0.00	0.72	0.00	63.20	142.49	2.76	1.36	0.21	0.00	82.79	143.85	226.64
1995	11.16	0.00	5.98	0.00	0.16	0.00	56.34	135.19	2.01	1.10	0.00	0.00	76.58	136.29	212.87
1996	11.36	0.00	6.54	0.00	0.14	0.00	49.80	128.56	2.42	1.30	0.00	0.00	71.09	129.86	200.95
1997	11.62	0.00	6.90	0.00	0.11	0.00	33.72	85.90	2.07	1.10	0.00	0.00	53.24	87.00	140.24
1998	12.97	0.00	7.38	0.00	0.15	0.00	54.36	120.86	2.16	1.15	0.00	0.00	76.32	122.01	198.33
1999	13.37	0.00	9.15	0.00	0.13	0.00	38.47	95.78	1.69	0.91	0.00	0.00	59.90	96.69	156.59
2000	13.93	0.00	6.23	0.00	0.12	0.00	64.63	157.86	6.81	3.22	0.00	0.00	87.36	161.08	248.44

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources;

1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Jackson County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.50	0.00	1.10	0.00	0.40	0.00	0.05	0.87	---	---	0.16	115.00	3.21	115.87	119.08
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	1.60	0.00	2.33	0.00	1.20	0.00	0.70	0.02	---	---	1.40	144.00	7.23	144.02	151.25
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	1.77	0.00	2.48	0.00	0.80	0.00	5.70	0.71	---	---	0.30	120.42	11.05	121.13	132.18
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.99	0.00	2.59	0.00	0.80	0.00	4.47	1.71	---	---	0.32	109.20	10.17	110.91	121.08
1978	2.04	0.00	2.83	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	2.15	0.00	2.63	0.00	0.84	0.00	9.28	2.63	---	---	0.30	118.00	15.20	120.63	135.83
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	2.44	0.00	4.10	0.00	0.81	0.00	10.27	9.99	0.08	0.01	0.74	100.89	18.44	110.89	129.33
1986	2.58	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	2.47	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	2.60	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	2.55	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	2.33	0.00	3.87	0.00	1.38	0.00	22.13	3.38	0.00	0.00	0.37	107.99	30.08	111.37	141.45
1991	2.19	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	2.26	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	2.41	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	2.32	0.00	---	---	---	---	---	---	---	---	0.39	43.96	N/A	N/A	N/A
1995	2.31	0.00	4.30	0.00	1.67	0.00	19.31	2.55	0.34	0.00	0.29	50.28	28.22	52.83	81.05
1996	2.40	0.00	---	---	---	---	---	---	---	---	0.46	62.15	N/A	N/A	N/A
1997	2.79	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	2.87	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	2.78	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	2.46	0.00	3.22	0.00	1.50	0.00	14.00	1.70	0.39	0.00	0.30	89.53	21.87	91.23	113.10

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Jefferson County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.30	0.00	0.30	0.00	0.00	0.00	0.16	0.08	---	---	0.00	0.00	0.76	0.08	0.84
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.36	0.00	0.73	0.00	0.20	0.00	0.40	0.04	---	---	0.00	0.00	1.69	0.04	1.73
1971	0.38	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1972	0.43	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1973	0.39	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1974	0.43	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1975	0.44	0.00	0.42	0.00	0.02	0.00	1.51	0.14	---	---	0.00	0.00	2.39	0.14	2.53
1976	0.50	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1977	0.57	0.00	0.65	0.00	0.02	0.00	0.75	0.14	---	---	0.00	0.00	1.99	0.14	2.13
1978	0.48	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	0.51	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1980	0.49	0.00	0.89	0.00	0.05	0.00	0.98	0.14	---	---	0.00	0.00	2.41	0.14	2.55
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.60	0.00	1.29	0.00	0.00	0.00	6.31	1.36	0.00	0.00	0.00	0.00	8.20	1.36	9.56
1986	0.67	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.63	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.68	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.70	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.72	0.00	1.04	0.00	0.08	0.00	8.34	1.81	0.11	0.00	0.00	0.00	10.29	1.81	12.10
1991	0.65	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.67	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.71	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.63	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.71	0.00	1.26	0.00	0.25	0.00	8.52	0.52	0.35	0.00	0.00	0.00	11.09	0.52	11.61
1996	0.74	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.67	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.69	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.66	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.72	0.00	0.84	0.00	0.20	0.00	6.43	0.10	0.20	0.07	0.00	0.00	8.39	0.17	8.56

Jefferson County is located within the Northwest Florida and Suwannee River Water Management Districts. The values shown represent the county totals as reported by the Districts. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Lafayette County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.10	0.00	0.00	0.00	0.00	0.00	0.34	0.10	----	----	0.00	0.00	0.54	0.10	0.64
1966	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1970	0.10	0.00	0.00	0.00	0.00	0.00	0.90	0.20	----	----	0.00	0.00	1.24	0.20	1.44
1971	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1975	0.14	0.00	0.00	0.00	0.00	0.00	2.60	0.17	----	----	0.00	0.00	2.95	0.17	3.12
1976	----	----	----	----	----	----	----	----	----	----	0.00	0.00	4.12	0.12	4.24
1977	0.17	0.00	0.00	0.00	0.00	0.00	3.66	0.12	----	----	0.00	0.00	N/A	N/A	N/A
1978	0.12	0.00	0.00	0.00	----	----	----	----	----	----	0.00	0.00	----	----	----
1979	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1980	0.12	0.00	0.00	0.00	0.00	0.00	1.82	0.29	----	----	0.00	0.00	2.28	0.29	2.57
1981	0.14	0.00	0.00	0.00	0.00	0.00	----	----	----	----	0.00	0.00	N/A	N/A	N/A
1982	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1985	0.15	0.00	0.00	0.00	0.08	0.00	6.50	0.26	0.00	0.00	0.00	0.00	7.31	0.26	7.57
1986	0.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1987	0.16	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1988	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1989	0.19	0.00	0.00	0.00	0.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1990	0.18	0.00	0.00	0.00	0.14	0.00	8.08	0.54	0.00	0.00	0.00	0.00	9.10	0.54	9.64
1991	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1992	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1993	0.17	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1994	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1995	0.18	0.00	0.00	0.00	0.23	0.00	5.92	0.34	0.00	0.00	0.00	0.00	7.11	0.34	7.45
1996	0.17	0.00	0.00	0.00	0.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1997	0.18	0.00	0.00	0.00	0.20	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1998	0.20	0.00	0.00	0.00	0.19	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1999	0.24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
2000	0.20	0.00	0.00	0.00	0.20	0.00	5.77	0.16	0.00	0.00	0.00	0.00	6.78	0.16	6.94

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Lake County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	8.00	0.00	1.72	0.00	19.30	0.00	14.08	12.32	----	----	0.00	0.00	43.10	12.32	55.42
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	10.00	0.00	3.25	0.00	19.40	0.00	13.40	7.70	----	----	0.00	0.00	46.05	7.70	53.75
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	9.85	0.00	4.20	0.00	20.65	0.00	38.54	19.27	----	----	0.00	0.00	73.24	19.27	92.51
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	9.94	0.00	5.89	0.00	20.65	0.00	107.20	35.89	----	----	0.00	0.00	143.68	35.89	179.57
1978	11.20	0.00	3.31	0.00	17.47	0.00	87.60	29.50	----	----	0.00	0.00	119.58	29.50	149.08
1979	10.80	0.00	3.11	0.00	13.22	0.00	85.60	28.80	----	----	0.00	0.00	112.73	28.80	141.53
1980	11.39	0.00	5.66	0.00	11.18	0.00	65.31	7.61	----	----	0.00	0.00	93.54	7.61	101.15
1981	12.31	0.00	5.92	0.00	20.30	0.00	99.65	21.73	----	----	0.00	0.00	138.18	21.73	159.91
1982	11.67	0.00	8.89	0.00	15.19	4.13	85.24	17.97	----	----	0.00	0.00	120.99	22.10	143.09
1983	11.94	0.00	8.70	0.00	15.65	3.40	83.82	17.85	----	----	0.00	0.00	120.11	21.25	141.36
1984	14.21	0.00	9.11	0.00	16.43	0.00	86.85	18.70	----	----	0.00	0.00	126.60	18.70	145.30
1985	15.34	0.00	8.52	0.00	12.24	3.41	28.78	4.89	----	----	0.00	0.00	64.88	8.30	73.18
1986	16.25	0.00	8.27	0.00	11.53	3.94	33.61	8.45	----	----	0.00	0.00	69.66	12.39	82.05
1987	17.66	0.00	8.76	0.00	11.68	0.00	51.66	12.74	----	----	0.00	0.00	89.76	12.74	102.50
1988	18.21	0.00	8.52	0.00	13.33	0.00	47.89	11.33	----	----	0.00	0.00	87.95	11.33	99.28
1989	20.77	0.00	9.47	0.00	12.29	0.00	53.47	13.52	----	----	0.00	0.00	96.00	13.52	109.52
1990	20.67	0.00	9.56	0.00	9.51	0.00	43.22	11.64	1.39	1.04	0.00	0.00	84.35	12.68	97.03
1991	19.19	0.00	8.64	0.00	8.66	0.64	59.22	15.46	1.57	1.10	0.00	0.00	97.28	17.20	114.48
1992	18.95	0.00	4.94	0.00	5.62	0.72	40.33	10.32	0.66	0.47	0.00	0.00	70.50	11.51	82.01
1993	21.78	0.00	5.70	0.00	4.40	1.14	40.78	9.78	1.35	1.10	0.00	0.00	74.01	12.02	86.03
1994	20.23	0.00	5.20	0.00	7.89	1.14	42.33	10.37	1.35	1.10	0.00	0.00	77.00	12.61	89.61
1995	26.46	0.00	2.71	0.00	10.23	1.14	34.97	5.80	1.06	0.74	0.00	0.00	75.43	7.68	83.11
1996	29.35	0.00	2.00	0.00	8.51	0.73	38.40	6.61	1.48	1.04	0.00	0.00	79.74	8.38	88.12
1997	30.50	0.00	1.85	0.00	8.81	0.73	36.20	6.10	1.38	0.99	0.00	0.00	78.74	7.82	86.56
1998	35.49	0.00	2.74	0.00	8.58	0.49	54.52	9.77	1.90	1.34	0.00	0.00	103.23	11.60	114.83
1999	36.50	0.00	2.93	0.00	10.38	0.21	28.76	5.11	1.14	0.80	0.00	0.00	79.71	6.12	85.83
2000	39.92	0.00	4.27	0.00	10.44	0.60	30.85	5.30	5.36	3.87	0.00	0.00	90.84	9.77	100.61

Lake County is located within the St. Johns River and the Southwest Florida Water Management Districts. The values shown represent the County totals as reported by the Districts.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1989.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District and the Southwest Florida Water Management District.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004), St. Johns River Water Management District (2004), and the Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Lee County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	4.30	0.00	1.40	0.00	1.00	0.00	0.04	28.20	---	---	0.15	0.00	6.89	28.20	35.09
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	6.70	1.60	1.64	0.00	0.30	4.00	27.30	8.00	---	---	0.04	0.00	35.98	13.60	49.58
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	12.06	3.59	N/A	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1975	9.97	6.85	2.00	0.00	0.40	8.00	48.83	15.56	---	---	0.04	0.00	61.24	30.41	91.65
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	12.27	6.74	2.44	0.00	0.40	8.00	71.34	22.54	---	---	0.04	0.00	86.49	37.28	123.77
1978	13.48	3.28	5.40	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	24.53	5.20	6.14	0.00	4.09	0.00	39.41	7.18	---	---	0.68	0.00	74.85	12.38	87.23
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	29.47	2.28	8.81	0.00	6.87	0.00	18.75	5.07	8.29	0.00	0.12	0.00	72.31	7.35	79.66
1986	28.88	2.27	9.48	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	30.96	2.57	10.37	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	34.52	3.08	11.57	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	37.02	3.08	13.59	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	38.88	3.25	20.44	8.14	7.73	2.75	48.49	13.74	19.56	18.51	0.25	0.00	123.05	38.25	161.30
1991	36.14	3.38	15.17	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	35.97	3.11	14.89	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	36.32	3.13	20.98	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	37.71	3.41	21.09	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	37.64	3.09	19.77	7.55	4.14	5.13	52.07	10.28	11.08	2.95	0.08	0.00	112.56	21.45	134.01
1996	40.73	3.11	21.80	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	41.23	3.03	21.47	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	43.90	3.20	23.12	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	46.26	2.95	23.20	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	49.09	3.28	24.37	8.86	0.00	8.76	7.19	42.51	18.00	11.36	0.22	0.00	120.80	39.77	160.57

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted from under Agricultural from 1965 through 1984.

- Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).
 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.
 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).
 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Leon County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	9.00	0.00	1.60	0.00	0.00	0.00	0.17	0.40	----	----	0.00	0.00	10.77	0.40	11.17
1966	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1970	12.12	0.00	3.04	0.00	0.00	0.00	0.04	0.01	----	----	0.00	0.00	15.20	0.01	15.21
1971	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1975	15.83	0.00	3.21	0.00	0.06	0.00	0.48	0.27	----	----	1.18	0.00	20.76	0.27	21.03
1976	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1977	17.03	0.00	4.30	0.00	1.22	0.00	0.46	0.30	----	----	1.85	0.00	24.86	0.30	25.16
1978	17.08	0.00	----	----	----	----	----	----	----	----	0.00	0.00	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1980	17.20	0.00	3.34	0.00	1.86	0.00	2.42	0.19	----	----	4.20	0.00	29.02	0.19	29.21
1981	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	0.00	0.00	----	----	----
1985	22.06	0.00	4.42	0.00	0.33	0.00	1.25	2.17	2.11	0.00	4.27	0.00	34.44	2.17	36.61
1986	23.44	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1987	23.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1988	25.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1989	24.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1990	25.02	0.00	5.74	0.00	0.26	0.00	1.96	0.48	1.39	0.53	4.11	0.00	38.48	1.01	39.49
1991	24.26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1992	26.88	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1993	28.50	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1994	26.24	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1995	28.74	0.00	4.79	0.00	0.23	0.00	0.87	0.21	1.28	0.54	2.64	0.00	38.55	0.75	39.30
1996	28.67	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1997	27.92	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1998	33.83	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
1999	33.45	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	N/A	N/A	N/A
2000	35.70	0.00	4.29	0.00	0.12	0.00	1.27	0.00	1.24	0.30	2.74	0.00	45.36	0.30	45.66

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Levy County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.40	0.00	0.33	0.00	0.02	0.00	1.32	0.07	---	---	0.01	42.00	2.08	42.07	44.15
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.90	0.00	0.65	0.00	0.00	0.00	0.40	0.00	---	---	0.00	0.00	1.95	0.00	1.95
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.98	0.00	0.85	0.00	0.00	0.00	1.80	0.13	---	---	0.00	0.00	3.63	0.13	3.76
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.05	0.00	0.86	0.00	0.00	0.00	2.30	0.54	---	---	0.00	0.00	4.21	0.54	4.75
1978	0.95	0.00	0.97	0.00	0.00	0.00	1.80	0.70	---	---	0.00	0.00	3.72	0.70	4.42
1979	1.00	0.00	1.09	0.00	0.00	0.00	1.59	1.05	---	---	0.00	0.00	3.68	1.05	4.73
1980	1.10	0.00	1.32	0.00	0.22	0.00	1.88	1.08	---	---	0.00	0.00	4.52	1.08	5.60
1981	1.19	0.00	1.36	0.00	0.18	0.00	1.80	1.32	---	---	0.00	0.00	4.53	1.32	5.85
1982	---	---	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1983	---	---	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1984	---	---	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1985	1.19	0.00	2.17	0.00	0.41	2.28	10.37	2.17	0.00	0.00	0.00	0.00	14.14	4.45	18.59
1986	1.25	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	1.33	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	1.35	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	1.48	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	1.58	0.00	2.15	0.00	0.36	1.70	17.14	1.34	0.29	0.00	0.00	0.00	21.52	3.04	24.56
1991	1.88	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	1.59	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	1.76	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	1.69	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	1.85	0.00	3.66	0.00	0.07	2.00	15.19	0.49	0.16	0.00	0.00	0.00	20.93	2.49	23.42
1996	1.86	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	1.94	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	1.96	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	1.92	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	2.16	0.00	3.95	0.00	0.06	1.96	21.82	0.63	0.45	0.04	0.00	0.00	28.44	2.63	31.07

Levy County is located within the Southwest Florida and the Suwannee River Water Management Districts. The values shown represent the County totals as reported by the Districts.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Liberty County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.10	0.00	0.10	0.00	0.74	0.01	0.01	0.03	----	----	0.00	0.00	0.95	0.04	0.99
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.16	0.00	0.22	0.00	0.80	0.50	0.00	0.00	----	----	0.00	0.00	1.18	0.50	1.68
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.09	0.00	0.24	0.00	0.33	0.00	0.00	0.01	----	----	0.00	0.00	0.66	0.01	0.67
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	0.09	0.00	0.29	0.00	0.33	0.00	0.01	0.03	----	----	0.00	0.00	0.72	0.03	0.75
1978	0.09	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	0.13	0.00	0.27	0.00	0.33	0.00	0.00	0.03	----	----	0.00	0.00	0.73	0.03	0.76
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	0.25	0.00	0.40	0.00	0.00	0.00	0.44	0.00	0.00	0.00	0.00	0.00	1.09	0.00	1.09
1986	0.24	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	0.21	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	0.24	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	0.24	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	0.26	0.00	0.46	0.00	0.22	0.00	2.30	0.00	0.02	0.00	0.00	0.00	3.26	0.00	3.26
1991	0.27	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	0.28	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	0.26	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	0.27	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	0.32	0.00	0.50	0.00	0.38	0.00	0.34	0.07	0.02	0.00	0.00	0.00	1.56	0.07	1.63
1996	0.36	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	0.37	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	0.41	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	0.41	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	0.39	0.00	0.45	0.00	0.82	0.00	0.16	0.00	0.00	0.00	0.00	0.00	1.82	0.00	1.82

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Madison County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.70	0.00	0.40	0.00	0.00	0.00	0.16	0.52	----	----	0.00	0.00	1.26	0.52	1.78
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.60	0.00	0.85	0.00	0.00	0.00	1.60	0.10	----	----	0.00	0.00	3.05	0.10	3.15
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	1.09	0.00	0.75	0.00	0.03	0.00	1.93	0.21	----	----	0.00	0.00	3.80	0.21	4.01
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	1.09	0.00	0.83	0.00	0.03	0.00	3.38	0.40	----	----	0.00	0.00	5.33	0.40	5.73
1978	1.07	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	0.96	0.00	0.93	0.00	0.04	0.00	1.45	1.06	----	----	0.00	0.00	3.38	1.06	4.44
1981	0.99	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	1.18	0.00	1.90	0.00	0.46	0.00	2.46	0.00	0.00	0.00	0.00	0.00	6.00	0.00	6.00
1986	1.32	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	1.36	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	1.42	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	1.39	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	1.43	0.00	2.56	0.00	0.23	0.00	2.97	0.27	0.00	0.00	0.00	0.00	7.19	0.27	7.46
1991	1.37	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	1.87	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	1.55	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1.54	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1.58	0.00	1.54	0.00	0.11	0.00	5.44	0.55	0.11	0.00	0.00	0.00	8.78	0.55	9.33
1996	1.56	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	1.53	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	1.61	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	1.54	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	1.65	0.00	1.23	0.00	0.15	0.00	5.83	0.11	0.20	0.06	0.00	0.00	9.06	0.17	9.23

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Manatee County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	3.80	2.60	0.00	0.00	1.30	0.00	35.70	2.50	----	----	0.00	0.00	41.63	5.10	46.73
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.30	9.60	0.00	3.88	3.00	0.00	49.50	0.00	----	----	0.00	0.00	56.68	9.60	66.28
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.00	19.89	0.98	4.40	1.99	0.00	24.43	1.38	----	----	0.02	25.00	30.84	46.27	77.11
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	0.00	24.46	2.10	4.68	3.35	0.00	42.78	2.36	----	----	0.01	14.00	50.82	40.82	91.64
1978	0.00	24.30	6.30	4.37	5.20	0.00	47.80	5.34	----	----	0.05	2.60	57.42	32.24	89.66
1979	0.00	27.20	6.60	2.16	0.20	0.00	61.91	6.88	----	----	0.01	9.80	64.28	43.88	108.16
1980	0.00	28.07	7.17	2.41	0.20	0.00	67.59	0.96	----	----	0.25	3.60	70.45	32.63	103.08
1981	0.00	30.34	8.34	2.30	0.50	0.00	84.15	1.12	----	----	0.40	10.00	87.35	41.46	128.81
1982	0.00	32.91	9.46	2.38	4.45	0.00	73.16	0.74	----	----	0.59	6.99	80.58	40.64	121.22
1983	0.00	31.05	9.61	3.78	4.34	0.00	66.81	0.67	----	----	0.66	4.00	75.59	35.72	111.31
1984	0.00	33.04	10.00	2.94	3.62	0.00	101.50	1.03	----	----	0.69	1.53	108.75	35.60	144.35
1985	0.00	30.89	9.40	0.43	3.30	0.00	87.79	0.34	1.22	0.57	0.44	2.51	93.18	34.31	127.49
1986	0.00	32.77	10.20	0.28	4.75	0.00	84.69	0.86	----	----	0.57	10.60	90.29	44.23	134.52
1987	0.00	35.28	10.29	0.45	2.16	0.00	73.89	0.89	----	----	0.55	12.41	77.05	48.58	125.63
1988	0.00	38.44	11.67	1.04	3.27	0.00	89.48	0.90	----	----	0.58	6.58	94.37	45.92	140.29
1989	0.00	37.58	11.35	4.92	2.41	0.00	95.12	1.77	----	----	0.15	10.95	102.60	50.30	152.90
1990	0.09	40.30	12.55	0.08	2.58	0.00	92.25	1.50	1.53	0.24	0.00	3.27	96.53	45.31	141.84
1991	0.00	54.79	12.41	0.08	5.22	2.77	59.85	14.00	1.53	1.89	0.00	1.53	66.68	74.98	141.66
1992	0.93	48.28	11.69	0.20	3.48	2.99	86.10	7.22	1.63	1.33	0.00	1.52	92.34	61.34	153.68
1993	3.74	39.88	10.44	3.29	0.25	0.78	67.87	1.00	1.32	0.98	0.27	0.11	76.74	42.75	119.49
1994	12.11	32.87	11.37	3.09	0.16	0.14	72.15	0.79	1.18	1.02	0.43	0.00	89.12	34.82	123.94
1995	12.22	29.88	9.00	3.00	0.57	0.26	73.67	0.94	1.24	0.94	0.00	0.01	90.70	32.03	122.73
1996	13.90	29.85	9.35	2.48	0.39	0.27	79.62	0.83	2.10	1.20	0.30	0.19	98.79	32.34	131.13
1997	12.84	29.82	9.22	2.30	0.33	0.57	75.04	0.98	1.84	0.65	0.20	0.20	92.55	32.22	124.77
1998	18.97	24.40	9.23	0.08	2.15	0.48	78.63	1.58	2.42	0.77	0.31	0.26	102.56	27.49	130.05
1999	13.10	29.40	9.72	0.19	5.00	0.81	91.00	1.15	2.53	0.94	0.00	0.00	111.82	32.30	144.12
2000	13.87	36.05	9.71	0.17	0.47	0.53	104.37	1.35	3.42	1.62	0.00	25.10	122.30	64.65	186.95

Public supply transfers represent water withdrawn in Manatee County and exported for use in Sarasota County.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989. Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Marion County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	3.80	0.00	0.00	0.00	4.50	0.00	7.60	1.60	----	----	0.00	0.00	18.10	1.60	19.70
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	3.90	0.00	0.00	0.00	2.10	0.00	5.70	1.10	----	----	0.00	0.00	16.57	1.10	17.67
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	6.23	0.00	0.00	0.00	0.30	0.00	18.07	0.85	----	----	0.00	0.00	31.18	0.85	32.03
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	6.08	0.00	0.00	0.00	0.31	0.00	19.38	0.92	----	----	0.00	0.00	33.24	0.92	34.16
1978	5.90	0.00	0.00	0.00	0.90	0.00	25.10	1.30	----	----	0.00	0.00	37.77	1.30	39.07
1979	5.90	0.00	0.00	0.00	0.30	0.00	16.34	1.60	----	----	0.00	0.00	29.36	1.60	30.96
1980	6.78	0.00	0.00	0.00	1.57	0.00	20.19	1.06	----	----	0.00	0.00	35.87	1.06	36.93
1981	7.40	0.00	0.00	0.00	0.91	0.00	26.47	0.98	----	----	0.00	0.00	42.78	0.98	43.76
1982	9.37	0.00	0.00	0.00	0.20	0.00	21.81	2.95	----	----	0.00	0.00	40.04	2.95	42.99
1983	9.83	0.00	0.00	0.00	0.18	0.00	22.43	2.88	----	----	0.00	0.00	42.06	2.88	44.94
1984	10.79	0.00	0.00	0.00	0.22	0.00	29.57	3.31	----	----	0.00	0.00	53.17	3.31	56.48
1985	11.89	0.00	0.00	0.00	0.93	1.93	19.01	0.96	2.05	1.00	0.00	0.00	49.14	3.89	53.03
1986	12.92	0.00	0.00	0.00	0.73	1.93	21.53	2.78	----	----	0.00	0.00	48.93	4.71	53.64
1987	13.21	0.00	0.00	0.00	0.88	0.00	14.42	1.88	----	----	0.00	0.00	43.23	1.88	45.11
1988	14.62	0.00	0.00	0.00	1.06	0.00	9.92	1.85	----	----	0.00	0.00	40.60	1.85	42.45
1989	15.48	0.00	0.00	0.00	1.02	0.00	14.05	1.76	----	----	0.00	0.00	47.58	1.76	49.34
1990	16.47	0.00	0.00	0.00	1.07	0.00	12.86	1.14	1.87	0.73	0.00	0.00	49.20	1.87	51.07
1991	16.46	0.00	0.00	0.00	1.61	0.00	8.64	1.37	1.82	1.61	0.00	0.00	44.70	2.98	47.68
1992	20.30	0.00	0.00	0.00	1.75	0.00	10.50	0.95	1.32	1.21	0.00	0.00	52.83	2.16	54.99
1993	17.77	0.11	0.00	0.00	1.70	0.00	7.49	0.63	3.37	0.70	0.00	0.00	47.16	1.44	48.60
1994	16.31	0.10	0.00	0.00	3.72	0.00	7.62	0.64	2.90	0.61	0.00	0.00	49.21	1.35	50.56
1995	20.13	0.14	0.00	0.00	1.91	0.00	6.20	0.44	2.46	0.46	0.00	0.00	51.03	1.04	52.07
1996	21.48	0.08	0.00	0.00	1.85	0.00	7.41	0.55	3.17	0.61	0.00	0.00	55.77	1.24	57.01
1997	23.14	0.00	0.00	0.00	1.82	0.00	8.34	0.59	3.51	0.68	0.00	0.00	58.20	1.27	59.47
1998	24.87	0.03	0.00	0.00	1.18	0.00	9.85	0.84	3.67	0.70	0.00	0.00	62.87	1.57	64.44
1999	25.85	0.00	0.00	0.00	1.51	0.00	6.80	0.50	3.57	0.50	0.00	0.00	62.97	1.00	63.97
2000	27.99	0.00	0.00	0.00	2.08	0.00	16.31	0.90	4.88	1.20	0.00	0.00	67.68	2.10	69.78

Marion County is located within the St. Johns River and the Southwest Florida Water Management Districts. The values shown represent the County totals as reported by the Districts.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965 - 1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991 - 1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District and the Southwest Florida Water Management District.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004), St. Johns River Water Management District (2004), and the Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Martin County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.90	0.00	1.10	0.00	0.50	0.00	11.10	44.53	---	---	0.00	0.00	14.60	44.53	59.13
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	1.60	0.00	1.92	0.00	0.50	0.00	15.40	74.90	---	---	0.00	0.00	19.42	74.90	94.32
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	5.72	0.00	2.40	0.00	0.08	0.00	7.29	76.76	---	---	0.00	0.00	15.49	76.76	92.25
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	5.51	0.00	2.66	0.00	0.10	0.00	8.77	81.36	---	---	0.00	0.00	17.04	81.36	98.40
1978	5.40	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	6.15	0.00	5.96	0.00	0.12	0.00	32.34	107.01	---	---	0.00	40.30	44.57	147.31	191.88
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	9.33	0.00	7.50	0.00	0.17	0.00	27.47	117.10	1.33	1.33	0.49	24.50	46.29	142.93	189.22
1986	9.26	0.00	0.21	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	10.11	0.00	0.22	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	11.20	0.00	0.27	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	12.99	0.00	0.25	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	13.66	0.00	0.31	0.00	0.77	28.77	36.77	101.36	5.35	4.21	0.53	19.56	65.71	153.90	219.61
1991	12.68	0.00	0.31	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	13.18	0.00	0.27	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	12.90	0.00	0.33	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	12.24	0.00	0.43	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	14.00	0.00	1.22	0.00	1.66	0.00	26.36	100.98	0.00	0.00	0.16	19.25	49.80	120.23	170.03
1996	15.18	0.00	2.00	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	14.22	0.00	1.56	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	15.86	0.00	3.21	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	15.75	0.00	3.73	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	18.45	0.00	2.82	0.00	1.20	1.95	15.02	125.00	4.19	3.69	0.54	24.63	43.60	155.27	198.87

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Miami-Dade County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	202.30	0.00	5.90	0.00	5.00	2.70	67.90	0.00	----	----	0.30	0.00	285.10	2.70	287.80
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	212.10	0.00	5.20	0.00	7.70	2.70	44.80	0.00	----	----	0.04	0.00	273.77	2.70	276.47
1971	----	----	5.66	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	5.81	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	6.02	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	5.66	----	----	----	----	----	----	----	----	----	----	----	----
1975	270.51	0.00	5.96	0.00	3.38	0.00	87.66	2.90	----	----	0.04	0.00	371.09	2.90	373.99
1976	----	----	5.79	----	----	----	----	----	----	----	----	----	----	----	----
1977	280.15	0.00	5.96	0.00	6.73	0.00	101.06	2.96	----	----	0.00	0.00	391.92	2.96	394.88
1978	268.11	0.00	6.44	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	314.29	0.00	8.61	0.00	19.73	0.00	86.98	2.76	----	----	0.00	0.00	439.38	2.76	442.14
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	339.77	0.00	11.34	0.00	15.78	0.00	103.68	0.00	13.50	4.50	0.00	0.00	486.05	4.50	490.55
1986	345.41	0.00	12.72	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	365.24	0.00	13.14	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	365.62	0.00	13.17	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	367.00	0.00	13.37	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	337.69	0.00	12.07	0.00	40.34	0.00	115.01	7.14	20.55	7.23	2.26	0.00	526.60	14.37	540.97
1991	352.06	0.00	11.77	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	378.57	0.00	12.96	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	388.16	0.00	14.15	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	388.72	0.00	14.05	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	386.60	0.00	14.07	0.00	38.82	4.56	95.95	11.72	14.24	2.55	2.10	0.00	550.42	18.83	569.25
1996	389.16	0.00	14.39	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	386.40	0.00	14.68	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	390.33	0.00	15.42	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	389.41	0.00	16.31	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	394.29	0.00	17.02	4.85	41.65	0.00	86.55	23.80	8.51	4.88	2.08	0.00	537.93	28.68	566.61

Public supply transfers represent water withdrawn in Miami-Dade County and exported for use in Monroe County.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural self-supplied includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) purposes. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Monroe County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	---	---	0.20	0.00	0.20	0.00	0.20
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	1.58	0.00	0.00	0.00	0.00	0.00	0.00	0.00	---	---	0.00	0.00	1.58	0.00	1.58
1971	0.98	0.00	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	1.16	0.00	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	1.60	0.00	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	1.57	0.00	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	1.71	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00	1.81	0.00	1.81
1976	1.25	0.00	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.27	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.10	0.00	1.43	0.00	1.43
1978	1.96	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	2.46	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.49	0.00	0.00	0.00	2.95	0.00	2.95
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.60	0.00	0.00	0.00	1.60	0.00	1.60
1986	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	0.00	0.00	0.00	0.00	0.07	0.00	0.00	0.00	1.22	0.00	0.00	0.00	1.29	0.00	1.29
1991	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	0.00	0.00	0.51	0.00	0.07	0.00	0.04	0.00	1.16	0.00	0.00	0.00	1.78	0.00	1.78
1996	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	0.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	0.00	0.00	0.08	0.00	0.10	0.00	0.03	0.00	1.23	0.62	0.00	0.00	1.44	0.62	2.06

Public supply transfers represent water withdrawn in Miami-Dade County and imported for use in Monroe County.

Agricultural self-supplied includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) purposes. This category was accounted for under Agricultural from 1965 through 1974.

Ground water withdrawn for public supply and recreational irrigation in Monroe County may be saline water and is usually treated through a desalinations process for use.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Nassau County, 1965-2000 (Continued)
 [Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.20	0.00	0.00	0.00	37.80	0.00	0.81	0.09	----	----	0.00	0.00	40.31	0.09	40.40
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	2.00	0.00	1.40	0.00	50.00	0.00	0.00	0.00	----	----	0.00	0.00	53.40	0.00	53.40
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	2.40	0.00	1.83	0.00	57.93	0.00	0.93	0.00	----	----	0.00	0.00	63.09	0.00	63.09
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	3.07	0.00	2.36	0.00	58.11	0.00	1.61	1.44	----	----	0.00	0.00	65.15	1.44	66.59
1978	2.54	0.00	2.58	0.00	58.89	0.00	1.58	1.43	----	----	0.00	0.00	65.59	1.43	67.02
1979	2.32	0.00	4.09	0.00	55.41	0.00	1.42	0.86	----	----	0.00	0.00	63.24	0.86	64.10
1980	2.58	0.00	4.55	0.00	44.97	0.00	1.59	0.86	----	----	0.00	0.00	53.69	0.86	54.55
1981	2.90	0.00	5.19	0.00	41.41	0.00	1.59	0.86	----	----	0.00	0.00	51.09	0.86	51.95
1982	3.29	0.00	5.72	0.00	41.90	0.00	1.60	0.86	----	----	0.00	0.00	52.51	0.86	53.37
1983	2.76	0.00	3.81	0.00	32.01	0.00	1.47	0.89	----	----	0.00	0.00	40.05	0.89	40.94
1984	2.89	0.00	3.75	0.00	36.00	0.00	1.30	0.87	----	----	0.00	0.00	43.94	0.87	44.81
1985	3.04	0.00	3.97	0.00	37.40	0.00	1.01	0.83	0.72	0.10	0.00	0.00	46.14	0.93	47.07
1986	3.12	0.00	3.95	0.00	38.42	0.00	0.98	0.80	1.19	0.17	0.00	0.00	47.66	0.97	48.63
1987	3.54	0.00	3.98	0.00	33.93	0.00	1.07	0.80	1.12	0.16	0.00	0.00	43.64	0.96	44.60
1988	3.58	0.00	4.08	0.00	33.25	0.00	1.06	0.80	0.94	0.13	0.00	0.00	42.91	0.93	43.84
1989	3.59	0.00	4.11	0.00	31.64	0.00	0.60	0.33	1.06	0.16	0.00	0.00	41.00	0.49	41.49
1990	3.85	0.00	3.83	0.00	32.69	0.00	0.65	0.27	1.75	0.33	0.00	0.00	42.77	0.60	43.37
1991	4.04	0.00	3.96	0.00	32.71	0.00	0.81	0.00	1.53	0.23	0.00	0.00	43.05	0.23	43.28
1992	4.02	0.00	3.85	0.00	36.65	0.00	0.81	0.00	0.40	0.06	0.00	0.00	45.73	0.06	45.79
1993	4.43	0.00	4.39	0.00	38.10	0.00	0.25	0.00	1.39	0.20	0.00	0.00	48.56	0.20	48.76
1994	4.53	0.00	4.51	0.00	34.55	0.00	0.25	0.00	1.36	0.21	0.00	0.00	45.20	0.21	45.41
1995	4.96	0.00	4.23	0.00	34.49	0.00	0.15	0.00	0.71	0.11	0.00	0.00	44.54	0.11	44.65
1996	5.01	0.00	4.58	0.00	35.73	0.00	0.18	0.00	1.29	0.19	0.00	0.00	46.79	0.19	46.98
1997	5.06	0.00	4.96	0.00	35.23	0.00	0.18	0.00	1.39	0.21	0.00	0.00	46.82	0.21	47.03
1998	5.98	0.00	5.41	0.00	35.03	0.00	0.19	0.00	1.38	0.21	0.00	0.00	47.99	0.21	48.20
1999	6.86	0.00	6.58	0.00	30.56	0.00	0.14	0.00	1.10	0.16	0.00	0.00	45.24	0.16	45.40
2000	6.81	0.00	3.48	0.00	32.46	0.00	0.68	0.02	3.10	0.46	0.00	0.00	46.53	0.48	47.01

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Okaloosa County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	7.50	0.00	2.00	0.00	1.40	0.00	0.16	0.16	----	----	0.00	0.00	11.06	0.16	11.22
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	7.90	0.00	3.00	0.00	4.70	0.00	0.00	0.00	----	----	0.00	0.00	15.60	0.00	15.60
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	9.31	0.00	2.21	0.00	6.05	0.00	0.45	0.45	----	----	0.00	0.00	18.02	0.45	18.47
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	12.79	0.00	3.15	0.00	5.45	0.00	0.34	0.57	----	----	0.00	0.00	21.73	0.57	22.30
1978	13.13	0.00	4.95	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	12.97	0.00	3.76	0.00	4.62	0.00	0.17	0.16	----	----	0.00	0.00	21.52	0.16	21.68
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	17.36	0.00	2.19	0.00	5.82	0.00	0.92	0.00	1.38	0.00	0.00	0.00	27.67	0.00	27.67
1986	17.65	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	18.53	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	19.27	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	19.74	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	20.92	0.00	2.83	0.00	5.98	0.00	0.24	0.00	1.62	0.00	0.00	0.00	31.59	0.00	31.59
1991	19.33	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	19.85	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	20.86	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	20.22	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	21.20	0.00	1.85	0.00	4.84	0.00	0.96	0.23	0.86	0.00	0.00	0.00	29.71	0.23	29.94
1996	21.98	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	21.74	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	21.80	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	22.23	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	22.97	0.00	1.27	0.00	4.14	0.00	0.75	0.00	3.02	0.46	0.00	0.00	32.15	0.46	32.61

The estimation procedure for domestic self-supplied withdrawals changed for 2000 as a different per capita was used to calculate withdrawals compared to previous years.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Okeechobee County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.00	0.90	0.00	0.00	0.00	0.00	32.60	5.60	----	----	0.00	0.00	32.90	6.50	39.40
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.00	0.61	0.00	0.24	0.00	0.00	28.90	5.90	----	----	0.00	0.00	29.14	6.51	35.65
1971	0.00	0.72	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1972	0.00	0.66	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1973	0.00	0.98	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1974	0.00	1.06	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1975	0.00	1.04	0.00	0.90	0.00	0.00	67.95	17.02	----	----	0.00	0.00	68.85	18.06	86.91
1976	0.00	1.11	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1977	0.00	1.33	0.00	1.44	0.00	0.00	77.87	19.89	----	----	0.00	0.00	79.31	21.22	100.53
1978	0.00	1.45	0.00	1.62	0.00	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	0.00	1.49	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1980	0.00	1.50	0.00	1.64	0.00	0.18	74.81	13.32	----	----	0.00	0.00	76.63	14.82	91.45
1981	0.00	1.55	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1982	0.00	1.57	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1983	0.00	1.65	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1984	0.00	1.74	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1985	0.00	1.93	0.00	1.58	0.00	0.12	22.63	3.23	0.11	0.11	0.00	0.00	24.44	5.27	29.71
1986	0.00	2.19	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	0.00	2.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	0.00	2.02	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	0.00	2.14	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	0.00	2.09	0.00	1.56	0.00	0.15	37.19	2.77	0.00	0.50	0.00	0.00	38.90	5.36	44.26
1991	0.00	2.15	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	0.00	2.08	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	0.00	2.18	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	0.07	1.82	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	0.75	1.20	0.00	1.11	0.00	0.13	32.64	4.27	1.12	0.00	0.00	0.00	35.75	5.47	41.22
1996	0.72	1.25	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	0.73	1.26	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	0.57	1.25	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	0.58	1.25	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	0.54	1.69	0.00	1.52	0.00	0.36	57.29	9.75	0.09	0.59	0.00	0.00	59.80	12.03	71.83

Okeechobee County is located within the St Johns River and South Florida Water Management Districts. The values shown represent the County totals as reported by the Districts.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Orange County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	62.93	0.00	13.93	4.00	0.00	0.00	8.90	8.70	---	---	0.00	95.00	81.83	103.70	185.53
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	65.80	0.00	15.30	7.60	0.00	0.00	11.20	8.90	---	---	0.00	128.00	91.60	136.90	228.50
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	77.80	0.00	14.40	8.55	0.00	0.60	13.18	19.84	---	---	0.08	77.40	113.79	97.84	211.63
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	83.30	0.00	14.72	8.62	0.00	0.60	50.43	11.79	---	---	0.08	65.60	156.61	77.99	234.60
1978	84.13	0.00	14.97	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	93.80	0.00	17.51	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1980	87.14	0.00	17.73	10.70	---	1.01	60.26	7.25	---	---	0.00	22.60	172.33	30.86	203.19
1981	88.45	0.00	17.03	---	---	---	---	---	---	---	---	---	---	---	---
1982	94.40	0.00	16.44	---	---	---	---	---	---	---	---	---	---	---	---
1983	98.23	0.00	16.70	---	---	---	---	---	---	---	---	---	---	---	---
1984	112.72	0.00	20.34	---	---	---	---	---	---	---	---	---	---	---	---
1985	122.60	0.00	22.10	6.07	0.00	1.02	43.14	51.87	4.78	1.03	0.00	0.00	191.82	53.92	245.74
1986	132.11	0.00	22.47	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	139.74	0.00	23.49	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	139.81	0.00	25.02	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	153.65	0.00	24.85	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	161.29	0.00	23.52	5.01	0.00	18.88	22.69	60.97	2.67	0.50	0.33	0.00	210.87	61.47	272.34
1991	148.10	0.00	22.46	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	153.11	0.00	24.92	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	154.59	0.00	25.06	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	155.01	0.00	23.54	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	165.02	0.00	23.97	12.95	0.00	19.67	22.09	28.54	8.42	2.46	0.41	0.00	228.56	31.00	259.56
1996	172.06	0.00	25.34	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	168.86	0.00	24.61	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	190.67	0.00	25.73	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	198.60	0.00	24.50	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	211.76	0.00	25.61	8.82	0.00	24.21	19.67	7.32	16.14	9.40	0.76	0.00	281.36	16.72	298.08

Orange County is located within the St Johns River and South Florida Water Management Districts. The values shown represent the County totals as reported by the Districts.

Public supply transfers represent water withdrawn in Orange County and exported for use in Brevard County.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural self-supplied includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) purposes. This category was accounted for under Agricultural from 1965 through 1984.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Osceola County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-Industrial-mining		Agricultural		Recreational Irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	3.30	0.00	0.00	0.00	0.00	0.00	9.40	6.80	---	---	0.20	0.00	13.35	6.80	20.15
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	2.70	0.00	0.00	0.00	0.10	0.00	8.00	5.40	---	---	0.10	0.00	12.91	5.40	18.31
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	3.65	0.00	0.00	0.00	0.70	0.00	8.89	4.17	---	---	0.50	0.00	15.54	4.17	19.71
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	3.96	0.00	0.00	0.00	0.70	0.00	3.30	2.21	---	---	1.00	0.00	12.93	2.21	15.14
1978	4.19	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	4.86	0.00	0.00	0.00	0.70	0.00	18.01	4.06	---	---	1.00	0.00	28.28	4.06	32.34
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	7.03	0.00	0.00	0.00	3.20	0.00	40.04	5.58	0.00	0.00	0.00	0.00	53.69	5.58	59.27
1986	8.30	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	9.48	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	9.25	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	10.98	0.00	0.00	0.00	2.22	0.00	---	---	---	---	---	---	N/A	N/A	N/A
1990	12.20	0.00	0.00	0.00	2.21	0.00	44.42	12.04	0.30	1.36	0.00	0.00	63.44	13.40	76.84
1991	14.50	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	16.80	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	18.31	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	17.63	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	19.15	0.00	0.00	0.00	0.73	0.00	40.53	12.80	1.05	0.41	0.04	0.00	68.27	13.21	81.48
1996	20.38	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	21.58	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	23.02	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	25.88	0.00	0.00	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	30.00	0.00	0.00	0.00	0.84	0.00	80.47	25.49	1.57	0.54	0.03	0.00	117.52	26.03	143.55

Osceola County is located within the St Johns River and South Florida Water Management Districts. The values shown represent the County totals as reported by the Districts.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Palm Beach County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	22.80	15.90	0.00	0.00	2.70	3.00	29.20	333.40	----	----	0.60	0.00	62.20	352.30	414.50
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	34.90	20.40	0.00	0.00	1.80	26.60	123.20	286.60	----	----	0.20	0.00	170.78	333.60	504.38
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	62.73	26.11	0.00	0.00	1.79	44.75	35.96	469.41	----	----	0.18	0.00	113.54	540.72	654.26
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	66.07	30.24	13.36	0.45	2.18	45.43	39.13	472.45	----	----	0.18	0.00	120.92	548.57	669.49
1978	76.22	26.73	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	88.34	27.95	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1980	94.01	29.48	0.00	0.00	1.75	0.92	55.82	559.48	----	----	0.19	0.00	162.51	589.88	752.39
1981	93.28	28.15	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1982	91.90	26.64	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	117.38	29.16	0.00	0.00	2.22	0.54	28.76	493.22	0.00	31.36	0.00	0.00	153.92	554.28	708.20
1986	119.66	29.37	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	136.34	32.29	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	140.32	31.21	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	156.12	32.68	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	135.09	29.90	0.31	0.00	1.99	30.18	13.22	705.46	28.94	31.06	0.00	0.00	200.58	796.60	997.18
1991	145.12	30.75	1.75	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	150.46	31.81	3.19	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	154.65	30.20	2.78	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	150.36	29.21	5.35	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	153.86	33.02	6.75	17.19	1.95	21.00	15.88	655.56	30.15	31.23	0.00	0.00	219.03	740.81	959.84
1996	162.85	31.59	7.66	18.27	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	162.94	32.30	8.79	18.41	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	177.83	35.96	11.78	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	180.87	34.98	16.14	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	194.57	35.27	18.27	10.17	5.76	13.21	42.07	904.09	21.18	52.91	0.00	0.00	273.75	1,005.48	1,279.23

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.
 Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.
 Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).
 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.
 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).
 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Pasco County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	2.10	0.00	0.83	0.00	16.40	0.00	12.55	3.06	----	----	0.00	0.00	31.88	3.06	34.94
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	2.00	0.00	6.20	0.00	30.00	0.00	9.60	0.00	----	----	0.00	0.00	47.80	0.00	47.80
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	18.66	0.00	10.39	0.00	25.01	0.00	39.96	10.43	----	----	0.00	0.00	94.02	10.43	104.45
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	21.67	0.00	11.55	0.00	13.19	0.00	31.09	7.31	----	----	0.00	0.00	77.50	7.31	84.81
1978	40.60	0.00	11.28	0.00	14.50	0.00	27.78	0.00	----	----	0.00	0.00	94.16	0.00	94.16
1979	57.44	0.00	3.89	0.00	13.00	0.00	22.49	0.00	----	----	0.00	0.00	96.82	0.00	96.82
1980	54.65	0.00	5.74	0.00	15.80	0.00	21.26	0.78	----	----	0.00	0.00	97.45	0.78	98.23
1981	61.04	0.00	48.54	0.00	7.72	0.38	24.61	1.25	----	----	0.00	0.00	99.64	1.63	101.27
1982	64.73	0.00	46.30	0.00	23.72	0.00	17.22	0.70	----	----	0.00	0.00	111.39	0.70	112.09
1983	58.82	0.00	41.54	0.00	27.87	0.00	16.15	0.66	----	----	0.00	0.00	108.72	0.66	109.38
1984	63.40	0.00	44.97	0.00	30.28	0.00	23.94	0.92	----	----	0.00	0.00	124.42	0.92	125.34
1985	63.87	0.00	44.10	0.00	19.91	0.00	20.63	0.87	1.79	0.00	0.00	0.00	114.45	0.87	115.32
1986	68.93	0.00	47.86	0.00	20.03	0.00	20.75	0.82	----	----	0.00	0.00	115.22	0.82	116.04
1987	83.87	0.00	62.11	0.00	27.40	0.00	11.61	0.85	----	----	0.00	0.00	127.68	0.85	128.53
1988	83.92	0.00	59.25	0.00	32.51	0.00	21.31	1.80	----	----	0.00	0.00	144.29	1.80	146.09
1989	87.36	0.00	62.86	0.00	38.12	0.00	14.60	2.10	----	----	0.00	0.00	150.19	2.10	152.29
1990	90.65	0.00	65.26	0.00	18.68	0.00	15.10	1.66	1.30	0.65	0.00	0.00	135.92	2.38	138.30
1991	93.84	0.00	58.91	0.13	13.01	39.26	25.78	4.60	2.13	2.14	0.00	0.00	134.89	46.00	180.89
1992	102.26	0.00	68.61	2.89	12.26	38.54	24.06	4.03	2.46	1.49	0.00	0.00	143.93	44.06	187.99
1993	97.06	0.00	58.93	10.76	11.67	8.46	18.16	0.69	3.47	0.47	0.17	0.00	141.29	9.62	150.91
1994	94.44	0.00	62.07	7.99	14.94	10.91	15.88	0.37	4.10	0.56	0.61	0.00	137.96	11.84	149.80
1995	93.93	0.00	67.79	8.81	9.38	11.56	14.49	0.20	2.56	0.43	0.23	0.00	129.40	12.19	141.59
1996	98.53	0.00	66.34	9.72	8.59	11.53	14.06	0.38	2.98	0.67	0.20	0.00	134.08	12.58	146.66
1997	97.29	0.00	61.51	11.30	8.47	8.71	13.87	0.37	3.73	0.84	0.21	0.00	134.87	9.92	144.79
1998	93.20	0.00	57.82	4.39	8.29	4.02	15.23	0.48	3.13	0.85	0.16	0.00	124.40	5.35	129.75
1999	97.94	0.00	60.70	3.47	7.36	1.65	15.57	0.54	4.71	1.11	0.14	0.00	129.19	3.30	132.49
2000	102.67	0.00	67.44	4.50	4.72	0.81	22.37	0.36	5.29	1.07	0.14	0.00	139.69	2.24	141.93

Public supply transfers represent water withdrawn in Pasco County and exported for use in Pinellas County.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989. Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Pinellas County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	32.60	0.00	13.20	0.00	1.00	0.00	12.54	0.24	----	----	0.10	0.00	46.34	0.24	46.58
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	32.00	0.00	28.00	13.12	2.00	0.00	4.00	0.04	----	----	0.00	0.00	51.12	0.04	51.16
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	36.93	0.00	39.97	6.46	1.30	0.00	34.27	0.02	----	----	0.00	0.00	78.96	0.02	78.98
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	42.62	0.00	46.06	2.91	1.28	0.00	28.62	0.00	----	----	0.00	0.00	75.43	0.00	75.43
1978	39.20	0.00	58.60	0.73	1.00	0.00	19.73	0.00	----	----	0.00	0.00	60.66	0.00	60.66
1979	39.85	0.00	61.65	3.00	1.00	0.00	20.38	0.00	----	----	0.00	0.00	64.23	0.00	64.23
1980	36.71	0.00	66.19	2.72	0.90	0.00	9.42	0.46	----	----	0.00	0.00	49.75	0.46	50.21
1981	32.55	0.00	70.85	2.00	0.70	0.00	11.27	0.56	----	----	0.00	0.00	46.52	0.56	47.08
1982	32.21	0.00	70.60	2.27	0.12	0.00	10.92	0.58	----	----	0.00	0.00	45.52	0.58	46.10
1983	56.34	0.00	64.61	2.30	0.11	0.04	11.20	0.59	----	----	0.00	0.00	69.95	0.63	70.58
1984	41.93	0.00	75.79	3.52	0.21	0.03	11.02	0.58	----	----	0.00	0.00	56.68	0.61	57.29
1985	40.13	0.00	76.28	3.72	0.05	0.00	0.95	0.35	9.30	0.20	0.00	0.00	54.15	0.55	54.70
1986	39.51	0.00	77.91	2.37	0.04	0.00	16.22	0.86	----	----	0.00	0.00	58.14	0.86	59.00
1987	40.02	0.00	79.55	1.70	0.07	0.00	6.72	0.63	----	----	0.00	0.00	48.51	0.63	49.14
1988	46.57	0.00	73.68	7.54	0.03	0.00	4.24	0.91	----	----	0.00	0.00	58.38	0.91	59.29
1989	46.88	0.00	69.93	1.77	0.07	0.00	1.54	3.81	----	----	0.00	0.00	50.26	3.81	54.07
1990	40.97	0.00	77.91	1.98	0.08	0.00	1.29	0.00	3.78	1.35	0.00	0.00	48.10	1.35	49.45
1991	39.79	0.00	70.36	0.76	0.14	0.14	0.36	0.85	1.80	11.42	0.00	0.00	42.85	12.41	55.26
1992	32.49	0.00	81.12	1.77	0.17	0.27	0.33	0.77	1.64	4.90	0.00	0.00	36.40	5.94	42.34
1993	37.67	0.00	73.83	8.89	0.08	0.16	0.55	0.03	2.23	4.85	0.00	0.00	49.42	5.04	54.46
1994	38.71	0.00	84.33	9.06	0.17	0.44	1.31	0.02	2.20	3.87	0.00	0.00	51.45	4.33	55.78
1995	34.83	0.00	78.52	4.99	0.04	0.07	0.59	0.03	2.28	1.92	0.00	0.00	42.73	2.02	44.75
1996	32.99	0.00	86.68	0.66	0.12	0.24	0.40	0.03	2.00	2.40	0.00	0.00	36.17	2.67	38.84
1997	33.61	0.00	81.31	2.99	0.06	0.07	0.52	0.04	2.43	5.92	0.00	0.00	39.61	6.03	45.64
1998	34.23	0.00	79.30	0.35	0.12	0.18	0.52	0.03	2.32	2.36	0.00	0.00	37.54	2.57	40.11
1999	34.35	0.00	86.27	0.40	0.11	0.26	0.45	0.04	2.89	3.29	0.00	0.00	38.20	3.59	41.79
2000	39.88	0.00	78.74	0.41	0.09	0.55	0.43	0.03	2.39	3.00	0.00	0.00	43.20	3.58	46.78

Public supply transfers represent water withdrawn in Hillsborough and Pasco Counties and imported for use in Pinellas County.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Polk County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee; all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational Irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	26.40	0.00	0.00	0.00	270.00	0.00	51.80	9.10	----	----	0.10	265.00	350.44	274.10	624.54
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	27.70	0.00	0.00	7.02	236.00	71.00	156.10	17.30	----	----	0.10	110.00	426.92	198.30	625.22
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	31.23	0.00	0.00	9.30	270.38	1.85	96.89	5.10	----	----	0.10	298.50	407.90	305.45	713.35
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	30.37	0.00	0.00	9.30	233.43	0.00	91.34	5.65	----	----	0.20	222.90	364.64	228.55	593.19
1978	32.50	0.00	0.00	9.36	223.60	4.10	85.80	4.40	----	----	0.10	168.00	351.36	176.50	527.86
1979	34.90	0.00	0.00	11.32	199.91	23.60	78.07	3.99	----	----	0.10	221.60	324.30	249.19	573.49
1980	35.54	0.00	0.00	13.84	208.71	0.00	53.96	3.52	----	----	0.20	295.00	312.25	298.52	610.77
1981	41.38	0.00	0.00	14.53	187.41	14.70	125.93	7.55	----	----	0.30	315.00	369.55	337.25	706.80
1982	65.15	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1983	46.53	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1984	50.80	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1985	54.90	0.00	0.00	18.17	135.93	1.14	105.92	7.58	4.06	0.75	1.38	68.97	320.36	78.44	398.80
1986	57.15	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	60.17	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	64.75	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	65.61	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	65.52	0.00	0.00	23.26	143.35	0.00	114.35	11.94	5.78	1.27	1.41	70.62	353.67	83.83	437.50
1991	69.43	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	67.34	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	63.18	0.19	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	58.52	0.30	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	58.15	0.27	0.00	7.67	67.61	24.12	100.14	4.32	8.71	1.38	0.70	118.78	242.98	148.87	391.85
1996	63.26	0.22	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	65.68	0.21	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	71.03	0.14	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	71.22	0.15	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	75.43	0.06	0.00	12.45	71.20	6.39	157.53	9.33	9.65	2.29	4.27	15.66	330.53	33.73	364.26

Polk County is located within the St. Johns River, South Florida and the Southwest Florida Water Management Districts. The values shown represent the County totals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District and the Southwest Florida Water Management District.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004), St. Johns River Water Management District (2004), and the Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Putnam County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	2.00	0.00	0.92	0.00	3.00	54.00	11.81	1.52	----	----	0.06	13.00	17.79	68.52	86.31
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	2.70	0.00	2.69	0.00	15.50	16.00	7.60	2.10	----	----	0.05	123.00	28.54	141.10	169.64
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	2.58	0.00	0.00	0.00	16.20	21.00	18.86	0.18	----	----	0.06	120.00	40.61	141.18	181.79
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	3.14	0.00	0.00	0.00	16.20	21.00	18.96	4.71	----	----	0.17	1.20	41.50	26.91	68.41
1978	2.50	0.00	3.96	0.00	19.23	21.00	25.47	3.05	----	----	0.17	0.33	51.33	24.38	75.71
1979	2.55	0.00	7.16	0.00	20.50	23.14	48.10	5.58	----	----	0.33	0.69	78.64	29.41	108.05
1980	2.86	0.00	4.76	0.00	19.23	21.00	32.97	3.90	----	----	0.34	0.69	60.16	25.59	85.75
1981	2.77	0.00	8.16	0.00	29.38	8.07	24.31	1.10	----	----	0.24	0.41	64.86	9.58	74.44
1982	3.45	0.00	7.40	0.00	19.02	36.57	20.58	1.71	----	----	0.31	0.42	50.76	38.70	89.46
1983	3.06	0.00	6.64	0.00	16.15	39.27	21.01	1.67	----	----	0.36	1.70	47.22	42.64	89.86
1984	2.80	0.00	6.09	0.00	26.02	12.82	21.42	1.10	----	----	0.62	1.77	56.95	15.69	72.64
1985	2.97	0.00	6.35	0.00	43.07	9.98	16.92	0.96	0.28	0.00	0.68	9.13	70.27	20.07	90.34
1986	3.18	0.00	6.44	0.00	37.37	10.20	18.38	1.52	0.33	0.00	0.53	12.09	66.23	23.81	90.04
1987	3.28	0.00	6.54	0.00	42.71	4.00	15.54	1.04	0.26	0.00	0.52	11.62	68.85	16.66	85.51
1988	3.44	0.00	6.34	0.00	43.64	4.00	18.53	1.14	0.24	0.00	1.26	12.04	73.45	17.18	90.63
1989	3.38	0.00	6.23	0.00	43.74	4.01	19.11	1.37	0.26	0.00	0.45	12.31	73.17	17.69	90.86
1990	3.15	0.00	5.95	0.00	33.52	10.27	20.17	1.35	0.31	0.00	0.53	7.71	63.63	19.33	82.96
1991	3.59	0.00	6.71	0.00	23.35	26.37	13.14	0.64	0.27	0.00	0.61	14.93	47.67	41.94	89.61
1992	3.70	0.00	6.94	0.00	21.67	37.66	16.54	0.84	0.09	0.00	0.46	14.19	49.40	52.69	102.09
1993	3.71	0.00	6.58	0.00	18.94	33.14	12.12	0.84	0.21	0.00	0.55	12.27	42.11	46.25	88.36
1994	3.41	0.00	6.05	0.00	8.44	33.94	11.56	0.66	0.20	0.00	0.66	12.56	30.32	47.16	77.48
1995	3.59	0.00	8.23	0.00	11.19	34.74	14.20	1.08	0.20	0.00	0.70	14.50	38.11	50.32	88.43
1996	3.91	0.00	8.60	0.00	20.36	17.26	16.14	1.12	0.32	0.00	0.55	15.91	49.88	34.29	84.17
1997	3.25	0.00	6.72	0.00	17.54	17.47	13.17	0.70	0.32	0.00	0.60	17.99	41.60	36.16	77.76
1998	3.87	0.00	8.48	0.00	19.55	33.98	14.76	0.77	0.36	0.00	0.61	16.33	47.63	51.08	98.71
1999	2.96	0.00	5.78	0.00	24.16	33.65	15.31	0.88	0.27	0.00	0.58	14.39	49.06	48.92	97.98
2000	3.20	0.00	4.99	0.00	19.14	31.12	11.69	3.50	0.64	0.40	0.69	13.90	40.35	48.92	89.27

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources; 1965 -1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991 -1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in St. Johns County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.20	1.80	0.00	0.00	0.00	0.00	13.98	0.02	----	----	0.00	0.00	15.78	1.82	17.60
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	1.40	1.10	0.00	0.00	0.00	0.00	22.10	0.00	----	----	0.00	0.00	25.18	1.10	26.28
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	2.67	0.00	0.00	0.00	2.00	0.00	28.82	0.10	----	----	0.00	0.00	35.84	0.10	35.94
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	2.89	0.00	0.00	0.00	2.00	0.00	33.58	0.05	----	----	0.00	0.00	41.14	0.05	41.19
1978	2.73	0.00	0.00	0.00	2.94	0.00	22.96	0.09	----	----	0.00	0.00	28.63	0.09	28.72
1979	2.83	0.00	0.00	0.00	2.73	0.00	29.47	0.49	----	----	0.00	0.00	35.03	0.49	35.52
1980	3.50	0.00	0.00	0.00	2.38	0.00	24.85	0.49	----	----	0.00	0.00	30.75	0.49	31.24
1981	4.96	0.00	0.00	0.00	2.62	0.00	29.86	0.49	----	----	0.00	0.00	37.45	0.49	37.94
1982	5.69	0.00	0.00	0.00	2.60	0.00	32.55	0.49	----	----	0.00	0.00	41.06	0.49	41.55
1983	6.12	0.00	0.00	0.00	1.80	0.00	36.10	0.72	----	----	0.00	0.00	44.06	0.72	44.78
1984	6.98	0.00	0.00	0.00	2.52	0.00	34.49	1.43	----	----	0.00	0.00	44.05	1.43	45.48
1985	7.01	0.00	0.00	0.00	2.13	0.00	38.98	0.26	2.02	0.81	0.00	0.00	50.22	1.07	51.29
1986	7.28	0.00	0.00	0.00	2.13	0.00	37.88	0.10	2.17	1.19	0.00	0.00	49.53	1.29	50.82
1987	7.61	0.00	0.00	0.00	2.23	0.00	25.57	0.10	1.72	0.98	0.00	0.00	37.19	1.08	38.27
1988	7.49	0.00	0.00	0.00	2.12	0.00	33.41	0.10	1.41	0.80	0.00	0.00	44.47	0.90	45.37
1989	7.79	0.00	0.00	0.00	2.10	0.00	35.97	0.10	1.74	1.00	0.00	0.00	47.65	1.10	48.75
1990	8.39	0.00	0.00	0.00	2.24	0.00	38.29	0.10	2.25	1.29	0.00	0.00	51.26	1.39	52.65
1991	7.63	0.00	0.00	0.00	2.62	0.00	28.60	0.00	1.97	1.12	0.00	0.00	40.89	1.12	42.01
1992	8.62	0.00	0.00	0.00	2.50	0.00	35.57	0.00	0.62	0.35	0.00	0.00	47.39	0.35	47.74
1993	11.54	0.00	0.00	0.00	0.49	0.00	27.00	0.00	1.77	1.03	0.00	0.00	40.86	1.03	41.89
1994	9.72	0.00	0.00	0.00	0.50	0.00	28.20	0.00	1.75	1.02	0.00	0.00	40.22	1.02	41.24
1995	10.30	0.00	0.00	0.00	2.89	0.00	31.35	0.00	1.13	0.64	0.00	0.00	45.73	0.64	46.37
1996	11.53	0.00	0.00	0.00	2.69	0.00	28.24	0.00	1.65	0.93	0.00	0.00	44.16	0.93	45.09
1997	12.32	0.00	0.00	0.00	2.80	0.00	26.23	0.00	1.48	0.84	0.00	0.00	42.89	0.84	43.73
1998	13.88	0.00	0.00	0.00	2.82	0.00	32.73	0.00	1.84	1.04	0.00	0.00	51.33	1.04	52.37
1999	14.89	0.00	0.00	0.00	3.10	0.00	32.36	0.00	1.34	0.77	0.00	0.00	51.75	0.77	52.52
2000	16.49	0.00	0.00	0.00	1.91	0.00	28.59	0.00	5.55	3.16	0.00	0.00	52.55	3.16	55.71

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

- Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.
- Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.
- Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).
- 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.
- 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).
- 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in St. Lucie County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	3.20	0.00	1.70	0.00	0.80	0.00	46.40	139.30	----	----	0.00	0.00	52.10	139.30	191.40
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	4.30	0.00	2.02	0.00	1.20	0.00	25.90	159.50	----	----	0.00	0.00	33.42	159.50	192.92
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	6.14	0.00	3.97	0.00	0.19	0.00	49.60	319.31	----	----	0.00	0.00	59.90	319.31	379.21
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	6.65	0.00	4.64	0.00	0.19	0.00	64.57	332.55	----	----	0.00	0.00	76.05	332.55	408.60
1978	5.69	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	9.69	0.00	3.18	0.00	2.05	0.00	48.89	183.13	----	----	0.00	0.00	63.81	183.13	246.94
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	10.83	0.00	6.40	0.00	2.25	0.00	48.03	162.85	0.77	0.77	0.00	0.00	68.28	163.62	231.90
1986	11.06	0.00	0.09	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	12.38	0.00	0.14	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	13.10	0.00	0.21	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	14.34	0.00	0.25	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	14.39	0.00	11.49	0.00	1.59	1.50	52.04	166.87	4.33	2.29	0.00	0.00	83.84	170.66	254.50
1991	14.13	0.00	0.45	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	14.57	0.00	0.43	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	14.50	0.00	0.72	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	13.92	0.00	0.85	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	15.31	0.00	9.15	0.00	0.07	5.03	51.53	222.10	4.70	1.98	0.00	0.00	80.76	229.11	309.87
1996	15.54	0.00	1.26	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	15.69	0.00	1.86	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	15.85	0.00	1.87	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	16.40	0.00	2.83	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	17.95	0.00	5.65	0.00	6.35	2.53	45.16	192.18	3.99	4.21	0.00	0.00	80.84	198.92	279.76

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Data sources: 1965 -1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991 -1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Santa Rosa County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.10	0.00	1.50	0.00	7.20	0.00	0.40	0.02	---	---	0.00	0.00	10.20	0.02	10.22
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	2.40	0.00	2.75	0.00	10.30	0.00	0.20	0.00	---	---	0.00	0.00	15.65	0.00	15.65
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	3.40	0.00	1.05	0.00	17.67	0.00	0.40	0.17	---	---	0.00	0.00	22.52	0.17	22.69
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	4.39	0.00	1.18	0.00	18.99	0.00	1.56	0.21	---	---	0.00	0.00	26.12	0.21	26.33
1978	4.56	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	5.83	0.00	0.68	0.00	18.52	0.00	0.69	0.32	---	---	0.00	0.00	25.72	0.32	26.04
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	7.80	0.00	0.82	0.00	6.59	0.00	1.20	0.15	0.00	0.00	0.00	0.00	16.41	0.15	16.56
1986	8.70	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	9.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	8.88	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	8.79	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	9.63	0.00	0.58	0.00	6.48	0.00	2.08	0.22	3.83	0.07	0.00	0.00	22.60	0.29	22.89
1991	9.41	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	10.62	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	10.78	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	10.62	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	11.28	0.00	0.43	0.00	6.02	0.00	0.81	0.15	3.45	0.07	0.00	0.00	22.23	0.22	22.45
1996	11.05	0.00	1.03	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	11.34	0.00	1.15	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	12.57	0.00	1.17	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	12.42	0.00	1.05	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	13.47	0.00	1.07	0.81	5.57	0.00	5.00	0.00	6.00	0.35	0.00	0.00	30.85	0.35	31.20

Public supply transfers represent water withdrawn in Escambia County and imported for use in Santa Rosa County.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data source; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Sarasota County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	6.70	0.20	0.00	0.00	0.40	0.00	40.00	0.80	----	----	0.00	0.00	47.98	1.00	48.98
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	10.90	0.30	0.00	0.00	7.60	0.00	27.30	2.30	----	----	0.00	0.00	47.73	2.60	50.33
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	9.33	0.00	0.98	0.00	2.99	0.00	18.35	2.34	----	----	0.00	0.00	38.00	2.34	40.34
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	9.58	1.26	2.10	0.00	2.92	0.00	22.65	2.68	----	----	0.00	0.00	41.85	3.94	45.79
1978	9.10	1.40	6.30	0.00	0.30	0.00	30.95	3.52	----	----	0.00	0.00	45.81	4.92	50.73
1979	9.90	1.80	6.60	0.00	0.30	0.00	25.05	2.85	----	----	0.00	0.00	40.46	4.65	45.11
1980	11.10	1.30	7.17	0.00	0.10	0.00	20.89	0.47	----	----	0.00	0.00	33.17	1.77	34.94
1981	11.60	1.26	8.34	0.00	0.10	0.00	19.52	0.44	----	----	0.00	0.00	32.28	1.70	33.98
1982	14.53	0.00	9.46	0.00	0.45	0.00	21.51	0.44	----	----	0.00	0.00	37.14	0.44	37.58
1983	14.04	0.76	12.95	0.00	0.91	0.00	19.74	0.41	----	----	0.00	0.00	35.36	1.17	36.53
1984	13.70	0.74	11.92	0.00	0.71	0.00	23.58	0.48	----	----	0.00	0.00	39.03	1.22	40.25
1985	17.09	0.78	9.40	0.00	0.49	0.00	13.68	0.00	5.96	0.47	0.00	0.00	40.36	1.25	41.61
1986	19.14	1.58	10.20	0.00	0.05	0.00	23.42	0.48	----	----	0.00	0.00	44.78	2.06	46.84
1987	18.65	1.32	10.29	0.00	2.48	0.00	14.78	0.68	----	----	0.00	0.00	40.53	2.00	42.53
1988	17.43	0.00	11.67	0.00	4.55	0.00	15.48	3.17	----	----	0.00	0.00	40.36	3.17	43.53
1989	17.92	0.00	13.25	0.00	2.81	0.48	15.42	2.49	----	----	0.00	0.00	39.86	2.97	42.83
1990	21.43	0.00	14.18	0.00	1.87	0.48	27.48	2.39	4.85	0.31	0.00	0.00	59.49	3.18	62.67
1991	28.59	0.00	14.16	0.00	0.39	3.76	18.54	1.73	2.78	2.95	0.00	0.00	53.55	8.44	61.99
1992	27.01	0.00	11.69	0.00	0.37	4.27	7.46	4.93	3.00	3.15	0.00	0.00	42.18	12.35	54.53
1993	22.89	2.73	11.63	0.00	0.14	2.91	6.00	2.30	3.36	2.70	0.00	0.00	42.29	10.64	52.93
1994	21.36	1.28	12.56	0.00	0.10	4.03	3.47	2.48	3.41	2.74	0.00	0.00	34.14	10.53	44.67
1995	25.21	0.77	10.20	0.00	0.07	5.05	3.41	0.24	2.98	2.96	0.00	0.00	32.34	9.02	41.36
1996	23.00	1.30	10.66	0.00	0.09	2.25	5.07	0.84	3.28	3.68	0.00	0.00	34.73	8.07	42.80
1997	24.64	1.01	10.21	0.00	0.11	1.02	5.26	0.43	4.23	3.32	0.00	0.00	38.81	5.78	44.59
1998	25.15	1.19	10.23	0.00	0.11	0.66	5.21	0.59	4.27	3.81	0.00	0.00	35.99	6.25	42.24
1999	23.85	1.50	10.89	0.00	0.29	1.13	5.41	0.68	4.64	3.24	0.00	0.00	35.18	6.55	41.73
2000	27.56	0.85	10.62	0.43	0.31	0.33	6.88	0.66	4.89	4.07	0.00	0.00	40.07	5.91	45.98

Public supply transfers represent water withdrawn in DeSoto and Manatee Counties and imported for use in Sarasota County.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Seminole County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	4.20	0.00	1.81	0.04	0.00	0.00	4.90	5.98	----	----	0.00	0.00	10.91	6.02	16.93
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	6.30	0.00	2.65	0.00	0.50	0.00	3.40	2.70	----	----	0.00	0.00	12.85	2.70	15.55
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	10.50	0.00	8.05	0.00	2.59	0.00	10.98	0.00	----	----	0.00	0.00	32.12	0.00	32.12
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	12.60	0.00	8.63	0.00	2.59	0.00	12.97	0.67	----	----	0.00	0.00	36.79	0.67	37.46
1978	13.98	0.00	8.54	0.00	3.77	0.00	18.38	0.92	----	----	0.00	0.00	44.67	0.92	45.59
1979	18.78	0.00	7.57	0.00	3.10	0.00	27.53	1.69	----	----	0.00	0.00	56.98	1.69	58.67
1980	25.65	0.00	11.68	0.00	4.13	0.00	18.68	0.40	----	----	0.00	0.00	60.14	0.40	60.54
1981	25.97	0.00	5.67	0.00	4.03	0.00	20.07	0.40	----	----	0.00	0.00	55.74	0.40	56.14
1982	25.16	0.00	5.63	0.00	3.37	0.00	26.94	0.40	----	----	0.00	0.00	61.10	0.40	61.50
1983	28.19	0.00	3.30	0.00	4.39	0.00	32.80	0.40	----	----	0.00	0.00	68.68	0.40	69.08
1984	31.85	0.00	4.62	0.00	5.29	0.00	31.49	0.75	----	----	0.00	0.00	73.25	0.75	74.00
1985	34.86	0.00	3.63	0.00	4.96	0.00	20.87	1.18	3.52	0.88	0.00	0.00	67.84	2.06	69.90
1986	38.68	0.00	3.72	0.00	2.54	0.00	20.16	1.18	4.98	1.13	0.00	0.00	70.08	2.31	72.39
1987	40.90	0.00	4.12	0.00	0.62	0.00	16.62	1.21	4.05	0.96	0.00	0.00	66.31	2.17	68.48
1988	42.86	0.00	5.03	0.00	0.61	0.00	15.16	1.19	3.54	0.83	0.00	0.00	67.20	2.02	69.22
1989	47.39	0.00	5.00	0.00	0.53	0.00	8.33	1.10	4.37	1.02	0.00	0.00	65.62	2.12	67.74
1990	50.79	0.00	3.14	0.00	0.49	0.00	8.25	1.10	2.90	0.70	0.00	0.00	65.57	1.80	67.37
1991	46.44	0.00	0.62	0.00	0.53	0.00	5.53	0.14	4.25	1.02	0.00	0.00	57.37	1.16	58.53
1992	47.15	0.00	2.32	0.00	0.43	0.00	5.61	0.07	1.62	0.38	0.00	0.00	57.13	0.45	57.58
1993	49.10	0.00	1.36	0.00	0.41	0.00	6.55	0.09	4.01	1.01	0.00	0.00	61.43	1.10	62.53
1994	48.49	0.00	4.13	0.00	0.17	0.00	6.65	0.19	3.98	1.00	0.00	0.00	63.42	1.19	64.61
1995	50.69	0.00	8.63	0.00	0.14	0.00	6.99	0.26	2.46	0.62	0.00	0.00	68.91	0.88	69.79
1996	50.76	0.00	5.64	0.00	0.15	0.00	6.25	0.21	3.74	0.89	0.00	0.00	66.54	1.10	67.64
1997	54.82	0.00	5.95	0.00	0.16	0.00	6.73	0.37	4.12	0.98	0.00	0.00	71.78	1.35	73.13
1998	61.01	0.00	5.90	0.00	0.15	0.00	7.67	0.22	4.70	1.11	0.00	0.00	79.43	1.33	80.76
1999	60.08	0.00	3.89	0.00	0.09	0.00	4.39	0.14	3.07	0.73	0.00	0.00	71.52	0.87	72.39
2000	66.90	0.00	2.73	0.00	0.08	0.00	11.06	0.02	7.70	1.76	0.00	0.00	88.47	1.78	90.25

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Sumter County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.50	0.00	0.79	0.00	26.00	0.00	3.97	0.21	---	---	0.00	0.00	31.26	0.21	31.47
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.80	0.00	1.20	0.00	18.50	0.00	4.60	0.00	---	---	0.00	0.00	25.10	0.00	25.10
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	0.61	0.00	1.35	0.00	16.06	0.00	3.98	0.17	---	---	0.00	0.00	22.00	0.17	22.17
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.92	0.00	1.45	0.00	16.06	0.00	3.96	0.17	---	---	0.00	0.00	22.39	0.17	22.56
1978	0.90	0.00	1.57	0.00	0.10	0.00	3.94	0.20	---	---	0.00	0.00	6.51	0.20	6.71
1979	0.80	0.00	1.63	0.00	2.32	0.00	16.53	0.83	---	---	0.00	0.00	21.28	0.83	22.11
1980	1.00	0.00	1.89	0.00	1.14	0.00	17.01	0.34	---	---	0.00	0.00	21.04	0.34	21.38
1981	1.10	0.00	1.93	0.00	10.50	0.00	19.72	0.39	---	---	0.00	0.00	33.25	0.39	33.64
1982	1.52	0.00	1.39	0.00	45.00	0.00	13.64	0.28	---	---	0.00	0.00	61.55	0.28	61.83
1983	1.08	0.00	1.53	0.00	32.34	0.00	11.62	0.24	---	---	0.00	0.00	46.57	0.24	46.81
1984	1.18	0.00	2.60	0.00	32.87	0.00	19.73	0.40	---	---	0.00	0.00	56.38	0.40	56.78
1985	1.31	0.00	2.41	0.00	57.76	0.00	16.25	0.35	0.54	0.00	0.00	0.00	78.27	0.35	78.62
1986	1.25	0.00	2.50	0.00	39.03	0.00	15.73	0.32	---	---	0.00	0.00	58.51	0.32	58.83
1987	1.13	0.00	2.29	0.00	44.70	0.00	8.47	0.26	---	---	0.00	0.00	56.59	0.26	56.85
1988	1.94	0.00	3.68	0.00	82.58	0.00	5.78	0.20	---	---	0.00	0.00	93.98	0.20	94.18
1989	2.01	0.00	3.83	0.00	69.89	0.00	7.40	0.38	---	---	0.00	0.00	83.13	0.38	83.51
1990	1.90	0.00	3.00	0.00	60.02	0.00	7.59	0.47	0.62	0.00	0.00	0.00	73.13	0.47	73.60
1991	2.36	0.00	1.85	0.00	0.63	56.85	25.05	1.84	0.33	0.49	0.00	0.00	30.22	0.49	30.71
1992	2.42	0.00	1.78	0.00	0.64	64.54	27.02	1.04	0.27	0.52	0.00	0.00	32.13	0.52	32.65
1993	2.33	0.00	1.88	0.00	0.12	50.23	9.00	0.21	0.58	0.00	0.00	0.00	13.91	0.58	14.49
1994	2.70	0.00	1.49	0.00	0.11	30.80	6.70	0.21	1.07	0.00	0.00	0.00	12.07	1.07	12.54
1995	2.45	0.00	0.93	0.00	0.20	53.39	7.76	0.20	1.21	0.00	0.00	0.00	12.55	1.21	13.76
1996	2.52	0.00	2.14	0.00	0.21	26.94	8.68	0.36	1.19	0.00	0.00	0.00	14.74	1.19	15.93
1997	2.76	0.00	3.05	0.00	0.25	23.85	9.12	0.33	1.68	0.00	0.00	0.00	16.86	1.68	18.54
1998	4.04	0.00	2.51	0.00	0.26	35.32	9.57	0.40	1.54	0.21	0.00	0.00	17.92	1.75	19.67
1999	4.61	0.00	5.10	0.00	0.35	33.54	9.30	0.38	2.78	0.00	0.00	0.00	22.14	2.78	24.92
2000	4.44	0.00	4.57	0.00	0.36	16.98	14.08	0.54	3.35	0.17	0.00	0.00	26.80	3.52	27.32

Water withdrawals for mining operations were classified as ground water between 1965 and 1990, and were re-classified as surface water beginning in 1991.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965-1984 and 1986-1989.

Data source: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee, and published data from the Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 1. Freshwater withdrawals by category in Suwannee County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.60	0.00	0.50	0.03	0.00	0.00	0.56	0.39	----	----	0.01	180.02	1.67	180.44	182.11
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.60	0.00	0.94	0.00	7.10	0.00	4.10	0.00	----	----	0.01	173.00	12.75	173.00	185.75
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	1.13	0.00	0.99	0.00	2.39	0.00	1.95	0.02	----	----	0.01	172.80	6.47	172.82	179.29
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	1.23	0.00	1.29	0.00	1.98	0.00	8.05	0.38	----	----	0.01	172.80	12.56	173.18	185.74
1978	1.13	0.00	1.42	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	1.06	0.00	1.54	0.00	0.79	0.00	8.38	1.22	----	----	0.01	172.80	11.78	174.02	185.80
1981	1.20	0.00	----	----	0.90	0.00	----	----	----	----	----	----	N/A	N/A	N/A
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	1.33	0.00	2.45	0.00	0.95	1.43	16.33	0.22	0.16	0.00	0.08	64.95	21.30	66.60	87.90
1986	1.35	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	1.33	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	1.24	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	1.32	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	1.36	0.00	2.95	0.00	0.81	0.30	27.06	0.65	0.05	0.00	0.07	108.51	32.30	109.46	141.76
1991	1.31	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	1.31	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	1.44	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1.46	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1.42	0.00	3.25	0.00	1.22	0.00	23.15	0.41	0.05	0.00	0.14	112.97	29.23	113.38	142.61
1996	1.48	0.00	----	----	1.24	0.00	----	----	----	----	0.17	97.60	N/A	N/A	N/A
1997	1.30	0.00	----	----	1.25	0.00	----	----	----	----	0.19	120.00	N/A	N/A	N/A
1998	1.43	0.00	----	----	1.27	0.00	----	----	----	----	0.15	116.89	N/A	N/A	N/A
1999	1.38	0.00	----	----	1.38	0.00	----	----	----	----	0.16	122.93	N/A	N/A	N/A
2000	1.40	0.00	2.70	0.00	1.54	0.00	20.64	0.38	0.07	0.02	0.06	101.01	26.41	101.41	127.82

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Taylor County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee. all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	1.00	0.00	0.20	0.00	50.00	0.00	0.12	0.08	---	---	0.00	0.00	51.32	0.08	51.40
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	1.20	0.00	0.38	0.00	53.70	0.00	0.10	0.00	---	---	0.00	0.00	55.38	0.00	55.38
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	1.37	0.00	0.42	0.00	57.02	0.00	0.29	0.13	---	---	0.00	0.00	59.10	0.13	59.23
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	1.67	0.00	0.53	0.00	48.40	0.00	0.17	0.16	---	---	0.00	0.00	50.77	0.16	50.93
1978	1.45	0.00	0.91	0.00	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	1.49	0.00	0.84	0.00	51.05	0.00	0.15	0.13	---	---	0.00	0.00	53.53	0.13	53.66
1981	---	---	---	---	48.27	0.00	---	---	---	---	0.00	0.00	N/A	N/A	N/A
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	1.58	0.00	1.13	0.00	45.21	1.63	0.61	0.00	0.00	0.00	0.00	0.00	48.53	1.63	50.16
1986	1.50	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	1.59	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	1.30	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	1.46	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	1.42	0.00	0.99	0.00	46.37	1.20	0.47	0.08	0.13	0.00	0.00	0.00	49.38	1.28	50.66
1991	1.44	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	1.32	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	1.24	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	1.39	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	1.93	0.00	1.33	0.00	47.40	1.98	0.33	0.00	0.11	0.00	0.00	0.00	51.10	1.98	53.08
1996	1.73	0.00	---	---	45.59	1.85	---	---	---	---	---	---	N/A	N/A	N/A
1997	1.70	0.00	---	---	44.37	1.79	---	---	---	---	---	---	N/A	N/A	N/A
1998	1.80	0.00	---	---	42.12	1.93	---	---	---	---	---	---	N/A	N/A	N/A
1999	1.79	0.00	---	---	43.75	1.73	---	---	---	---	---	---	N/A	N/A	N/A
2000	1.73	0.00	0.95	0.00	42.15	2.96	1.89	0.04	0.07	0.02	0.00	0.00	46.79	3.02	49.81

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Union County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.15	0.00	0.00	0.00	0.80	0.00	0.06	0.09	----	----	0.00	0.00	1.21	0.09	1.30
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.10	0.00	0.00	0.00	0.60	0.00	0.09	0.02	----	----	0.00	0.00	1.57	0.02	1.59
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.55	0.00	0.00	0.00	0.00	0.00	0.23	0.02	----	----	0.00	0.00	1.65	0.02	1.67
1976	0.55	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1977	0.56	0.00	0.00	0.00	0.00	0.00	0.26	0.13	----	----	0.00	0.00	1.22	0.13	1.35
1978	0.59	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	0.56	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1980	0.57	0.00	0.00	0.00	0.00	0.00	0.17	0.09	----	----	0.00	0.00	1.21	0.09	1.30
1981	0.50	0.00	0.00	0.00	----	----	----	----	----	----	0.00	0.00	N/A	N/A	N/A
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	0.52	0.00	0.00	0.00	0.02	0.00	0.71	0.00	0.00	0.00	0.00	0.00	2.11	0.00	2.11
1986	0.65	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	0.61	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	0.62	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	0.61	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	0.63	0.00	0.00	0.00	1.18	0.00	1.23	0.68	0.00	0.00	0.00	0.00	3.05	0.68	3.73
1991	0.62	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	0.33	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	0.31	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	0.29	0.00	0.00	0.00	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	0.38	0.00	0.00	0.00	0.82	0.00	0.35	0.00	0.00	0.00	0.00	0.00	2.57	0.13	2.70
1996	0.40	0.00	0.00	0.00	0.30	0.00	----	----	----	----	----	----	N/A	N/A	N/A
1997	0.45	0.00	0.00	0.00	0.31	0.00	----	----	----	----	----	----	N/A	N/A	N/A
1998	0.54	0.00	0.00	0.00	0.35	0.00	----	----	----	----	----	----	N/A	N/A	N/A
1999	0.39	0.00	0.00	0.00	0.37	0.00	----	----	----	----	----	----	N/A	N/A	N/A
2000	0.36	0.00	0.00	0.00	1.10	0.00	1.05	0.02	0.00	0.00	0.00	0.00	2.91	0.02	2.93

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Volusia County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	13.00	0.00	0.00	0.00	0.60	0.00	6.40	1.40	---	---	0.20	350.00	21.94	351.40	373.34
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	19.20	0.00	0.00	0.00	0.50	0.00	6.90	0.30	---	---	0.50	406.00	30.77	406.30	437.07
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	25.10	0.00	0.00	0.00	0.14	0.00	5.56	0.00	---	---	0.03	314.00	37.33	314.00	351.33
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	26.60	0.00	0.00	0.00	0.14	0.00	4.87	5.03	---	---	0.00	286.00	37.96	291.03	328.99
1978	26.49	0.00	0.00	0.00	0.31	0.00	13.47	5.32	---	---	0.32	283.77	44.53	289.09	333.62
1979	28.26	0.00	0.00	0.00	0.39	0.00	20.97	6.87	---	---	0.33	75.03	55.89	81.90	137.79
1980	30.22	0.00	0.00	0.00	0.54	0.00	22.90	2.46	---	---	0.55	154.10	58.74	156.56	215.30
1981	31.79	0.00	0.00	0.00	0.54	0.00	26.25	1.36	---	---	0.34	119.56	63.44	120.92	184.36
1982	31.23	0.00	0.00	0.00	0.32	0.00	26.11	1.36	---	---	0.34	116.85	63.12	118.21	181.33
1983	32.98	0.00	0.00	0.00	0.42	0.00	26.43	1.36	---	---	0.38	108.58	67.12	109.94	177.06
1984	34.58	0.00	0.00	0.00	0.55	0.00	23.01	4.85	---	---	0.25	86.55	64.05	91.40	155.45
1985	36.39	0.00	0.00	0.00	0.64	0.00	32.39	5.32	4.17	1.18	0.16	111.25	79.07	117.75	196.82
1986	37.74	0.00	0.00	0.00	0.94	0.00	28.75	5.31	4.84	1.41	0.36	116.92	77.98	123.64	201.62
1987	40.30	0.00	0.00	0.00	1.07	0.00	28.14	5.40	3.41	1.18	0.36	117.42	78.71	124.00	202.71
1988	43.67	0.00	0.00	0.00	0.55	0.00	28.29	5.52	2.84	1.01	0.41	117.45	81.18	123.98	205.16
1989	43.68	0.00	0.00	0.00	0.83	0.00	27.75	5.13	3.32	1.18	0.43	118.64	81.41	124.95	206.36
1990	44.21	0.00	0.00	0.00	0.69	0.00	19.33	3.24	3.70	1.26	0.44	198.85	75.12	203.35	278.47
1991	42.06	0.00	0.00	0.00	0.82	0.00	16.93	2.77	3.78	1.27	0.51	118.19	66.73	122.23	188.96
1992	44.14	0.00	0.00	0.00	0.85	0.00	17.70	2.75	1.27	0.45	0.34	116.06	70.30	119.26	189.56
1993	46.63	0.00	0.00	0.00	0.83	0.00	17.91	2.78	3.41	1.09	0.59	117.69	77.40	121.56	198.96
1994	46.27	0.00	0.00	0.00	0.67	0.00	16.32	2.42	3.36	1.07	0.57	117.70	75.72	121.19	196.91
1995	48.78	0.00	0.00	0.00	0.69	0.00	25.53	4.45	2.16	0.69	0.37	70.30	81.16	75.44	156.60
1996	49.59	0.00	0.00	0.00	5.20	0.00	27.69	4.49	4.28	1.50	0.42	1.80	89.55	7.79	97.34
1997	51.28	0.00	0.00	0.00	0.40	0.00	21.05	3.07	4.31	1.50	0.43	3.06	80.10	7.63	87.73
1998	55.90	0.00	0.00	0.00	0.56	0.00	24.95	3.34	5.56	1.95	0.51	2.94	90.41	8.23	98.64
1999	55.50	0.00	0.00	0.00	0.58	0.00	22.05	3.58	3.52	1.26	0.51	3.39	84.90	8.23	93.13
2000	54.90	0.00	0.00	0.00	0.49	0.00	29.03	3.88	9.04	3.21	0.54	130.63	97.02	137.72	234.74

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Power generation withdrawals for 1996-99 do not include all fresh surface water withdrawn within the County.

Data sources: 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 1. Freshwater withdrawals by category in Wakulla County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-Industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.02	0.00	0.30	0.00	0.00	1.00	0.01	0.02	----	----	0.00	0.00	0.33	1.02	1.35
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.20	0.00	0.47	0.00	0.70	0.40	0.00	0.00	----	----	0.30	159.92	1.67	160.32	161.99
1971	----	----	----	----	----	----	----	----	----	----	0.32	N/A	N/A	N/A	N/A
1972	----	----	----	----	----	----	----	----	----	----	0.31	N/A	N/A	N/A	N/A
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.26	0.00	0.42	0.00	0.80	0.43	0.01	0.03	----	----	0.28	104.64	1.77	105.10	106.87
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	0.36	0.00	0.45	0.00	0.78	0.00	0.06	0.05	----	----	0.47	95.30	2.12	93.35	95.47
1978	0.37	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	0.44	0.00	0.69	0.00	0.81	0.00	0.02	0.03	----	----	0.06	86.70	2.02	86.73	88.75
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	0.59	0.00	0.65	0.00	0.55	0.00	0.04	0.00	0.00	0.00	0.01	28.97	1.84	28.97	30.81
1986	0.70	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	0.68	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	0.64	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	0.83	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	0.77	0.00	0.71	0.00	0.79	0.00	0.11	0.00	0.00	0.00	0.31	51.80	2.69	51.80	54.49
1991	0.77	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	0.84	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	0.92	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1.05	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1.06	0.00	1.05	0.00	0.63	0.00	0.42	0.00	0.23	0.00	0.21	68.92	3.60	68.92	72.52
1996	1.26	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	1.33	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	1.89	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	1.86	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	2.19	0.00	1.44	0.00	0.62	0.00	0.28	0.00	0.20	0.00	0.06	28.38	4.79	28.38	33.17

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals. Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes. Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984. Power generation withdrawals include all surface water withdrawn for power generation within the County as the water may have been reported as saline in past years. Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995). 1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee. 1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999). 2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Walton County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ---, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.90	0.00	0.50	0.00	0.00	0.00	0.17	0.03	---	---	0.00	0.00	1.57	0.03	1.60
1966	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1967	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1968	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1969	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1970	0.70	0.00	0.83	0.00	1.20	0.00	1.01	0.00	---	---	0.00	0.00	3.74	0.00	3.74
1971	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1972	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1973	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1974	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1975	1.08	0.00	0.74	0.00	0.41	0.00	0.67	0.33	---	---	0.00	0.00	2.90	0.33	3.23
1976	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1977	0.99	0.00	0.95	0.00	0.80	0.00	3.75	1.03	---	---	0.00	0.00	6.49	1.03	7.52
1978	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1979	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1980	1.58	0.00	0.86	0.00	0.74	0.00	8.19	1.14	---	---	0.00	0.00	11.37	1.14	12.51
1981	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1982	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1983	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1984	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
1985	2.99	0.00	0.48	0.00	0.69	0.00	1.05	0.39	0.84	0.13	0.00	0.00	6.05	0.52	6.57
1986	3.00	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1987	3.14	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1988	3.15	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1989	3.21	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1990	3.71	0.00	0.41	0.00	0.86	0.00	2.40	0.87	0.55	0.25	0.00	0.00	7.93	1.12	9.05
1991	3.57	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1992	3.97	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1993	4.37	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1994	4.53	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1995	4.35	0.00	0.64	0.00	0.92	0.00	4.05	0.83	0.63	0.35	0.00	0.00	10.59	1.18	11.77
1996	4.82	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1997	4.60	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1998	4.90	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
1999	6.58	0.00	---	---	---	---	---	---	---	---	---	---	N/A	N/A	N/A
2000	7.35	0.00	0.17	0.00	0.92	0.00	1.80	0.29	0.27	1.52	0.00	0.00	10.51	1.81	12.32

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 1. Freshwater withdrawals by category in Washington County, 1965-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1965	0.30	0.00	0.40	0.00	0.00	0.00	0.06	0.26	----	----	0.00	0.00	0.76	0.26	1.02
1966	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1967	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1968	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1969	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1970	0.40	0.00	0.92	0.00	0.00	0.00	0.02	0.00	----	----	0.00	0.00	1.34	0.00	1.34
1971	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1972	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1973	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1974	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1975	0.59	0.00	0.80	0.00	0.00	0.00	0.10	0.07	----	----	0.00	0.00	1.49	0.07	1.56
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	0.78	0.00	0.93	0.00	0.00	0.00	0.07	0.08	----	----	0.00	0.00	1.78	0.08	1.86
1978	0.81	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	0.92	0.00	0.94	0.00	0.04	0.00	0.02	0.07	----	----	0.00	0.00	1.92	0.07	1.99
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	1.03	0.00	1.15	0.00	0.00	0.00	0.34	0.14	0.55	0.00	0.00	0.00	3.07	0.14	3.21
1986	1.00	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	1.00	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	1.03	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	1.09	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	1.18	0.00	1.65	0.00	0.01	0.00	0.19	0.11	0.22	0.00	0.00	0.00	3.25	0.11	3.36
1991	1.52	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	1.20	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	1.49	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	1.09	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	1.14	0.00	1.73	0.00	0.13	0.00	0.97	0.27	0.35	0.00	0.00	0.00	4.32	0.27	4.59
1996	1.13	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	1.13	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	1.16	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	1.09	0.00	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	1.16	0.00	1.42	0.00	0.22	0.00	1.13	0.00	0.39	0.00	0.00	0.00	4.32	0.00	4.32

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1965 through 1984.

Data sources; 1965-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).


1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 2. Freshwater withdrawals by category for each Water Management District in Florida, 1975-2000

Water Management District	Page
Northwest Florida Water Management District	127
St. Johns River Water Management District	128
South Florida Water Management District	129
Southwest Florida Water Management District	130
Suwannee River Water Management District	131



Appendix 2. Freshwater withdrawals by category, Northwest Florida Water Management District, Florida, 1975-2000

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1975	64.72	34.76	18.70	0.85	73.24	67.27	14.14	7.39	----	----	1.30	490.10	172.10	600.37	772.47
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	75.65	38.25	24.71	0.11	78.07	60.75	14.53	10.70	----	----	2.20	452.20	195.16	562.01	757.17
1978	75.44	38.95	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	79.23	37.75	24.73	0.10	65.22	70.91	25.43	13.21	----	----	7.76	540.30	202.37	662.27	864.64
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	101.38	37.50	31.41	0.00	52.32	49.49	26.67	22.55	11.91	1.31	7.96	371.34	231.65	482.19	713.84
1986	104.92	43.08	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	107.84	40.83	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	112.45	41.55	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	112.26	40.06	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	117.03	41.40	31.95	0.00	52.27	53.69	46.23	20.03	15.24	1.18	8.13	350.22	270.85	466.52	737.37
1991	108.53	41.20	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	114.99	42.21	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	121.12	38.11	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	115.15	40.19	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	122.05	45.66	0.00	0.00	54.31	51.98	39.78	11.89	13.93	2.05	5.95	278.80	269.28	390.38	659.66
1996	123.85	43.01	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	123.96	46.37	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	136.47	38.45	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	135.79	47.26	0.00	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	145.90	46.17	0.00	21.75	51.86	26.30	36.32	9.94	18.57	4.50	6.41	317.80	280.81	404.71	685.52

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1975 through 1984.

Data sources; 1975-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 2. Freshwater withdrawals by category, St. Johns River Water Management District, Florida, 1975-2000 (Continued)

[U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1975	258.66	8.90	55.20	0.00	167.34	26.04	229.32	317.60	----	----	4.30	511.40	714.82	863.94	1,578.76
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	269.50	10.80	59.13	0.00	165.14	25.90	343.43	128.83	----	----	6.80	352.80	844.00	518.33	1,362.33
1978	239.35	9.46	75.28	0.00	158.71	21.00	440.82	226.40	----	----	3.99	306.70	918.15	563.56	1,481.71
1979	245.47	11.88	109.67	0.00	149.76	23.14	667.48	314.70	----	----	4.71	149.33	1,177.09	499.05	1,676.14
1980	281.90	12.97	85.37	0.00	137.79	25.57	416.91	190.77	----	----	4.94	117.39	926.91	346.70	1,273.61
1981	294.86	12.41	89.80	0.00	144.97	15.04	801.68	298.13	----	----	4.21	119.97	1,335.52	445.55	1,781.07
1982	280.19	11.33	88.20	0.00	128.82	40.87	484.01	214.76	----	----	4.05	117.27	985.27	384.23	1,369.50
1983	287.49	11.91	80.99	0.00	120.91	42.76	493.47	254.98	----	----	4.54	110.28	987.40	419.93	1,407.33
1984	318.85	13.21	87.72	0.00	137.33	12.91	492.58	261.32	----	----	4.55	88.32	1,041.03	375.76	1,416.79
1985	354.44	14.09	81.76	0.00	152.59	19.75	351.90	198.53	23.24	11.01	4.03	120.38	967.96	363.76	1,331.72
1986	366.52	15.47	82.33	0.00	127.96	20.50	370.90	201.53	32.43	13.11	4.71	129.01	984.85	379.62	1,364.47
1987	384.78	15.61	85.71	0.00	141.67	4.00	349.40	194.34	26.85	10.65	5.50	128.87	993.91	353.47	1,347.38
1988	393.14	16.15	86.73	0.00	146.11	4.00	375.95	219.11	25.46	10.40	6.29	129.49	1,033.68	379.15	1,412.83
1989	415.63	15.49	90.24	0.00	144.65	4.01	362.78	198.77	27.29	11.25	6.16	130.95	1,046.75	360.47	1,407.22
1990	427.90	16.24	85.98	0.00	126.09	10.27	347.41	214.08	27.20	10.79	6.75	206.56	1,021.33	457.94	1,479.27
1991	398.91	15.24	84.51	0.00	117.23	27.01	333.43	185.94	29.65	12.10	6.87	133.12	970.60	373.41	1,344.01
1992	409.80	14.83	84.92	0.00	109.82	38.38	344.25	280.35	12.03	5.41	6.18	130.25	967.00	469.22	1,436.22
1993	425.94	14.92	82.20	0.00	99.46	34.28	354.68	212.66	27.51	12.33	7.00	129.96	996.79	404.15	1,400.94
1994	418.35	15.71	85.35	0.00	90.79	35.08	357.76	210.28	27.23	12.29	12.11	130.26	991.59	403.62	1,395.21
1995	449.43	12.15	93.37	0.00	95.76	35.88	302.34	189.31	19.34	8.26	7.66	84.80	967.90	330.40	1,298.30
1996	475.07	11.07	86.05	0.00	108.19	17.99	313.27	189.30	28.20	12.23	19.07	17.71	1,029.85	248.30	1,278.15
1997	473.90	11.85	82.80	0.00	100.50	18.20	230.30	130.00	27.20	11.50	21.70	20.20	936.40	191.75	1,128.15
1998	517.82	12.15	88.25	0.00	97.03	34.49	417.26	197.30	32.77	13.69	7.89	19.27	1,161.02	276.90	1,437.92
1999	531.09	12.11	92.03	0.00	93.86	33.86	228.32	116.27	22.66	9.76	8.00	17.78	975.96	189.78	1,165.74
2000	558.51	14.08	51.70	0.00	90.64	31.80	392.33	214.20	72.66	31.94	10.86	144.53	1,176.70	436.55	1,613.25

Values presented in this table may not be identical to those reported or published by the water management district due to differences in data-collection procedures, categories of use, or revisions in reported or published values.

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1975 through 1984.

Surface-water values for power generation for 1991-94 and 1996-99 do not include all of the fresh surface water withdrawn for power generation within the District.

Data sources; 1975-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - Published water-use data from the St. Johns River Water Management District, Palatka.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and St. Johns River Water Management District (2004).

Appendix 2. Freshwater withdrawals by category, South Florida Water Management District, Florida, 1975-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power Generation		Total freshwater withdrawn		
	Ground	Surface	Treated	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total	
1975	513.78	42.83	----	48.99	1.20	18.26	54.35	576.63	1,274.71	----	1.36	0.00	1,159.02	1,373.09	2,532.11
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	549.11	39.62	----	42.90	0.90	22.73	55.14	675.70	1,288.23	----	1.89	0.00	1,292.33	1,383.89	2,676.22
1978	549.96	32.90	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	648.14	45.59	----	65.12	0.00	41.11	1.93	648.11	1,145.00	----	4.28	116.70	1,406.76	1,309.22	2,715.98
1981	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	746.22	37.37	10.82	59.92	0.00	48.80	1.56	498.53	1,020.45	47.91	46.23	24.50	1,402.06	1,130.11	2,532.17
1986	769.16	37.53	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	828.43	40.64	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	853.82	40.43	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	884.21	42.05	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	815.82	39.71	21.49	89.10	0.00	77.59	66.92	729.59	1,403.19	135.78	80.09	19.56	1,851.04	1,609.47	3,460.51
1991	848.46	40.80	17.69	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	897.46	40.85	18.78	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	913.97	38.72	24.81	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	923.13	37.93	35.74	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	933.59	39.38	37.29	74.81	0.00	66.58	54.52	696.84	1,496.75	127.43	59.37	19.28	1,902.05	1,669.30	3,571.35
1996	959.47	42.52	43.28	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1997	953.10	43.60	43.03	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1998	1,003.41	48.67	50.19	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1999	1,019.45	48.05	54.76	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
2000	1,070.28	49.24	73.66	52.43	0.00	88.83	35.18	784.37	1,682.22	86.21	123.68	24.63	2,085.07	1,914.95	4,000.02

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1975 through 1984.

Data sources; 1975-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Appendix 2. Freshwater withdrawals by category, Southwest Florida Water Management District, Florida, 1975-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-industrial-mining		Agricultural		Recreational irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1975	136.26	76.49	68.81	0.00	390.93	9.95	453.29	37.95	----	----	0.76	418.70	1,050.05	543.09	1,593.14
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	153.95	84.15	54.52	0.00	343.77	7.40	414.28	37.13	----	----	0.80	334.08	967.32	462.76	1,430.08
1978	176.36	72.77	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	194.95	72.04	45.15	0.00	344.27	31.51	408.87	23.70	----	----	0.80	307.80	994.04	435.05	1,429.09
1980	205.85	84.13	44.38	0.00	281.47	6.44	417.82	18.41	----	----	0.87	298.60	950.39	407.58	1,357.97
1981	222.29	96.06	45.64	0.00	289.50	24.16	582.55	31.06	----	----	1.13	392.00	1,141.11	543.28	1,684.39
1982	267.15	87.32	38.80	0.00	262.85	12.19	428.23	20.54	----	----	3.70	251.53	1,000.73	371.58	1,372.31
1983	256.30	89.78	57.97	0.00	251.02	2.71	438.23	22.29	----	----	3.60	175.69	1,007.12	290.47	1,297.59
1984	278.61	89.45	65.02	0.00	279.80	5.94	653.96	30.25	----	----	4.91	75.34	1,282.30	200.98	1,483.28
1985	286.50	104.65	62.88	0.00	284.87	1.14	592.50	27.92	35.67	3.29	3.24	80.59	1,264.66	217.59	1,482.25
1986	288.30	96.26	47.84	0.00	243.19	0.00	613.54	32.06	----	----	2.91	188.25	1,195.78	316.57	1,512.35
1987	298.97	104.01	69.08	0.00	267.60	0.06	472.78	24.80	----	----	2.79	116.53	1,111.22	245.40	1,356.62
1988	317.49	115.01	67.20	0.00	318.65	8.62	543.00	36.26	----	----	3.45	87.83	1,249.79	247.72	1,497.51
1989	327.54	120.17	84.36	0.00	319.31	4.84	624.69	69.45	----	----	2.01	63.49	1,357.91	257.95	1,615.86
1990	323.88	128.98	76.19	0.00	189.36	99.74	573.17	59.37	22.07	13.46	2.98	75.87	1,187.65	377.42	1,565.07
1991	334.49	143.58	32.18	0.00	147.35	228.58	486.77	85.26	23.52	33.65	2.20	1.36	1,026.51	492.43	1,518.94
1992	348.37	141.13	37.29	0.00	159.01	193.47	514.83	68.14	29.65	21.51	0.65	1.52	1,089.80	425.77	1,515.57
1993	337.03	123.15	90.00	0.00	138.72	102.52	433.92	20.29	37.57	20.07	1.16	0.11	1,038.40	266.14	1,304.54
1994	334.50	113.15	79.25	0.00	133.64	100.49	424.33	19.10	36.67	17.97	2.08	0.00	1,010.47	250.71	1,261.18
1995	336.21	112.51	72.24	0.00	127.52	107.35	408.19	15.70	34.68	14.82	2.70	118.79	981.54	369.17	1,350.71
1996	349.55	112.37	65.12	0.00	94.78	80.05	489.41	19.38	40.68	20.06	3.57	1.50	1,043.11	233.36	1,276.47
1997	363.68	119.22	71.89	0.00	122.77	67.62	465.07	17.80	45.32	21.54	3.81	1.88	1,072.54	228.06	1,300.60
1998	383.85	119.11	48.10	0.00	120.26	66.22	499.77	20.08	41.94	19.10	4.11	1.96	1,098.03	226.47	1,324.50
1999	385.94	132.02	50.62	0.00	115.69	57.45	524.79	19.36	48.29	18.97	6.36	2.07	1,131.69	229.87	1,361.56
2000	408.88	127.94	52.74	0.00	117.52	34.40	683.06	24.83	51.95	20.79	6.77	40.76	1,320.92	248.72	1,569.64

Values presented in this table may not be identical to those reported or published by the water management district due to differences in data-collection procedures, categories of use, or revisions in reported or published values.

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards. Water withdrawals for mining operations were classified as ground water between 1965 and 1990, and were re-classified as surface water beginning in 1991.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1975-1984 and 1986-1989. Surface-water values for power generation for 1991-94 and 1996-99 do not include all of the fresh surface water withdrawn for power generation within the District.

Data sources; 1975-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, and 1996-1999 - published water-use data, Southwest Florida Water Management District, Brooksville.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004) and Southwest Florida Water Management District (2002).

Appendix 2. Freshwater withdrawals by category, Suwannee River Water Management District, Florida, 1975-2000 (Continued)

[Compiled by the U.S. Geological Survey, Tallahassee, all values in million gallons per day; ----, no or partial data were available; N/A, totals not available]

Year	Public supply		Domestic self-supplied		Commercial-Industrial-mining		Agricultural		Recreational Irrigation		Power generation		Total freshwater withdrawn		
	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Ground	Surface	Total
1975	9.42	0.00	9.23	0.00	95.60	3.09	16.46	3.01	----	----	0.61	172.80	131.32	178.90	310.22
1976	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1977	10.86	0.00	10.53	0.00	93.97	4.30	29.32	4.48	----	----	1.11	172.80	145.79	181.58	327.37
1978	11.45	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1979	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1980	11.05	0.00	13.35	0.00	94.18	0.00	23.26	5.87	----	----	0.94	172.80	142.78	178.67	321.45
1981	10.88	0.00	14.90	0.00	96.93	0.00	30.63	7.78	----	----	1.00	172.80	154.34	180.58	334.92
1982	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1983	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1984	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----
1985	12.26	0.00	24.32	0.00	93.50	5.34	58.98	1.70	0.00	0.00	2.83	64.95	191.89	71.99	263.88
1986	12.81	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1987	13.16	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1988	12.84	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1989	14.56	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1990	14.19	0.00	24.75	0.00	97.01	3.20	80.27	4.04	0.65	0.00	2.19	108.51	219.06	115.75	334.81
1991	13.37	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1992	13.66	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1993	13.75	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1994	13.47	0.00	----	----	----	----	----	----	----	----	----	----	N/A	N/A	N/A
1995	14.29	0.00	23.06	0.00	93.95	3.98	80.37	2.93	1.00	0.00	2.27	113.21	214.94	120.12	335.06
1996	14.48	0.00	----	----	87.22	3.82	----	----	----	----	2.56	97.60	N/A	N/A	N/A
1997	14.53	0.00	----	----	84.08	3.74	----	----	----	----	2.44	120.00	N/A	N/A	N/A
1998	15.51	0.00	----	----	79.73	4.08	----	----	----	----	2.41	116.89	N/A	N/A	N/A
1999	15.36	0.00	----	----	84.49	1.73	----	----	----	----	2.50	122.93	N/A	N/A	N/A
2000	15.79	0.00	20.06	0.00	81.85	4.92	93.87	1.87	1.06	0.37	2.54	101.01	215.17	108.17	323.34

Public supply treated includes water that does not meet public drinking water standards and is either treated through a desalination process or is diluted with fresher water to meet standards.

The procedure used to estimate domestic self-supplied withdrawals changed for 2000 from previous years as a different per capita was used to calculate withdrawals.

Agricultural includes water withdrawn for crop irrigation, livestock, and fish farming purposes.

Recreational includes water used for all turf grass (golf, commercial, industrial, and public) irrigation. This category was accounted for under Agricultural from 1975 through 1984.

Data sources: 1975-1990 - Modified from USGS Open-File Report 94-521 (Marella, 1995).

1991-1994, 1996-1999 - USGS unpublished water-use data files, Tallahassee.

1995 - USGS Water-Resources Investigations Report 99-4002 (Marella, 1999).

2000 - USGS Scientific Investigations Report 2004-5151 (Marella, 2004).

Glossary

A

Advanced wastewater treatment Any treatment of sewage that goes beyond the secondary or biological water treatment stage and includes the removal of nutrients, such as phosphorus and nitrogen and a high percentage of suspended solids. This treatment is more stringent than secondary treatment, and requires an 85 percent reduction in conventional pollutant concentrations or a significant reduction in nonconventional pollutants (U.S. Environmental Protection Agency, 1997, written commun.).

Agriculture water use Includes water used for agricultural irrigation and nonirrigation purposes. Irrigation water use includes the artificial application of water on lands to assist in the growing of crops, plants, and pasture. Nonirrigation water use includes water used for livestock, fish farming, and other farm needs. Livestock water use includes water used for stock watering, feedlots, and dairy operations. The water can be obtained from a public supply or be self-supplied.

C

Commercial water use Water for motels, hotels, restaurants, office buildings, commercial facilities and civilian and military institutions. The water may be obtained from a public supply or be self-supplied.

Community water system A public water system which serves at least 15 service connections used by year-round residents or regularly serves at least 25 year-round residents (Florida Department of Environmental Regulation, 1990).

Consumptive use That part of water withdrawn that is evaporated, transpired, incorporated into products or crops, consumed by humans or livestock, or otherwise removed from the immediate water environment. Sometimes called water consumed or water depleted.

Cooling pond Usually a manmade water body used by power plants or large industrial plants that enables the facility to recirculate

once-through cooling water. The water levels in the pond are usually maintained by rainfall or augmented by pumping (withdrawal of) water from another source (fresh, saline, or reclaimed).

Cooling tower A large tower or stack that is used for heat exchange of once-through cooling water generated by steam condensers. Hot water from the plant is sprayed into the top of the tower and exchanges heat with the passing air as it falls. The water is then collected at the bottom of the tower and used again. A small amount of water is lost (consumed) through evaporation in this process. See cooling water or once-through cooling.

Cooling water Water used for cooling purposes by electric generators, steam condensers, large machinery or products at power or industrial plants. Water used for cooling purposes can be either fresh, saline, or reclaimed and may be used only once or recirculated multiple times. See cooling pond or once-through cooling water.

D

Desalination The removal of salts from water. Desalination is primarily used for public-supply water to ensure that it meets Florida Department of Environmental Protection secondary drinking standards. The three primary types of desalination used in Florida are: (1) distillation, (2) electro dialysis processes, and (3) reverse osmosis processes (Buros, 1989, South Florida Water Management District, 1990). The reverse osmosis processes are the most commonly used in Florida followed by electro dialysis (Dykes and Conlon, 1989). In addition to these three desalination processes, many public suppliers also dilute or blend brackish or saline water with fresher water to produce potable water. Also see reverse osmosis.

Dewatering The deliberate attempt to lower the ground-water level in or below land surface for selected purposes such as agricultural, construction, mining or other activities. For mining operation, dewatering usually is accomplished by pumping the water out of the

ground and discharging to a surface-water body. However, some dewatering involves gravity feeding water from the surficial aquifer into a deeper aquifer (usually the Floridan aquifer system) through recharge wells (Campbell, 1986). In Florida, this discharge usually requires a permit from the Florida Department of Environmental Protection.

Domestic wastewater facility Facilities that receive or dispose of wastewater derived principally from residential dwellings, business or commercial buildings, institutions, and the like (Florida Department of Environmental Regulation, 1991). Can also include some wastewater derived from industrial facilities. May also be referred to as a municipal wastewater facility.

Domestic water use Water for normal household purposes, such as drinking, food preparation, bathing, washing clothes and dishes, flushing toilets, and watering lawns and gardens. Also called residential water use. The water can be obtained from a public supply or be self-supplied.

E

Effluent Water that flows out of a wastewater treatment facility or other works used for the purpose of treating, stabilizing, or holding waste.

F

Flood irrigation Irrigation systems that control the water table with lateral supply ditches. These include open field ditch systems (furrows), semi-closed conveyance systems, subsurface conduit systems, crown flood systems, and continuous flood systems. Also includes seepage or subsurface irrigation systems. The efficiencies of these flood irrigation systems range from 20 to 80 percent (Smajstrla and others, 1988), however, an average of 60 percent is commonly used for estimating water requirements. May also be referred to as subsurface irrigation.

Freshwater Water that contains less than 1,000 milligrams per liter (mg/L) of dissolved solids; generally, more than 500 mg/L is considered undesirable for drinking and many industrial uses. Generally, freshwater is considered potable.

G

Gigawatthour (GWh) A measure of electricity, one billion watthours.

Ground water Specifically, that part of the subsurface water that is in the saturated zone (a zone in which all voids are filled with water).

Ground-water disposal Wastewater that is disposed of through the ground either by injection or seepage. This includes the following discharge methods; absorption beds, injection wells, drainfields, percolation ponds, rapid infiltration basins, spray fields, and land application systems (Marella, 1994). Land application systems or reuse systems are considered a ground-water disposal as treated wastewater used to irrigate is generally intended to filter down through the soil.

H

Hydroelectric power water use The use of water in the generation of electricity at plants where the turbine generators are driven by falling water. This is considered an instream use of water.

I

Industrial wastewater facility Facilities that produce, treat or dispose of wastewater not otherwise defined as a domestic wastewater; includes the runoff and leachate from areas that receive pollutants associated with industrial or commercial storage, handling, or processing (Florida Department of Environmental Regulation, 1991).

Industrial water use Water used for industrial purposes such as fabricating, processing, washing, and cooling, and includes such industries as steel, chemical and allied products, paper and allied products, mining, and petroleum refining. The water can be obtained from a public supply or be self-supplied.

Instream use Water use taking place within a stream channel for such purposes as hydroelectric power generation, navigation, water-quality improvement, fish and wildlife propagation, and recreation. Sometimes called nonwithdrawal use or in-channel use.

M

Micro irrigation Irrigation systems that apply water directly to, or very near, the soil surface, either above the ground or into the air,

in discrete drops, continuous drops, small streams, mist, or sprays. These include drip systems, spray systems, jet systems, and bubbler systems. Also referred to as drip, low pressure or low volume irrigation. The efficiencies of these micro-irrigation systems range from 75 to 95 percent (Smajstrla and others, 1988), however, an average of 80 percent is commonly used for estimating water requirements.

Mining water use Water used for the extraction of minerals and liquids. Mining also includes water used for milling (such as crushing, screening, washing, and flotation), environmental purposes (such as dust control and wetland restoration or maintenance), material conveyance, dewatering, and domestic uses on the premises. Generally, most of the water used at a mining operation is self-supplied.

N

Navigational water use Water utilized as a means of commercial (and sometimes recreational) transportation. Includes water used to lift a vessel in a lock, or maintain a navigable channel level. Navigational water use is considered a nonconsumptive instream use of water and is generally not measured.

Net water use Water withdrawals plus or minus water transfers. In most counties, the net water use and water withdrawals are equal. However, in counties involved in water transfers (imports and exports), the net water use represents the actual amount of water used regardless of the amount of water withdrawn. In Florida, water transfers are mostly found in the public supply category. Also see water transfers.

Non-Community water system A public water system which provides piped water for human consumption to at least 15 service connections or which serves at least 25 individuals at least 60 days out of the year but which is not a community water system. The difference between a community water system and a non-community water system is that the former serves inhabitants whereas the latter serves transients or non-residents who otherwise do not inhabit the building served by the system. (Florida Department of Environmental Regulation, 1990).

Non-Transient Non-Community water

system A public water system that is not a community water system and that regularly serves at least 25 of the same persons over a 6 month period. (Florida Department of Environmental Regulation, 1990).

O

Once-through cooling water Water (fresh or saline) that is withdrawn from a river, stream or other water body (manmade or natural), or a well, that is passed through a steam condenser one time, and then returned to the river or stream or other water body some distance from the intake (Hughes, 1975). Once-through cooling water is used to exchange the heat from the steam condensers to the cooler water. This method of cooling is commonly used in power production throughout Florida, and usually results in no consumption.

Offstream use Water withdrawn or diverted from a ground- or surface-water source for public-water supply, industry, irrigation, livestock, thermoelectric power generation, and other uses. Sometimes called off-channel use or withdrawal use.

Other water use Water used in Florida for such purposes as heating, cooling, irrigation (public-supplied only), lake augmentation, and other nonspecific uses. The water can be obtained from a public supply or be self-supplied.

P

Per capita use The average amount of water used per person during a standard time period, generally per day.

Potable water Water that meets the quality standards set by the Florida Department of Environmental Protection. Potable water is considered safe for human consumption and is often referred to as drinking water. In Florida, chloride and dissolved-solids concentrations in potable water must be less than or equal to 250 and 500 mg/L, respectively. Freshwater that exceeds these chloride and dissolved solids limits is often referred to as slightly saline, brackish, or nonpotable water and is either diluted with fresher water or treated through a desalination process to meet potable-water standards for public supply.

Power generation water use Water used in the process of the generation of electric power through a thermoelectric or hydroelectric facility. The majority of water used for this category is for cooling purposes (much of which is used for once-through cooling). Water is also used for boiler makeup or domestic purposes throughout the plant. Boiler makeup water and water used for domestic purposes are generally obtained from public supply, however, for plants located in remote areas, this water can be self-supplied. Cooling water is generally self-supplied, although some smaller plants use public-supply water for cooling purposes.

Primary wastewater treatment First step in wastewater treatment where screens and sedimentation tanks are used to remove most materials that float or settle. Primary treatment removes about 30 percent of carbonaceous biochemical oxygen demand from domestic sewage (U.S. Environmental Protection Agency, 1997, written commun.).

Public supply Water withdrawn by public or private water suppliers and delivered to users who do not supply their own water. Water suppliers provide water for a variety of uses, such as domestic, commercial, industrial, thermoelectric power (domestic and cooling purposes), and public-water use. According to the Florida Department of Environmental Protection, any water system that serves more than 25 people or has 15 year-round service connections is considered a community public supplier (Florida Department of Environmental Regulation, 1990a). For this report, public supply includes those systems that serve more than 400 people or use more than 10,000 gallons per day.

Public-water use Water supplied from a public-water supply and used for such purposes as firefighting, street washing, and municipal parks and swimming pools. Public-water use also includes system water losses (water lost to leakage) and unusable water discharged from desalination or lime-softening facilities. Also referred to as utility-water use.

R

Reclaimed water Water that has received at least secondary treatment and is reused after leaving a wastewater treatment facility.

Recycled water Water that is used more than one time before it passes back into the natural hydrological system or is discharged into a wastewater system. Also referred to as recirculated water.

Resident population The number of persons who live in a State who consider the State their permanent place of residence. College students, military personnel, and inmates of penal institutions are counted as permanent residents. According to this definition, tourist and seasonal or part-time residents are considered nonresident population.

Residential water use See domestic water use.

Reuse system The deliberate application of reclaimed water for a beneficial or other useful purpose. Reuse may encompass landscape irrigation (such as golf courses, cemeteries, highway medians, parks, playgrounds, school yards, nurseries, and residential properties), agricultural irrigation (such as food and fruit crops, wholesale nurseries, sod farms and pasture grass), aesthetic uses, ground-water recharge, environmental enhancement of surface water and wetland restoration, fire protection, and other useful purposes.

Reverse osmosis The process of removing salts from water using a membrane. With reverse osmosis, the product water passes through a fine membrane that the salts are unable to pass through, and the salt waste (brine) is removed and disposed. This differs from electrodialysis where the salts are extracted from the feedwater by using a membrane with an electrical current to separate the ions. During electrodialysis the positive ions flow through one membrane, while the negative ions flow through a different membrane, leaving freshwater as the end product. In this report, reverse osmosis includes any water treated through both reverse osmosis and electrodialysis and any water diluted or blended with fresher water that was used to obtain potable water. Also see desalination.

S

Saline water Water that contains more than 1,000 mg/L of dissolved solids.

Secondary wastewater treatment The second step in most domestic wastewater treatment systems in which bacteria consume the organic parts of the waste. This treatment removes floating and settleable solids and about 90 percent of the oxygen-demanding substances and suspended solids. Disinfection is the final stage of secondary treatment (U.S. Environmental Protection Agency, 1997, written commun.).

Self-supplied water Water withdrawn from a ground- or surface-water source by a user and not obtained from a public supply.

Septic tank Refers to a buried watertight receptacle constructed to promote the separation of solids, grease, and liquid components of wastewater in the absence of oxygen. The liquid fraction from the septic tank is discharged to a drain field for further treatment or disposal. In Florida, septic tanks are regulated by the Florida Department of Health.

Settling pond A holding pond for wastewater where heavier particles sink to the bottom for removal and disposal.

Sprinkler irrigation A pressurized irrigation system where water is distributed through pipes to the field and applied through a variety of sprinkler heads or nozzles. Pressure is used to spread water droplets above the crop canopy to simulate a rainfall (Izuno and Haman, 1987). These systems include portable and traveling guns, solid or permanent fixtures (overhead or pop ups), center pivots, and periodic moving systems. Also referred to as overhead irrigation. The efficiencies of these sprinkler irrigation systems range from 15 to 85 percent (Sma-jstrla and others, 1988), however, an average of 70 percent is commonly used for estimating water requirements.

Surface-water disposal Refers to the release of reclaimed water or treated effluent directly into a surface water body (including marshes or wetlands). This does not include water discharged into ponds for holding or percolation purposes (Marella, 1994).

T

Tail-water runoff Unused irrigation water or rain water that is collected at the base or end of an irrigated system or field in a ditch or impoundment. This water may be reused again for irrigation purposes, be left to evaporate, percolate into the ground, receive treatment, and (or) be discharged to surface-water bodies.

Thermoelectric power Electrical power generated by using fossil fuel (coal, oil, natural gas or biomass), geothermal, or nuclear energy.

Treated (wastewater) effluent Water that has received primary, secondary, or advanced treatment and is released from a wastewater facility after treatment.

W

Wastewater A combination of liquid and water-carried pollutants from residential or commercial buildings, industrial plants, and institutions. Wastewater may include any ground water, surface runoff, or leachate that may be present in the system.

Water transfer Artificial conveyance of water from one area to another across a political or hydrological boundary. This is referred to as an import or export of water from one basin or county to another.

Water use 1) In a restrictive sense, the term refers to water that is actually used for a specific purpose such as domestic use, irrigation, or industrial processing. 2) More broadly, water use pertains to human's interaction with and influence on the hydrologic cycle, and includes elements such as water withdrawals, deliveries, consumptive use, wastewater releases, reclaimed wastewater, return flow and instream use.

Withdrawal Water removed from the ground or diverted from a surface-water source. The amount of water withdrawn may not equal the amount of water used due to water transfers or the recirculation or recycling of the same water. For example, a power plant may use the same water multiple times but withdraw a significantly different amount.



**125** *years of*
science
for America

1879–2004