

Figure A2-1. Timeseries plots of nondetections of acetochlor for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-2. Timeseries plots of nondetections of alachlor for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-3. Timeseries plots of nondetections of atrazine for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-4. Timeseries plots of nondetections of azinphos-methyl for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-5. Timeseries plots of nondetections of benfluralin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-6. Timeseries plots of nondetections of butylate for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-7. Timeseries plots of nondetections of carbaryl for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-8. Timeseries plots of nondetections of carbofuran for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-9. Timeseries plots of nondetections of chlorpyrifos for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-10. Timeseries plots of nondetections of cyanazine for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-11. Timeseries plots of nondetections of dacthal for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-12. Timeseries plots of nondetections of p,p'-DDE for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-13. Timeseries plots of nondetections of deethylatrazine for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-14. Timeseries plots of nondetections of desulfinylfipronil for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-15. Timeseries plots of nondetections of desulfinylfipronil amide for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-16. Timeseries plots of nondetections of diazinon for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-17. Timeseries plots of nondetections of dieldrin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-18. Timeseries plots of nondetections of 2,6-diethylaniline for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-19. Timeseries plots of nondetections of disulfoton for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-20. Timeseries plots of nondetections of EPTC for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-21. Timeseries plots of nondetections of ethalfluralin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-22. Timeseries plots of nondetections of ethoprophos for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-23. Timeseries plots of nondetections of fipronil for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-24. Timeseries plots of nondetections of fipronil sulfide for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-25. Timeseries plots of nondetections of fipronil sulfone for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-26. Timeseries plots of nondetections of fonofos for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-27. Timeseries plots of nondetections of alpha-HCH for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-28. Timeseries plots of nondetections of gamma-HCH for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-29. Timeseries plots of nondetections of linuron for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-30. Timeseries plots of nondetections of malathion for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-31. Timeseries plots of nondetections of metolachlor for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-32. Timeseries plots of nondetections of metribuzin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-33. Timeseries plots of nondetections of molinate for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-34. Timeseries plots of nondetections of napropamide for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-35. Timeseries plots of nondetections of parathion for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-36. Timeseries plots of nondetections of parathion-methyl for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-37. Timeseries plots of nondetections of pebulate for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-38. Timeseries plots of nondetections of pendimethalin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-39. Timeseries plots of nondetections of cis-permethrin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-40. Timeseries plots of nondetections of phorate for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-41. Timeseries plots of nondetections of prometon for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-42. Timeseries plots of nondetections of propachlor for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-43. Timeseries plots of nondetections of propanil for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-44. Timeseries plots of nondetections of propargite for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-45. Timeseries plots of nondetections of propyzamide for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-46. Timeseries plots of nondetections of simazine for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-47. Timeseries plots of nondetections of tebuthiuron for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-48. Timeseries plots of nondetections of terbacil for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-49. Timeseries plots of nondetections of terbufos for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-50. Timeseries plots of nondetections of thiobencarb for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-51. Timeseries plots of nondetections of triallate for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.




Figure A2-52. Timeseries plots of nondetections of trifluralin for all sites in the trend data set showing (1) original reporting levels; (2) rounded reporting levels and, for routine nondetections, reporting levels reassigned to the maximum value of the long-term method detection level (maxLT-MDL); and (3) raised reporting levels adjusted for temporal changes in recovery. Temporal changes in recovery are shown in panel 2 of the figures in appendix 3.