Appendix 3. Model Archive Summary for Total Suspended Solids at U.S. Geological Survey Site 07144780, North Fork Ninnescah River above Cheney Reservoir, Kansas, during January 1, 1999, through December 31, 2019

This model archive summary summarizes the total suspended solids (TSS) model developed to compute hourly or daily TSS during January 1, 1999, through December 31, 2019. This model is used concomitantly with other models during this period to calculate concentrations when other explanatory variables are not available for the purposes of load and concentration calculations. The methods used follow U.S. Geological Survey (USGS) guidance as referenced in relevant Office of Surface Water/Office of Water Quality Technical Memoranda and USGS Techniques and Methods, book 3, chapter C4 (Rasmussen and others, 2009), and other standard USGS methods (Sauer and Turnipseed, 2010; Turnipseed and Sauer, 2010).

Site and Model Information

Site number: 07144780

Site name: North Fork Ninnescah River above Cheney Reservoir, Kansas

Location: Lat 37°51'45", long 98°00'49" referenced to North American Datum of 1927, in NE 1/4 SE 1/4 NE 1/4 sec.19, T.25 S., R.6 W., Reno County, Kans., Hydrologic Unit 11030014, on right bank at upstream side of county highway bridge, 10 miles south of Hutchinson, 18.1 miles upstream from Cheney Dam.

Equipment: A Sutron Satlink 2 High Data Rate Collection Platform and a Design Analysis Water Log H350/355 nonsubmersible pressure transducer transfers real-time stage, precipitation, and water quality data via satellite. The primary reference gage is a Type-A wire-weight gage located on the downstream bridge guardrail. Check-bar elevation is 21.804 feet. The orifice is enclosed in 1 1/4-inch pipe, which runs from the gage house, under the bridge, and along an I-beam where it is attached to the concrete pier closest to the left edge of water.

Date model was developed: April 26, 2019

Model calibration data period: January 26, 1999, to September 28, 2017

Model Data

All data were collected using USGS protocols (U.S. Geological Survey, 2006; Wagner and others, 2006; Sauer and Turnipseed, 2010; Turnipseed and Sauer, 2010) and are stored in the National Water Information System (NWIS) database (https://doi.org/10.5066/F7P55KJN; U.S. Geological Survey, 2020). Explanatory variables were evaluated individually and in combination. Potential explanatory variables included streamflow, water temperature, specific conductance, pH, dissolved oxygen, and turbidity. Seasonal components (sine and cosine variables) were also evaluated as explanatory variables.

The regression model is based on 101 concomitant values of discretely collected TSS samples and continuously measured streamflow during January 26, 1999, through September 28, 2017. Discrete samples were collected over a range of streamflows. No samples were less than laboratory detection limits. Summary statistics and the complete model-calibration data are provided below. Outliers were identified using studentized residuals (for values greater than 3 or less than -3). None of the samples in this dataset were deemed outliers or removed from the model calibration dataset. Other data deemed outliers and removed in previously published versions of this model (Christensen and others, 2006; Stone and others, 2013) were examined and retained in the dataset if there were no clear issues, explanations, or conditions that would cause a result to be invalid for model calibration.

Total Suspended Solids

Discrete samples were collected from the downstream side of the bridge or instream within 50 feet of the bridge using equal-width-increment, multiple vertical, single vertical, or grab methods following U.S. Geological Survey (2006) and Rasmussen and others (2014). Discrete samples were collected on a semifixed to event-based schedule ranging from 1 to 17 samples per year with a Federal Interagency Sedimentation Project U.S. DH–95 or D–95 with a Teflon bottle, cap, and nozzle depth-integrating sampler; a DH–81 with a Teflon bottle, cap, and nozzle hand sampler; or a grab sample with a Teflon bottle depending on sample location. Samples were analyzed for TSS by

the Wichita Municipal Water and Wastewater Laboratory in Wichita, Kans., according to standard methods (American Public Health Association and others, 1995).

Continuous Data

Streamflow was measured using a nonsubmersible pressure transducer during January 1, 1999, through December 31, 2019. The continuous streamflow data used were time interpolated values from the continuous time series. If the continuous data were not available, the sample was not included in the dataset.

Model Development

Ordinary least squares regression analysis was done using R programming language (R Core Team, 2019) to relate discretely collected TSS to streamflow and other continuously measured data. The distribution of residuals was examined for normality and plots of residuals (the difference between the measured and model calculated values) compared to model calculated TSS were examined for homoscedasticity (departures from zero did not change substantially over the range of model calculated values). Previously published explanatory variables were also strongly considered for continuity; however, the best explanatory variable(s) was ultimately selected.

Streamflow was selected as a good predictor of logarithm base 10 (\log_{10}) (TSS) based on residual plots, relatively high coefficient of determination (R^2), and relatively low model standard percentage error (MSPE). This model was developed with the sole purpose to fill in gaps of missing data of the primary model for concentration and load estimations.

Model Summary

Summary of final TSS regression analysis at USGS site 07144780:

TSS-based model:

$$\log_{10}(TSS) = 0.464 \times \log_{10}(Q) + 0.877,$$

where.

TSS = total suspended solids, in milligrams per liter, and O = streamflow, in cubic feet per second.

The log-transformed model may be retransformed to original units so that TSS can be calculated directly. The retransformation introduces a bias in the calculated constituent. This bias may be corrected using Duan's bias correction factor (BCF; Duan, 1983). For this model, the calculated BCF is 1.35. The retransformed model, accounting for BCF, is as follows:

$$TSS = (Q^{0.464} \times 10^{0.877}) \times 1.35$$

Previous Models

Version	Model Equation	Reference
1.0	$\log_{10}(TSS) = 0.682 \times \log_{10}(Q) + 0.431$	Christensen and others (2006)
1.1	$\log_{10}(TSS) = 0.575 \times \log_{10}(Q) + 0.624$	Stone and others (2013)

Model Statistics, Data, and Plots

Definitions for terms used in this output can be found at the end of this document.


Model

 $\log TSS = +0.464 * \log Q + 0.877$


Variable Summary Statistics

	logTSS	TSS	logQ	Q
Minimum	0.924	8.4	0.724	5.29
1st Quartile	1.600	40.0	1.890	77.80
Median	2.050	112.0	2.310	205.00
Mean	1.960	143.0	2.320	643.00
3d Quartile	2.320	211.0	2.700	500.00
Maximum	2.840	696.0	3.910	8,220.00

Box Plots



Exploratory Plots



Red line shows the locally weighted scatterplot smoothing (LOWESS).

Basic Model Statistics

For a detailed definition and explanation of the terms used below, refer to Helsel and Hirsch (2002).

```
Number of Observations

Standard error (RMSE)

Average Model standard percentage error (MSPE)

Coefficient of determination (R<sup>2</sup>)

Adjusted Coefficient of Determination (Adj. R<sup>2</sup>)

Bias Correction Factor (BCF)

101

87.3

0.442

Adjusted Toefficient of Determination (Adj. R<sup>2</sup>)

1.35
```

Explanatory Variables

	Coefficients	Standard Error	t value F	Pr(> t)
(Intercept)	0.877	0.1260	6.94 4	4.21e-10
logQ	0.464	0.0524	8.86	3.35e-14

Correlation Matrix

	Intercept	E.vars
Intercept	1.000	-0.963
E.vars	-0.963	1.000


Outlier Test Criteria


Leverage C	Cook's D	DFFITS
0.0594	0.1944	0.2814


Flagged Observations

	logTSS	Estimate	Residual	Standard	Residual	Studentized	Residual	Leverage	Cook's D	DFFITS	
9/28/2000 10:30	0.924	1.48	-0.5510		-1.64		-1.650	0.0349	0.0485	-0.3140	
6/14/2010 11:30	1.960	2.62	-0.6540		-1.97		-1.990	0.0575	0.1180	-0.4930	
7/6/2010 10:30	2.110	2.69	-0.5800		-1.75		-1.770	0.0691	0.1140	-0.4830	


Statistical Plots







Cross Validation



Fold - equal partition of the data (10 percent of the data)


Large symbols - observed value of a data point removed in a fold

Small symbols - recomputed value of a data point removed in a fold

Recomputed regression lines - adjusted regression line with one fold removed

Minimum MSE of folds: 0.0192
Mean MSE of folds: 0.1200
Median MSE of folds: 0.1350
Maximum MSE of folds: 0.1720

(Mean MSE of folds) / (Model MSE): 1.0200



Red line - Model MSE

Blue line - Mean MSE of folds

Model-Calibration Data Set

	Date ⁻	logTSS	logQ	TSS	0	Computed	Computed	Residual	Normal	Censored
0	5466	106.33	-086	.55	٧	logTSS	TSS	csiadai	Quantiles	Values
	1999-01-26	1.23	2.02	16.8	104	1.81	87.5	-0.588	-1.69	
2	1999-01-31	2.37	2.78	232	602	2.17	198	0.199	0.575	
3	1999-02-03	1.8	2.38	62.4	241	1.98	129	-0.187	-0.575	
4	1999-03-17	2.21	2.58	162	384	2.08	161	0.133	0.354	
5	1999-04-06	2.29	2.63	194	428	2.1	169	0.189	0.518	
6	1999-04-16	2.49	2.84	309	697	2.2	212	0.294	0.901	
7	1999-05-13	1.74	1.99	54.8	97.7	1.8	85.1	-0.0618	-0.149	
8	1999-05-24	2.08	2.2	119	158	1.9	106	0.178	0.489	
9	1999-06-10	1.78	1.92	60.7	84	1.77	79.3	0.0132	0.0744	
10	1999-06-25	2.17	2.38	148	237	1.98	128	0.191	0.546	
11	1999-07-02	2.43	2.39	272	243	1.98	130	0.45	1.2	
12	1999-07-14	1.62	1.79	42	61.3	1.71	68.5	-0.0834	-0.25	
13	1999-07-29	1.68	1.78	48	60	1.7	67.8	-0.021	0	
14	1999-08-12	1.55	1.63	35.4	43	1.64	58.1	-0.086	-0.276	
	1999-08-26	1.32	1.52		33.2	1.58	51.5	-0.26	-0.864	
16	1999-09-22	1.48		30.5	49	1.66	61.7	-0.177	-0.489	
17	1999-12-02	1.4	1.79	25.2	61.2	1.71	68.5	-0.305	-0.901	
	2000-02-25	2.63	2.41	422	259	2	134	0.628	1.8	
19	2000-03-24	2.47	3.58	292	3790	2.54	464	-0.0723	-0.225	
_	2000-04-27	1.44		27.3	110	1.82	89.8	-0.388	-1.15	
	2000-05-25	1.59	1.78		60	1.7	67.8	-0.117	-0.302	
	2000-06-21	1.74	1.67		47	1.65	60.5	0.0897	0.25	
	2000-07-26	2.15	2.07	142	117	1.84	92.5	0.315	0.978	
	2000-08-29		0.968		9.3	1.33	28.5	-0.156	-0.434	
25	2000-09-28	0.924	1.29	8.4	19.5	1.48	40.3	-0.551	-1.51	

26 2000-10-26	2.21	3.43		2690	2.47	396	-0.259	-0.829	
27 2001-06-06	2.33	3.26		1820	2.39	331	-0.0642	-0.174	
28 2001-09-04	1.66	1.26		18	1.46	38.8	0.199	0.605	
29 2001-09-19	2.55	2.27	357	187	1.93	115	0.621	1.69	
30 2002-06-12	2.76	2.65	582	446	2.11	172	0.658	2.15	
31 2002-08-14	2.42	2.54	264	346	2.06	153	0.366	1.11	
32 2003-03-18	2.84	2.61	696	412	2.09	166	0.752	2.52	
33 2003-04-21	2.09	2.74	124	551	2.15	190	-0.0558	-0.124	
34 2004-03-05	2.57	3.29		1950	2.4	341	0.17	0.462	
35 2004-05-14	2	2.76	100	576	2.16	194	-0.158	-0.462	
36 2004-06-14	2.21	2.38	162	238	1.98	129	0.23	0.696	
37 2004-06-14	2.2	2.38	159	238	1.98	129	0.221	0.665	
38 2004-09-08	1.11	1.59 1		39	1.62	55.5	-0.508	-1.37	
39 2005-03-24	2.08	2.58	120	377	2.07	159	0.00664	0.0248	
40 2005-05-16	2.11	2.72	128	526	2.14	186	-0.0326	-0.0744	
41 2005-06-10	2.4	3.05		1120	2.29	263	0.106	0.276	
42 2005-06-13	2.16	3.5		3150	2.5	426	-0.339	-0.978	
43 2005-08-29	2.24	2.41	172	255	1.99	133	0.242	0.761	
44 2006-03-02	1.33	1.97 2			1.79	83	-0.463	-1.25	
45 2006-03-22	1.94	2.16 8		144	1.88	102	0.0618	0.124	
46 2006-05-01	1.74	1.93 5			1.77	79.6	-0.0299	-0.0248	
47 2006-05-12	2.06	2.19	114	155	1.89	106	0.163	0.434	
48 2006-06-05	1.88	1.59	75	39	1.62	55.5	0.26	0.829	
49 2006-07-31	1.38	1.15		14.2	1.41	34.8	-0.0322	-0.0496	
50 2006-09-07	1.46	1.59	29	39	1.62	55.5	-0.153	-0.407	
51 2006-09-21	1.18	1.36	15	23	1.51	43.5	-0.333	-0.939	
52 2006-09-21	1.15	1.36	14	23	1.51	43.5	-0.363	-1.06	
53 2007-01-09	1.48	1.85	30	70	1.73	72.8	-0.256	-0.795	
54 2007-03-14	1.36	1.82		66.7	1.72	71.2	-0.362	-1.02	
55 2007-03-22	1.87	1.99	74	97 206	1.8	84.8	0.0702	0.174	
56 2007-03-26	2.5	2.47	316	296	2.02	142	0.476	1.25	
57 2007-03-31 58 2007-04-16	2.66 1.97	3.12 2.87	94	1320 744	2.33	285 218	0.336	1.02	
59 2007-05-07	2.32	3.57		3720	2.21 2.53	461	-0.236 -0.21	-0.728 -0.635	
60 2007-06-29	1.96	2.6	92	401	2.33	461 164	-0.21 -0.121	-0.033 -0.328	
61 2007-09-04	1.35	1.45 2		28	1.55	47.6	-0.121 -0.198	-0.528 -0.605	
62 2008-04-24	2.13	2.43	136	267	2	136	0.131	0.328	
63 2008-05-09	2.45	3.49		3110	2.5	424	-0.0463	-0.0992	
64 2008-06-19	1.82	2.35		223	1.97	125	-0.0403	-0.381	
65 2008-09-15	2.16	2.59	146	390	2.08	162	0.0851	0.225	
66 2008-10-16	2.05	2.93	112	848	2.24	232	-0.187	-0.546	
67 2009-03-31	2.56	2.85	364	713	2.2	214	0.36	1.06	
68 2009-04-27	2.5	3.34		2180	2.43	359	0.0761	0.199	
69 2009-06-17	2.3	2.54	201	349	2.06	154	0.246	0.795	
70 2009-08-20	2.44	2.01	278	103	1.81	87.2	0.633	1.95	
71 2009-09-10	2.41	2.68	258	481	2.12	178	0.29	0.864	
72 2010-04-23	2.29	2.16	197	144	1.88	102	0.415	1.15	
73 2010-04-23	2.5	2.10	318	157	1.9	106	0.606	1.6	
74 2010 05 17	2.58	2.53	381	341	2.05	152	0.528	1.37	
75 2010-05-27	1.96	3.75		5630	2.62	558	-0.654	-1.8	
76 2010-06-16	2.06	3.36		2280	2.44	367	-0.374	-1.11	
77 2010-07-06	2.11	3.91		8220	2.69	665	-0.58	-1.6	
78 2010-08-25	2.31	3.05		1110	2.29	263	0.0185	0.0992	
79 2011-04-13	1.3	1.9		80.2	1.76	77.6	-0.46	-1.2	
80 2011-05-23	1.95	1.96		92.2	1.79	82.8	0.16	0.407	
,		•			=		33	- · · • ·	

81	2011-06-28	1.26	1.29	18	19.7	1.48	40.4	-0.222	-0.665	
82	2011-07-27	1.28	0.724	19	5.29	1.21	22	0.0661	0.149	
83	2012-02-06	1.82	2.31	66	205	1.95	120	-0.131	-0.354	
84	2012-03-23	2.21	2.53	161	338	2.05	151	0.156	0.381	
85	2012-06-20	1.46	1.41	29	25.9	1.53	45.9	-0.0707	-0.199	
86	2012-08-27	1.45	1.76	28	57.8	1.69	66.7	-0.248	-0.761	
87	2013-04-11	1	1.89	10	77.8	1.75	76.5	-0.755	-2.52	
88	2013-05-10	1.2	2.26	16	181	1.92	113	-0.721	-2.15	
89	2013-05-31	1.82	3.03	66	1080	2.28	259	-0.465	-1.31	
90	2013-08-05	1.83	3.48	67	3010	2.49	417	-0.665	-1.95	
91	2013-08-16	1.6	2.7	40	500	2.13	181	-0.528	-1.44	
92	2013-10-31	2.08	2.15	119	142	1.88	101	0.2	0.635	
93	2014-05-13	2.25	2.27	176	187	1.93	115	0.314	0.939	
94	2014-06-10	2.48	2.2	304	158	1.9	106	0.586	1.51	
95	2014-07-02	1.93	2.78	86	597	2.17	197	-0.231	-0.696	
96	2015-04-08	1.7	1.75	50	56.8	1.69	66.1	0.00798	0.0496	
97	2015-04-14	2.34	2	218	99.7	1.8	85.8	0.534	1.44	
98	2016-04-19	2	2.14	100	138	1.87	99.7	0.131	0.302	
99	2017-04-20	2.35	2.67	224	463	2.11	175	0.236	0.728	
100	2017-08-11	2.32	2.07	211	116	1.84	92.2	0.489	1.31	
101	2017-09-28	1.94	2.69	88	485	2.12	179	-0.179	-0.518	

Definitions

TSS: total suspended solids, in milligrams per liter (00530)
Q: streamflow, instantaneous, in cubic feet per second (00061)

Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References Cited

American Public Health Association, American Water Works Association, and Water Environment Federation, 1995, Standard methods for the examination of water and wastewater (19th ed.): Washington, D.C., American Public Health Association, 905 p.

Christensen, V.G., Graham, J.L., Milligan, C.R., Pope, L.M., and Ziegler, A.C., 2006, Water quality and relation to taste-and-odor compounds in the North Fork Ninnescah River and Cheney Reservoir, south-central Kansas, 1997–2003: U.S. Geological Survey Scientific Investigations Report 2006–5095, 43 p. [Also available at https://doi.org/10.3133/sir20065095.]

Duan, N., 1983, Smearing estimate—A nonparametric retransformation method: Journal of the American Statistical Association, v. 78, no. 383, p. 605–610. [Also available at https://doi.org/10.1080/01621459.1983.10478017.]

- Helsel, D.R., and Hirsch, R.M., 2002, Statistical methods in water resources: U.S. Geological Survey Techniques of Water-Resources Investigations, book 4, chap. A3, 522 p. [Also available at https://doi.org/10.3133/tm4A3.]
- R Core Team, 2019, R—A language and environment for statistical computing: Vienna, Austria, R Foundation for Statistical Computing, accessed August 2019 at https://www.R-project.org/.
- Rasmussen, T.J., Bennett, T.J., Stone, M.L., Foster, G.M., Graham, J.L., and Putnam, J.E., 2014, Quality-assurance and data-management plan for water-quality activities in the Kansas Water Science Center, 2014: U.S. Geological Survey Open-File Report 2014–1233, 41 p., accessed April 2020 at https://doi.org/10.3133/ofr20141233.
- Rasmussen, P.P., Gray, J.R., Glysson, G.D., and Ziegler, A.C., 2009, Guidelines and procedures for computing time-series suspended-sediment concentrations and loads from in-stream turbidity-sensor and streamflow data: U.S. Geological Survey Techniques and Methods, book 3, chap. C4, 52 p. [Also available at https://doi.org/10.3133/tm3C4.]
- Sauer, V.B., and Turnipseed, D.P., 2010, Stage measurement at gaging stations: U.S. Geological Survey Techniques and Methods, book 3, chap. A7, 45 p., accessed April 2020 at https://doi.org/10.3133/tm3A7.
- Stone, M.L., Graham, J.L., and Gatotho, J.W., 2013, Continuous real-time water-quality monitoring and regression analysis to compute constituent concentrations and loads in the North Fork Ninnescah River upstream from Cheney Reservoir, south-central Kansas, 1999–2012: U.S. Geological Survey Scientific Investigations Report 2013–5071, 44 p., accessed July 2020 at https://doi.org/10.3133/sir20135071.
- Turnipseed, D.P., and Sauer, V.B., 2010, Discharge measurements at gaging stations: U.S. Geological Survey Techniques and Methods, book 3, chap. A8, 87 p., accessed April 2020 at https://doi.org/10.3133/tm3A8.
- U.S. Geological Survey, 2006, Collection of water samples (ver. 2.0, September 2006): U.S. Geological Survey Techniques of Water Resources Investigations, book 9, chap. A4 [variously paged]. [Also available at https://doi.org/10.3133/twri09A4.]
- U.S. Geological Survey, 2020, USGS water data for the Nation: U.S. Geological Survey National Water Information System database, accessed April 20, 2020, at https://doi.org/10.5066/F7P55KJN.

Wagner, R.J., Boulger, R.W., Jr., Oblinger, C.J., and Smith, B.A., 2006, Guidelines and standard procedures for continuous water-quality monitors—Station operation, record computation, and data reporting: U.S. Geological Survey Techniques and Methods, book 1, chap D3, 51 p. plus 8 attachments. [Also available at https://doi.org/10.3133/tm1D3.]