

Mr. Lemmon
*(200)
T67K
70.292

Unclassified

Reconnaissance for Radioactive Deposits in Eastern Alaska, 1952

By A. E. Nelson, ^{Arthur Edward} W. S. West, ^{1922 - Walter Scott} and J. J. Matzko, ^{John Joseph} 1920-



Trace Elements Investigations Report 292

UNITED STATES DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY

TEI-292

K(200)
T67x
no. 292

Geology and Mineralogy

This document consists of 55 pages,
plus 2 figures.
Series A

UNITED STATES DEPARTMENT OF THE INTERIOR

✓ U.S. GEOLOGICAL SURVEY

RECONNAISSANCE FOR RADIOACTIVE DEPOSITS

IN EASTERN ALASKA, 1952*

By

Arthur E. Nelson, Walter S. West,
and John J. Matzko

April 1954

Trace Elements Investigations Report 292

This preliminary report is distributed
without editorial and technical review
for conformity with official standards
and nomenclature. It is not for public
inspection or quotation.

*This report concerns work done on behalf of the Division
of Raw Materials of the U. S. Atomic Energy Commission.

USGS-TEI-292

GEOLOGY AND MINERALOGY

Distribution (Series A)	No. of copies
American Cyanamid Company, Winchester	1
Argonne National Laboratory	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute, Columbus	1
Carbide and Carbon Chemicals Company, Y-12 Area	1
Division of Raw Materials, Albuquerque	1
Division of Raw Materials, Butte	1
Division of Raw Materials, Denver	1
Division of Raw Materials, Douglas	1
Division of Raw Materials, Hot Springs	1
Division of Raw Materials, Ishpeming	1
Division of Raw Materials, New York	6
Division of Raw Materials, Phoenix	1
Division of Raw Materials, Richfield	1
Division of Raw Materials, Salt Lake City	1
Division of Raw Materials, Washington	3
Dow Chemical Company, Pittsburg	1
Exploration Division, Grand Junction Operations Office	1
Grand Junction Operations Office	1
Technical Information Service, Oak Ridge	6
Tennessee Valley Authority, Wilson Dam	1
Terr. Dept. Mines, College (L. L. Patton)	1
Terr. Dept. Mines, Juneau (P. H. Holdsworth)	1
Terr. Dept. Mines, Ketchikan (A. E. Glover)	1
U. S. Geological Survey:	
Alaskan Geology Branch, Washington	8
Fuels Branch, Washington	1
Geochemistry and Petrology Branch, Washington	1
Geophysics Branch, Washington	1
Mineral Deposits Branch, Washington	1
K. L. Buck, Denver	1
R. M. Chapman, Fairbanks	1
R. P. Fischer, Grand Junction	1
G. O. Gates, Menlo Park	2
K. R. Klepper, Washington	1
D. M. Lemmon, Washington	1
L. H. Saarela, Anchorage	1
C. L. Sainsbury, Juneau	1
A. E. Weissenborn, Spokane	1
TEPCO, Denver	2
TEPCO, RPS, Washington	3
(Including master)	64

CONTENTS

	Page
Abstract	6
Chapter A.--Eastern Alaska Range, by A. E. Nelson	
Introduction	7
Geology	9
Areas investigated	10
Slana-Nabesna district	10
Mineral Point area	10
Slana area	11
Indian Group prospect	11
Silver Creek prospect	13
Rock Creek prospect	15
Nabesna mine	16
Orange Hill	18
Chisana district	19
Bonanza Creek	19
Conclusions	21
Literature cited	23
Unpublished reports	23
Chapter B.--Hope Creek and Miller House-Circle Hot Springs areas, by W. S. West and J. J. Matzko	
Introduction	24
Areas investigated	25
Hope Creek area	25
Miller House-Circle Hot Springs area	29
Conclusions	38
Literature cited	40
Chapter C.--Koyukuk-Chandalar region, by A. E. Nelson	
Introduction	41
Geology	42
Areas investigated	43
Chandalar mining district	43
Gold Bench	48
Conclusions	49
Literature cited	51
Unpublished reports	51
Chapter D.--Fortymile district, by J. J. Matzko	
Introduction	52
Areas investigated	52
Wade Creek	52
Walker Fork	54
Conclusions	55
Literature cited	55

ILLUSTRATIONS

	Page
Figure 1. Index map of eastern Alaska showing areas investigated in 1952	8
2. Geologic sketch map of the Chisana district showing sample locations	20
3. Geologic sketch map of the Hope Creek area showing sample locations	In envelope
4. Geologic sketch map of the Miller House-Circle Hot Springs area showing sample locations	In envelope
5. Sketch map of the Chandalar mining district showing sample locations	44
6. Topographic map of part of the Fortymile district, east-central Alaska	53

TABLES

Table 1. Radioactivity and mineralogy of rock and ore samples collected in the Slana-Nabesna district, 1952	12
2. Radioactivity and mineralogy of stream-gravel and placer concentrates collected in the eastern Alaska Range, 1952	14
3. Radioactivity data on rock and ore samples collected in the Chisana district	22
4. Equivalent-uranium analyses of samples, Hope Creek area, Alaska	27
5. Mineralogy of the heavy-mineral fraction of selected samples, Hope Creek area, Alaska	28
6. Equivalent-uranium analyses of samples, Miller House-Circle Hot Springs area, Alaska	32
7. Analyses of water samples, Miller House-Circle Hot Springs area, Alaska	34
8. Mineralogy of the heavy-mineral fractions of selected samples from the Miller House-Circle Hot Springs area, Alaska	36

TABLES--Continued

	Page
Table 9. Uranium-bearing minerals in samples from the Miller House-Circle Hot Springs area, Alaska	37
10. Radioactivity of stream and placer concentrate samples from the Koyukuk-Chandalar region, Alaska	46
11. Radioactivity of rock and ore samples from the Koyukuk-Chandalar region, Alaska	47

RECONNAISSANCE FOR RADIOACTIVE DEPOSITS

IN EASTERN ALASKA, 1952

By Arthur E. Nelson, Walter S. West, and John J. Matzko

ABSTRACT

Reconnaissance for radioactive deposits was conducted in selected areas of eastern Alaska during 1952. Examination of copper, silver, and molybdenum occurrences and of a reported nickel prospect in the Slana-Nabesna and Chisana districts in the eastern Alaska Range revealed a maximum radioactivity of about 0.003 percent equivalent uranium. No appreciable radioactivity anomalies were indicated by aerial and foot traverses in the area.

Reconnaissance for possible lode concentrations of uranium minerals in the vicinity of reported fluorite occurrences in the Hope Creek and Miller House-Circle Hot Springs areas of the Circle quadrangle and in the Fortymile district found a maximum of 0.055 percent equivalent uranium in a float fragment of ferruginous breccia in the Hope Creek area; analysis of samples obtained in the vicinity of the other fluorite occurrences showed a maximum of only 0.005 percent equivalent uranium.

No uraniumiferous lodes were discovered in the Koyukuk-Chandalar region, nor was the source of the monazite, previously reported in the placer concentrates from the Chandalar mining district, located. The source of the uranothorianite in the placers at Gold Bench on the South Fork of the Koyukuk River was not found during a brief reconnaissance, but a placer concentrate was obtained that contains 0.18 percent equivalent uranium. This concentrate is about ten times more radioactive than concentrates previously available from the area.

CHAPTER A.--EASTERN ALASKA RANGE

By Arthur E. Nelson

INTRODUCTION

An appraisal of uranium possibilities in Alaska made in 1950-51 (Wedow and others, 1951) indicated that certain mineral deposits in several localities in the eastern Alaska Range were favorable for the occurrence of uranium ores. Seven localities (nos. 1-7, fig. 1) in the eastern Alaska Range were investigated for possible radioactive deposits in the summer of 1952 on behalf of the Division of Raw Materials of the U. S. Atomic Energy Commission. The work was done by Arthur E. Nelson, geologist, and Richard S. Smith, geologic field assistant.

The investigations in the eastern Alaska Range consisted of radioactivity traverses of mines and prospects in and around mineralized zones, contacts, and aureoles. Specific outcrop tests were made at various points along the traverses. Supplemental check samples were collected for radioactivity analysis and mineralogic study. These included channel and wall-rock samples of veins and mineralized zones, and samples of ore dumps and various rock types. Concentrates were taken of certain stream gravels to determine if radioactive minerals were being released from the bedrock within the specific drainage basins. In addition, an airborne radioactivity traverse was flown over the Orange Hill area as well as over other larger mineralized zones in the region that were otherwise inaccessible with the equipment available.

The equipment used for detecting radioactivity in this work was standard commercial portable survey meters adapted to accept a variety of probes. Individual outcrop tests were made with a 6-inch beta-gamma probe; ground traverses were



FIGURE 1.--INDEX MAP OF EASTERN ALASKA SHOWING AREAS INVESTIGATED IN 1952

made with the survey meter and a 2- x 20-inch gamma probe lashed to a packboard; and airborne traverses were flown with a probe consisting of six 2- x 40-inch gamma tubes_s, connected in parallel_s, coupled to the survey meter and carried in light_s, fixed-wing aircraft. (See Wedow_s, 1951.)

GEOLOGY

The Slana-Nabesna and Chisana districts lie in that part of the eastern Alaska Range known as the Mentasta and Nutzotin Mountains (fig. 1). Here the Alaska Range has a southeasterly trend beginning at a point in the Mentasta Mountains known as Mentasta Pass and merging with the Nutzotin Mountains near the Nabesna River_s, which is a northeasterly flowing glacial stream originating in the Wrangell Mountains to the southwest. The Nutzotin Mountains continue the southeasterly trend of the Alaska Range toward the international boundary. On the northeast the range is bordered by the broad lowlands of the Tanana River and to the southwest by the Copper River and the Wrangell Mountains. The relief in this part of the Alaska Range is rugged_s and as a whole the mountains have been intensely glaciated in recent geologic time. In the higher altitudes there are numerous small glaciers.

The geology of the Nutzotin Mountains has been discussed in detail by Moffit (1943). In summary_s, a series of sedimentary beds_s, locally metamorphosed and ranging in age from Devonian to Quaternary_s, form a major part of the eastern Alaska Range. Shale_s, argillite_s, sandstone_s, arkose_s, graywacke_s, and conglomerate are the major bedrock types with minor amounts of limestone. These beds have been folded and locally faulted.

Igneous rock types are widely distributed throughout the region and occur as both flows and intrusives. The flow types are present throughout most of the section and in the main are represented by basalts and andesites. Intrusives of Paleozoic and Mesozoic age ranging from gabbro to granite intrude all the rock types with the intrusives of Mesozoic age being more extensively developed. Superimposed upon the consolidated sedimentary and igneous rocks are a series of unconsolidated stream and glacial deposits which occur in most of the valleys.

AREAS INVESTIGATED

Specific mines, properties, and mineralized zones in six localities (nos. 1-5 and 7, fig. 1) in the eastern Alaska Range were investigated for the occurrence of uranium ores, and one locality (no. 6, fig. 1) was traversed with airborne radiation equipment. The areas and specific properties investigated are discussed below.

Slana-Nabesna district

Mineral Point area

Thorne (1946, p. 8, 9) reported a nickel occurrence in a shear zone in argillite in the Mineral Point area (locality 1, fig. 1) located about 32 miles northeast of Slana on the eastern section of the Glenn Highway. The prospect is located on the north side of the road on a mountain that has two prominent limestone peaks which are separated by a saddle of reddish argillitic rock. Some of the rock in the shear zone has been altered to a red-yellow material.

Investigations conducted in the area revealed little mineralization. Radioactivity traverses were made over the entire area which included the shear zone as well as contacts and all rock types outcropping on the mountain. No anomalous radiation was detected during the course of the traverses. The radioactivity data on the rock specimens collected are shown in table 1.

Slana area

Slana, located near the confluence of the Slana and Copper Rivers, is on the south flank of the Alaska Range. Deposits of silver-bearing galena and copper minerals in quartz veins occur at several prospects located at different sites northwest of Slana. Two of these properties, the Indian Group and the Silver Creek prospects were examined for radioactive minerals. (See fig. 1.)

Indian Group prospect.--The Indian Group prospect (locality 2, fig. 1) is near the top of a ridge separating the Indian Creek and Ahtell Creek valleys and is 13 airline miles from the Indian Creek bridge on the eastern section of the Glenn Highway. The site of most of the open cuts is on the Indian Creek side of the ridge dividing the Indian Creek drainage from that of Ahtell Creek. A trail from Indian Creek leads up to the property, and the first of the open cuts was encountered about 400-500 feet below the crest of the divide. Little or no development work has been done on the Indian Group claims in recent years. At the present time there are at least six recognizable open cuts, all of which have either caved or have been partially filled with slide rock so that none of the mineralized portions of the veins are exposed. Sampling therefore was confined to the numerous ore dumps and float in and around the cuts.

Table 1.--Radioactivity^{1/} and mineralogy^{2/} of rock and ore samples collected in the Slana-Nabesna district, 1952

Sample number	Location	Type of sample	Equivalent uranium (percent)	Radioactive minerals
4510	Mineral Point (nickel prospect)	Rock occurring in shear zones	< 0.001	
4529	Indian Group, Indian Creek	Ore dump	.002	
4530	Indian Group, Indian Creek	Quartz diorite country rock	.004	Accessory minerals, mostly zircon
4532	Silver Creek (silver-lead prospect)	Ore dump	< .001	
4533	Silver Creek (silver-lead prospect)	Vein material	.001	
4534	Silver Creek (silver-lead prospect)	Wall rock next to vein	< .001	
4535	Rock Creek (molybdenite prospect)	Mafic phase of igneous country rock	.002	
4536	Rock Creek (molybdenite prospect)	Syenite country rock	.003	Accessory minerals, mostly sphene and zircon
4537	Rock Creek (molybdenite prospect)	Pegmatite vein	.003	Accessory minerals, mostly zircon and a trace of sphene
4538	Nabesna mine	Ore dump	< .001	
4539	Nabesna mine	Ore dump	< .001	
4540	Nabesna mine	Ore dump	< .001	

^{1/} Equivalent-uranium analyses by J. J. Matzko and others, U. S. Geological Survey laboratory, College, Alaska

^{2/} By A. E. Nelson

Moffit (1938) gives a detailed account of the geology of the Slana district. Briefly, the country rock at the Indian Group claims consists of quartz diorite which shows wide variations in texture. Most of the rock has a porphyritic texture with large phenocrysts of feldspar set in a coarse-granular groundmass. Locally a series of nearly vertical fracture planes that strike nearly east seem to have had some control in the placement of the mineralized quartz veins. The ore minerals are galena, tetrahedrite, and chalcopyrite, all of which are silver-bearing.

Radioactivity traverses were conducted over the area. These included checks in each of the open cuts as well as traverses along the veins and in the areas immediately adjacent to the prospect itself. No radioactivity anomalies were discovered. Supplemental check samples were collected from the country rock and ore dumps, and one concentrate (sample 4531, table 2) was collected from the sand and gravel in a stream draining the mineralized area. Results of the laboratory studies made on the samples collected during these investigations are shown in tables 1 and 2.

Silver Creek prospect.--The prospect on Silver Creek (locality 3, fig. 1) is located on the north side of the creek a little more than one mile upstream from its junction with Ahtell Creek. No work has been done recently on the prospect, but the trail to the property is in good condition and the site is easily accessible from the highway. The workings at the prospect site consist of two adits, several open cuts and prospect pits, all of which have caved, and an inclined shaft slightly more than 15 feet long. The site of the caved adit at creek level had been examined briefly by Wedow and Matzko in 1946 (Wedow, Killeen, and others, 1954, p. 16-18).

Table 2.--Radioactivity^{1/} and mineralogy^{2/} of stream-gravel and placer concentrates collected in the eastern Alaska Range, 1952

Sample number	Location	Type of sample	eU ^{3/} (percent)	eU (percent)		eU (percent) Magnetic fraction		Radioactive minerals
				BH ^{4/}	IH ^{5/}	1.6 mag.	1.6 non-mag.	
4515	Skookum Creek, Chisana district	placer	< 0.001	--	< 0.001	--	--	--
4522	Canyon Creek, Chisana district	stream-gravel	< .001	--	< .001	--	--	--
4523	Bonanza Creek, Chisana district	stream-gravel	.001	--	< .001	--	--	--
4524	Johnson Creek, Chisana district	stream-gravel	.002	< 0.001	--	--	--	--
4531	Unnamed creek near "Slana Lake", Slana-Nabesna district	stream-gravel	< .001	--	.003	0.01 (sphene)	0.09 (zircon)	Sphene and zircon

^{1/} Equivalent-uranium content of original concentrates determined by J. J. Matzko and others, U. S. Geological Survey laboratory, College, Alaska; of heavy-mineral and magnetic fractions by A. E. Nelson.

^{2/} By A. E. Nelson

^{3/} Equivalent uranium

^{4/} bromoform-heavy mineral fraction (specific gravity greater than 2.8)

^{5/} methylene iodide-heavy mineral fraction (specific gravity greater than 3.3)

The prospect is located in a northwest-trending fault zone about one hundred feet wide in the quartz diorite country rock. Bedrock in the fault zone has been altered to a relatively soft material, in places highly stained with iron oxide and with some copper carbonate stains. Prospecting has been carried out on several steeply dipping mineralized quartz veins which occur in the shear zone and have the same northwest trend. Although most of the workings were inaccessible, ore containing pyrite, galena, chalcopyrite, and a blue copper stain in quartz gangue was found on the lower dump.

No anomalous radiation was indicated in traverses made across the fault zone and adjacent areas, ore dumps, and along exposed portions of the veins. Check sampling was confined to ore dumps for the most part, but it was possible to take a channel sample across the face of the vein in the inclined shaft. This channel sample contained galena and chalcopyrite in a gangue of quartz. The results of laboratory studies on these samples are given in table 1.

Rock Creek prospect

The Rock Creek molybdenite prospect is located on the south flank of the Alaska Range (locality 4, fig. 1) about 3.5 miles north of mile 84.5 (measured from junction of the Richardson and Glenn Highways near Gakona--see fig. 1) on the Nabesna road. The prospect is near the head of the west fork of the east branch of Rock Creek at a height of 2,200 feet above the elevation of the road. No work has been done on the prospect in recent years, but former development work consists of an adit 170 feet long and one open cut which has since caved and in its present condition is 30 feet long.

A report by Van Alstine (1945) discusses the molybdenite deposits at Rock Creek. The bedrock at the head of Rock Creek is part of a mass of rock mapped by Moffit (1938) as undifferentiated granitic rocks of late Paleozoic and Mesozoic age. Permian lava flows are exposed in the middle portion of the creek. The prospect site is in a pink gneissic syenite which has interlamination of biotite schist. A pegmatite dike up to two feet wide intrudes the syenite and schist. The dike outcrops 150 feet north of the adit and is about 100 feet higher in elevation. The only metallic mineral noted was molybdenite. This mineral occurs in the pegmatite dike as tiny plates and blebs erratically distributed. The pegmatite dike was not found in the adit which was driven in the syenite to intersect it.

Radioactivity traverses in the area failed to indicate any significant anomalies. Laboratory studies on check samples collected from the prospect are listed in table 1.

Nabesna mine

The Nabesna Mining Corporation property (locality 5, fig. 1) is located in the northern part of the Wrangell Mountains of the eastern Alaska Range where the Wrangell Mountains and the Nutzotin Mountains merge. The property is accessible from Slana by a road which is about 54 miles long.

The Nabesna area has been prospected periodically since 1899. However, it was not until the fall of 1929 that a company was formed to mine gold at Nabesna. In 1931 a mill was in operation, and mining continued until about 1947 when most of the ore bodies had been exhausted. At present the several adits and shafts at the

main property are in such poor condition that access into the mine is impossible. At a newer development, about 1/2 mile north of the main mine, two adits have been driven into an ore body; one of these is caved, but the other is open for several hundred feet.

Wayland (Moffit, 1943) gives a detailed account of the geology of the Nabesna mine. Briefly, Triassic limestone, intruded by stocks and dikes of quartz diorite, overlies shale and basaltic lavas and is capped by Tertiary lavas. The limestone has been recrystallized in part with the development of tactite. Three types of ore bodies occur at the Nabesna mine, the most important being a series of discontinuous veins of auriferous pyrite and calcite with lesser amounts of chalcopyrite, sphalerite, and galena. These veins occur as replacements along fractures and contacts in the limestone. The other two types of ore deposits occur as bodies of magnetite and of pyrrhotite.

Minerals occurring at the Nabesna mine are pyrite, chalcopyrite, sphalerite, galena, pyrrhotite, arsenopyrite, stibnite, gold, and calcite with a suite of contact minerals such as andradite, vesuvianite, epidote, magnetite, specularite, wollastonite, spinel, brookite, and others.

Because entrance into the mine was impossible, investigations consisted of traverses along contacts on the property where possible and the ore dumps. At the newer development, 1/2 mile north of the main mine, several hundred feet of the adit was traversed as well as the ground in the immediate area. At no time during the traverses was any radioactivity anomaly detected in the area. Check sampling, of necessity, was confined to the ore dumps. The results of the laboratory studies of these samples are given in table 1.

Orange Hill

Orange Hill (locality 6, fig. 1) is located on the east side of the Nabesna River 12 miles south of the Nabesna airfield and 1/2 mile below the foot of Nabesna Glacier. It is elliptical in shape with its top about 600 feet above the level of the river. Access to Orange Hill is from Nabesna where the Nabesna River can be crossed in a boat or on horseback if appropriate crossings of the rapidly shifting channels of this glacial stream are selected carefully at stages of low water.

Development work done on the property includes adits, shafts, and open-cuts as well as some diamond-drill work. Many of the shafts and adits are reported to be caved. (See Van Alstine and Black, 1945.) High melt waters in the Nabesna River at the time of the author's visit made access to Orange Hill impossible. However, an airborne radioactivity traverse was flown over the mineralized area as well as over larger gossans in the region. No significant radioactivity anomalies were detected.

The oldest rocks of the region are thick lava flows and greenstones of Permian age. There is also a metamorphosed limestone of Permian age which is overlain and underlain by hornfels. The core of Orange Hill is a quartz diorite of probable Jurassic age. Dikes of alaskite intrude the quartz diorite as well as the sedimentary rocks, and all of these rocks are intruded by dacite and andesite dikes. Veinlets of quartz in the quartz diorite and silicic dikes contain pyrite, chalcopyrite, molybdenite, tetrahedrite, sphalerite, magnetite, and gypsum. The limestone has metamorphic deposits composed of magnetite, pyrrhotite, pyrite, chalcopyrite, sphalerite, and molybdenite. This summary of the geology and mineralogy at Orange Hill is taken from a detailed account by Van Alstine and Black (1945).

Chisana district

Bonanza Creek

Bonanza Creek in the Chisana district (locality 7, fig. 1) is a south-flowing creek in the Nutzotin Mountains emptying into Johnson Creek 7 miles east of the settlement of Chisana. (See fig. 2.) In this region most of the mining has been confined to placer-gold operations. At the time of the author's visit there were three small placer mines in operation. However, some lode prospecting has been done recently in several mineralized zones in the district, but at none of these prospects has there been any extensive development work done other than a few open-cuts. One prospect had two adits driven in on a mineralized zone many years ago, but these have since caved.

Moffit (1943) gives an account of the geology of this region. Briefly, volcanic and sedimentary rocks of Devonian and Permian age have been intruded by andesitic dikes and sills and by a large mass of granodiorite of Mesozoic age.

Minerals occurring in the area include galena, molybdenite, cinnabar, pyrite, copper, silver, and gold. There appears to be a definite structural control in the formation of the mineralized zones in the district. The mineralized fissure veins all have the same general trend with a strike of west to northwest and a northerly dip. The fissures are probably the result of the stresses and strains developed during the intrusion of the granodiorite into the Devonian and Permian rocks.

Investigations consisted of radioactivity traverses of the mineralized zones and rock types occurring in the district as well as collecting supplementary check samples from mineralized zones, outcrops, and stream gravels. Figure 2 shows the routes of the



Base and geology from U. S. Geol. Survey Bull. 933, plate 2




FIGURE 2.--GEOLOGIC SKETCH MAP OF THE CHISANA DISTRICT SHOWING SAMPLE LOCATIONS

EXPLANATION

SEDIMENTARY ROCKS


Unconsolidated materials



Gravel and sand



Conglomerate; sandstone with shaly partings; tuffaceous(?) sandstone; and sandy shale




Banded shale and argillite, arkosic sandstone, conglomerate and limestone



IGNEOUS ROCKS

Granitic intrusives



Basaltic rocks



○ Rock sample

● Concentrate

■ Buildings

..... Radioactivity traverse

--- Trail

QUATERNARY

TERTIARY

CRETACEOUS

TRASSIC, JURASSIC AND CRETACEOUS

MESOZOIC

PERMIAN DEVONIAN

radioactivity traverses and the location of the samples collected. The traverses conducted in the district did not indicate any anomalous radioactivity. The results of laboratory studies made on samples collected in this district are listed in tables 2 and 3.

CONCLUSIONS

Results of radioactivity investigations conducted in the eastern Alaska Range during the 1952 field season showed that no high-grade uranium deposits occur in association with the metalliferous lodes previously deemed favorable (Wedow and others, 1951) for the occurrence of uranium ores. Only three of the samples collected contain 0.003 or more percent equivalent uranium, and the radioactivity of these samples is due primarily to the accessory igneous rock minerals sphene and zircon.

Laboratory studies of the stream-gravel and placer concentrates indicated the presence of some radioactive material in one sample. The radioactivity of this sample also is due to trace amounts of sphene and zircon.

Therefore, the results of the studies made in the field and upon the samples collected indicate that there is little likelihood of finding high-grade uranium ores at the sites and prospects investigated in 1952 in the eastern Alaska Range and reported on herein. However, there are many other areas in the eastern Alaska Range which have not been studied and in which private prospecting should be encouraged.

Table 3.--Radioactivity data^{1/} on rock and ore samples collected in the Chisana district

Sample number	Location	Type of sample	Equivalent uranium (percent)
4511	Canyon in Johnson Creek	Stained material in fracture zone	< 0.001
4512	Big Eldorado Creek	Granite country rock	.001
4513	Big Eldorado Creek	Sulfide float	< .001
4514	Canyon in Bonanza Creek	Mafic dike	< .001
4516	On ridge 1 mile north of Little Eldorado Creek	Sulfide vein	< .001
4517	On ridge 1 mile north of Little Eldorado Creek	Float material	< .001
4518	On ridge 1 mile north of Little Eldorado Creek	Wall rock	< .001
4519	Canyon on lower Bonanza Creek	Sulfide vein	.001
4520	Canyon on lower Bonanza Creek	Sulfide vein	.003
4521	Canyon on lower Bonanza Creek	Sulfide vein	< .001
4525	Right limit of Johnson Creek, 1 mile west of Bonanza on hillside	Vein material	< .001
4526	Right limit of Johnson Creek, 1 mile west of Bonanza on hillside	Vein material	.001
4527	Right limit of Johnson Creek, 1 mile west of Bonanza on hillside	Vein material	< .001
4528	Right limit of Johnson Creek, 1 mile west of Bonanza on hillside	Wall rock	.003

^{1/} Equivalent-uranium analyses by J. J. Matzko and others, U. S. Geological Survey laboratory, College, Alaska.

LITERATURE CITED

- Moffit, F. H., 1938, Geology of the Slana-Tok district, Alaska: U. S. Geol. Survey Bull. 904.
- _____, 1943, Geology of the Nutzotin Mountains, Alaska, by F. H. Moffit, with a section on the igneous rocks by R. G. Wayland; Gold deposits near Nabesna, by R. G. Wayland: U. S. Geol. Survey Bull. 933-B, p. 103-199.
- Thorne, R. L., 1946, Exploration of argentiferous lead-copper deposits of the Slana district, Alaska: U. S. Bureau of Mines Rept. of Inv. 3940.
- Wedow, Helmuth, Jr., Killeen, P. L., and others, 1954, Reconnaissance for radioactive deposits in eastern interior Alaska, 1946: U. S. Geol. Survey Circ. 331.

UNPUBLISHED REPORTS

- Van Alstine, R. E., 1945, Report on the molybdenite occurrence at Rock Creek, Alaska: U. S. Geol. Survey unpublished report for the War Production Board.
- Van Alstine, R. E., and Black, R. F., 1945, Mineral deposits at Orange Hill, Alaska: U. S. Geol. Survey mimeographed report. (In open files of the U. S. Geol. Survey.)
- Wedow, Helmuth, Jr., 1951, Adaptation of portable survey meters for airborne reconnaissance with light planes in Alaska: U. S. Geol. Survey Trace Elements Memo. Rept. 323. (In open files--released through Technical Information Service, U. S. Atomic Energy Commission, Oak Ridge, Tennessee.)
- Wedow, Helmuth, Jr., and others, 1951, Interim report on an appraisal of the uranium possibilities of Alaska: U. S. Geol. Survey Trace Elements Memo. Rept. 235. (In open files of the U. S. Geol. Survey.)

CHAPTER B.--HOPE CREEK AND MILLER HOUSE-CIRCLE HOT SPRINGS AREAS

By Walter S. West and John J. Matzko

INTRODUCTION

The Hope Creek and Miller House-Circle Hot Springs areas, Circle quadrangle, Alaska, were investigated for the occurrence of uranium deposits by the U. S. Geological Survey in 1952.

The Hope Creek area is located approximately 55 miles northeast of Fairbanks, Alaska, and about 10 miles north of the Steese Highway (fig. 1). The area constitutes the headwaters sources of Hope, American, Champion, Little Champion, Nome, and Sourdough Creeks. The nearest accessible point to Hope Creek is 5 miles north of mile-post 67 on the Steese Highway. This point lies at the termination of a tractor trail along Sourdough Creek which will accommodate jeeps and trucks. (See fig. 3.)

The Miller House-Circle Hot Springs area is about 100 miles east-northeast of Fairbanks, Alaska (fig. 1), and is drained by Mammoth, Independence, Bedrock, Boulder, Deadwood, Holdem, Ketchem, Hot Springs, Portage, and Half Dollar Creeks. Part of the area is accessible from the Steese Highway by roads and trails up Mammoth, Independence, Boulder, Deadwood, Ketchem, Portage, and Half Dollar Creeks. An airfield at Circle Hot Springs will accommodate small planes. (See fig. 4.)

Fluorite, suggestive of the possibility of the presence of associated uranium minerals, has been reported in the Hope Creek area (Prindle, 1910). Uraniferous fluorite and several other uranium-bearing minerals were found in the Miller House-Circle Hot Springs area in 1949 by White and Tolbert (Wedow, White, and others, 1954).

These areas were examined for the possibilities of lode concentrations of uranium during the summer of 1952 by Walter S. West, geologist, and George M. Haselton, geologic field assistant. Part of the work in the Miller House-Circle Hot Springs area was done by John J. Matzko, geologist, and Fred Freitag, field assistant.

Field work in the Hope Creek area consisted of radioactivity traversing on foot with standard portable survey meters equipped with interchangeable beta-gamma and gamma probes, and of the collection of rock and mineral samples. Helicopter support for part of the work was furnished by the Topographic Division of the U. S. Geological Survey.

The examination of the Miller House-Circle Hot Springs area was accomplished by both foot and car-borne radioactivity traverses, and by the collection of rock, mineral, placer, and water samples. The equipment used for measuring radioactivity is the same as that described in Chapter A, page 7.

The equivalent-uranium analyses and mineralogic determinations were made at the Geological Survey's laboratories at College, Alaska, and Washington, D. C.

This work was done on behalf of the Division of Raw Materials of the U. S. Atomic Energy Commission.

AREAS INVESTIGATED

Hope Creek area

The geology of the Hope Creek area has been described by Prindle (1910, 1913) and Mertie (1937).

The bedrock in the area consists of the Birch Creek schist of pre-Cambrian age intruded by granitic rocks and dikes of Mesozoic(?) age (fig. 3).

Erosion has exposed granite at two localities within the area: one small exposure consisting entirely of granite talus near the headwaters of Sourdough Creek; the other, a larger mass of granite, forming pinnacles and talus on the tops and sides of the high ridges, at the headwaters of Hope, American, Champion, Little Champion, and Nome Creeks. The granite at both localities is a biotite-tourmaline variety. The texture of the granite ranges from medium-grained equigranular to coarse-grained porphyritic. The accessory minerals include zircon, sphene, garnet, and pyrrhotite. Quartz and quartz-tourmaline veins occur in the granite. Granitic and aplitic dikes ranging from slightly less than one inch to six feet in width are of rather common occurrence in the granite.

The Birch Creek formation consists mostly of quartz-mica schist in the Hope Creek area. The occurrence of quartz-pyrite-fluorite veins in the schist adjoining the granite has been reported (Prindle, 1910; Mertie, 1937) but was not confirmed during the 1952 investigation. Pyrite was observed at several places along the granite-schist contact.

Radioactivity traverses in the Hope Creek area indicated that the granite bedrock and granite talus were more radioactive than the adjacent schist bedrock and schist talus. For this reason sampling was confined to the granitic rock localities. Sample locations are shown on figure 3. Equivalent-uranium analyses of unconcentrated crushed rock samples from the Hope Creek area are given in table 4.

Laboratory studies revealed that most of the radioactivity was confined to the minerals present in the heavy-mineral fractions (those greater than 3.3 sp. gr.) of the samples. The mineralogy of the heavy-mineral fractions of selected samples is given in table 5.

Table 4.--Equivalent-uranium analyses^{1/} of samples, Hope Creek area, Alaska

Sample number	Location	Type of material	Equivalent uranium (percent)
4470	Headwater of Sourdough Creek	granite talus	0.003
4471	Divide between the north headwater fork of Hope Creek and the south headwater fork of American Creek	granite bedrock	.004
4472	-----do-----	granite bedrock	.003
4473	-----do-----	granite bedrock	.004
4474	Ridge between the south and middle headwater forks of American Creek	granite talus	.055
4475	-----do-----	granitic dike	.003
4476	-----do-----	quartz veins	.002
4477	Left limit, north fork of Hope Creek	granite bedrock	.004
4478	Granite-schist contact between the middle and north headwater forks of American Creek	granite and schist talus	.002
4479	Divide between American and Champion Creeks	granite bedrock	.003
4480	Divide between the middle and north headwater forks of American Creek	aplitic dike	.002
4481	-----do-----	granitic dike	.003
4482	Left limit, north fork of American Creek	granitic dike	.004
4483	Granite-schist contact at the head of the north fork of American Creek	granitic talus	.001
4484	Saddle on divide between the north headwater fork of Hope Creek and Nome Creek	granite bedrock	.003
4485	Divide between the south headwater fork of Hope Creek and Nome Creek	granitic dike talus	.004
4486	Headwaters of Sourdough Creek	granite talus	.003
4487	Headwaters of American Creek	quartz-tourmaline vein	.002
4488	-----do-----	granite bedrock	.003
4489	-----do-----	granite talus	.001
4490	-----do-----	granite bedrock	.003
4491	Headwaters of the north fork of Hope Creek	granite bedrock	.003

^{1/} By John J. Matzko and others, U. S. Geological Survey laboratory, College, Alaska.

Table 5.--Mineralogy^{1/} of heavy-mineral fractions (those greater than 3.3 specific gravity) of selected samples from the Hope Creek area, Alaska

Minerals	Estimated volume percent of minerals present in samples									
	4470	4471	4473	4474	4477	4478	4482	4483	4486	4487
Allanite(?)	--	--	--	--	9	--	--	--	--	--
Anatase	--	--	--	--	--	1	--	--	--	--
Apatite	--	--	tr ^{2/}	--	--	--	--	tr	--	--
Biotite	--	2	--	--	10	2	23	3	--	8
Diopside	--	tr	--	--	tr	--	--	--	--	--
Epidote	2	--	--	--	--	--	--	--	7	tr
Fluorite	--	--	1	--	--	--	--	--	tr	tr
Galena	--	tr	--	--	--	--	--	--	--	--
Garnet	1	--	--	--	--	--	tr	--	1	--
Goethite	--	--	10	X ^{3/}	--	--	--	25	--	--
Hematite	10	--	--	X	--	--	--	16	15	tr
Hornblende	9	--	--	X	--	--	--	--	12	--
Ilmenite	--	3	--	--	--	--	--	4	--	--
Limonite	1	--	--	--	--	21	--	11	tr	--
Malachite	--	--	--	--	--	--	tr	--	--	--
Molybdenite	--	tr	--	--	--	--	--	--	--	--
Muscovite	--	--	tr	--	--	--	--	--	--	tr
Pyrite	tr	tr	--	--	--	--	--	tr	tr	--
Pyrrhotite	50	40	1	--	20	20	50	16	45	50
Rutile	--	--	80	--	--	--	--	--	--	--
Scheelite	tr	--	--	--	--	--	--	--	tr	--
Sphene	15	1	--	--	--	--	2	--	20	--
Stibnite	1	--	--	--	--	1	--	--	--	--
Topaz	--	50	--	--	60	25	12	tr	--	--
Tourmaline	1	--	tr	X	--	30	11	25	--	40
Zircon	10	4	7	X	1	tr	2	--	tr	2

1/ by W. S. West and J. J. Matzko

2/ tr - trace

3/ X indicates presence of mineral in sample, amount not determined

Sample 4474, containing 0.055 percent equivalent uranium, was the most radioactive sample collected in the Hope Creek area. The rock occurs as float in a small saddle between two large granite pinnacles on the high divide between the middle and south headwater forks of American Creek (fig. 3). The granite float rock in this sample contains such an intricate network of hematite-goethite veinlets that it appears to be brecciated. No uranium-bearing minerals could be identified. An alpha plate made of the sample indicates that the radioactivity is probably due to a radioactive element or elements occurring as impurities in the goethite or in very minute discrete minerals mixed with the goethite but which are not discernible under the petrographic microscope. Zircon probably contributes a small amount of radioactivity to the sample.

The radioactivity of the remainder of the Hope Creek area samples is attributed to the presence of zircon, sphene, hematite, and allanite(?). In addition to these minerals, fluorite, topaz, limonite, apatite, garnet, scheelite, and malachite may contain trace amounts of uranium as an impurity as is the case in the Miller House-Circle Hot Springs area. However, none of the minerals from the Hope Creek area samples have been subjected to fluorimetric tests for uranium.

Niobium and tantalum have been detected spectroscopically in trace amounts in the heavy-mineral fraction of sample 4473 (fig. 3) and may be present in the rutile which constitutes an estimated 80 percent of this fraction of the sample (table 5).

Miller House-Circle Hot Springs area

The geology of the Miller House-Circle Hot Springs area has been described by Johnson (1910), Prindle (1913a), and Mertie (1937, 1938).

Bedrock in the area includes metamorphic and igneous rocks consisting of Birch Creek schist of pre-Cambrian age intruded by a granite stock and dikes of Mesozoic(?) age (fig. 4).

The exposed part of the igneous stock is composed principally of coarse-grained granular to porphyritic biotite granite. The common accessory minerals are zircon, sphene, allanite, garnet, scheelite, and pyrrhotite. The granite stock has a greater areal extent than has been shown on any previous geologic map of the region (Prindle, 1913a, pl. 2; Mertie, 1938, pl. 1).

The Birch Creek schist is mainly of the mica and quartz-mica varieties. Feldspathic, carbonaceous, and chloritic schists are also present in the area. Granitic and aplitic dikes, genetically related to the granite stock, occur in the granite and adjoining schist. The dikes range in width from a few inches to two feet.

Numerous quartz veins, ranging from a fraction of an inch to several feet in width, occur in the schist and to a lesser extent in the granite. Most of the veins show no signs of mineralization. However, pyrite, arsenopyrite, galena, wolframite, and gold have been found in a few of these veins.

Stream and bench gravels in the valleys of Porcupine, Bonanza, Miller, Mastodon, Independence, Mammoth, Crooked, Boulder, Deadwood, Switch, Ketchem, Portage, and Half Dollar Creeks and the North Fork of Harrison Creek have been mined for placer gold. During 1952 the only active mining in the area was on Independence, Mastodon, Crooked, Deadwood, and Portage Creeks.

Foot and car-borne radioactivity traversing in the Miller House-Circle Hot Springs area disclosed that the granite localities are more radioactive than those composed of schist. This conclusion was borne out by the results of radioactivity studies of the samples

collected in the area. The sample locations are shown on figure 4. Equivalent-uranium analyses of representative samples are given in table 6; with the exception of placer samples, the analyses listed in table 6 were made on unconcentrated crushed rock and vein material.

As shown in table 6, the equivalent-uranium content ranges from 0.001 to 0.005 percent for the granite samples, 0.002 to 0.004 percent for the dike-rock samples, 0.001 to 0.002 for the schist samples, and 0.001 to 0.01 percent for the placer samples. The equivalent-uranium content averages 0.004 percent for the granite samples and 0.003 percent for the dike-rock samples.

A limited number of water samples were taken for experimentation in the use of a water-sampling technique to locate lode concentrations of uranium in an area where radioactivity traversing and rock sampling were of no avail because of the wide-spread cover of disintegrated bedrock, alluvium, and vegetation. The analyses are shown in table 7.

Table 7 shows that sample 52AMz 15 from Portage Creek contains a relatively high uranium content compared to the average uranium content of fresh waters which is 1×10^{-8} percent (Irving May, personal communication) and the average uranium content of sea waters which is 3×10^{-7} percent (Nakanishi, 1951; and Pietsch and Grimaldi, 1952). This may indicate the desirability of using a more detailed water sampling technique in this and other areas where large expanses of ground cover effectively shield any possible radiation coming from the underlying rocks.

Radioactivity and mineralogic examinations of the rock, mineral, and placer samples indicate that most of the radioactive minerals are present in the heavy-mineral

Table 6.--Equivalent-uranium analyses^{1/} of samples, Miller House-Circle Hot Springs area, Alaska

Sample number	Location	Type of material	Equivalent uranium (percent)
4696	Independence Creek -----	placer	0.001
4699	Mammoth Creek, right limit-----	granite talus	.004
4698	-----do-----	dike talus	.003
4697	-----do-----	granite talus	.005
4722	Divide between Mammoth and Bedrock Creeks	granite talus	.005
4721	-----do-----	granite talus	.004
4720	Bedrock Creek, left limit -----	granite bedrock	.005
4719	-----do-----	granite bedrock	.005
4718	-----do-----	granite bedrock	.005
4717	-----do-----	vein in granite	.003
4716	-----do-----	granite bedrock	.004
4715	-----do-----	dike talus	.001
4714	-----do-----	granite bedrock	.004
4676	Boulder Creek, left limit-----	granite bedrock	.004
4705	-----do-----	granite bedrock	.004
4704	-----do-----	granite talus	.003
4702	-----do-----	granite bedrock	.003
4701	-----do-----	granite bedrock	.004
4706	Boulder Creek, stream bed -----	granitic dike	.004
4709	-----do-----	granite boulder	.005
4590	Deadwood Creek, 1.5 mi. below Switch Creek	schist bedrock	< .001
4690	Deadwood Creek, 0.5 mi. below Switch Creek	weathered dike	.003
4691	-----do-----	granite bedrock	.005
4692	-----do-----	granite bedrock	.003
4693	-----do-----	granite bedrock	.004
4689	Deadwood Creek, 0.25 mi. below Switch Creek	vein in granite	.005
4591	Deadwood Creek, 0.4 mi. below Discovery Creek	schist bedrock	< .001
4587	Discovery Creek, trib. to Deadwood Cr.	granite talus	.004
4588	-----do-----	granite talus	.001
4589	-----do-----	granite talus	.005
4600	-----do-----	placer	.005
4596	Deadwood Creek, 0.5 mi. upstream from road end	schist bedrock	.002
4597	Deadwood Creek, 1 mi. upstream from road end	schist bedrock	.001
4599	Deadwood Creek, west trib. below 25 Pup	schist bedrock	.001
4598	Deadwood Creek, 2 mi. upstream from road end	schist bedrock	< .001

See footnote at end of table.

Table 6.--Equivalent-uranium analyses^{1/} of samples, Miller House-Circle Hot Springs area, Alaska--Continued

Sample number	Location	Type of material	Equivalent uranium (percent)
4677	Portage Creek, Carstens' mining operation	granite bedrock	0.005
4678	-----do-----	mafic dike	.002
4679	-----do-----	placer	.01
4680	-----do-----	tin-iron nuggets	<.001
4688	Portage Creek, left limit-----	granite bedrock	.003
4687	-----do-----	granite talus	.003
4595	Divide between Portage and Half Dollar Creeks	granite talus	.002
4685	Half Dollar Creek, right limit -----	granite bedrock	.003
4683	-----do-----	granitic dike	.003
4681	-----do-----	granite bedrock	.002
4592	-----do-----	aplite dike	.003

^{1/} By J. J. Matzko and others at the Geological Survey laboratory, College, Alaska

Table 7. --Analyses^{1/} of water samples, Miller House-Circle Hot Springs area, Alaska

Sample number	Location	Uranium (percent)	Uranium (ppb)
52AMz 5	Discovery Creek-----	0.33×10^{-7}	0.33
52AMz 6	Switch Creek, tributary to Deadwood Creek	$.42 \times 10^{-7}$.42
52AMz 10	Deadwood Creek-----	1.56×10^{-7}	1.56
52AMz 11	-----do-----	2.01×10^{-7}	2.01
52AMz 12	Ketchem Creek-----	1.11×10^{-7}	1.11
52AMz 13	Holdem Creek, tributary to Ketchem Creek	1.74×10^{-7}	1.74
52AMz 14	Hot Springs Creek-----	1.44×10^{-7}	1.44
52AMz 15	Portage Creek ^{2/} -----	40.2×10^{-7}	40.2

^{1/} By Ivan Barlow, U. S. Geological Survey laboratory, Washington, D. C., September, 1952

^{2/} Original sample contained much clay-size material in suspension because of sluicing operation upstream from sample site; therefore, uranium content may be high because all solids may not have been removed in subsequent decanting to obtain clear sample, or because more uranium was taken into solution during the period of sluicing when more solid material than normal was in suspension and subject to corrosion. Another factor may be that some additional uranium was taken into solution from the suspended solids when the sample was acidified prior to decanting.

fractions (those greater than 3.3 specific gravity) of these samples. The minerals found in the heavy-mineral fractions of selected samples are listed in table 8.

Mineralogic studies of the samples revealed that fluorite, which was previously reported in the granite bedrock only on Deadwood Creek, also occurs in vugs of granitic rock on Portage and Boulder Creeks.

Fluorimetric tests proved that fluorite, limonite, malachite, zircon, sphene, topaz, allanite, garnet, uranothorianite, and scheelite contained in some of the Miller House-Circle Hot Springs samples were uraniferous. Other minerals listed in table 8, such as hematite, goethite, monazite, xenotime, pyrochlore-microlite, and some of the sulfides, may contain impurities or intergrowths of uranium but none have yet been subjected to fluorimetric examinations. The biotite in many of the samples contains pleochroic halos. The distribution of the uranium-bearing minerals in samples from the area is summarized in table 9.

In addition to the minerals listed in tables 8 and 9, a non-fluorescent, yellow-green, uranium mineral was found in granite bedrock, sample 4677 (fig. 4), from H. C. Carstens' gold-mining operation on Portage Creek. The mineral could not be identified because an insufficient quantity was available for X-ray determination. Practically all of the biotite in this sample contains pleochroic halos. The ilmenite contains trace amounts of niobium.

The only occurrence of uranothorianite, the most important uranium-bearing mineral found in the area, was in sample 4679, a sluice-box concentrate also from Carstens' mining operation on Portage Creek. The uranothorianite occurs as very small, black cubes, principally in the minus 100- plus 200- mesh size fraction of the placer concentrate. This fraction of the sample contains 0.091 percent

Table 8.--Mineralogy^{1/} of the heavy-mineral fractions of selected samples from the Miller House-Circle Hot Springs area, Alaska

Minerals	Estimated volume percent of minerals present in samples												
	4696	4697	4718	4701	4709	4691	4689	4600	4596	4677	4679	4685	4683
Allanite	tr ^{2/}	3	--	45	50	--	--	--	--	--	tr	25	20
Amphibole	--	11	--	--	--	--	--	--	--	--	--	--	--
Anatase	--	--	--	--	--	10	--	--	--	--	--	--	--
Apatite	--	--	--	--	--	--	--	1	--	tr	--	--	--
Arsenopyrite	tr	--	--	--	--	--	--	--	--	tr	1	--	--
Azurite	--	tr	--	--	--	--	--	--	--	--	--	--	--
Biotite	--	2	24	25	15	1	--	--	--	10	tr	tr	--
Bismuthinite	--	--	--	--	--	--	--	--	--	--	tr	--	--
Cassiterite	3	--	--	--	--	--	1	--	--	--	3	--	--
Cerussite	--	--	--	--	--	--	--	--	--	--	--	fr	--
Chalcopyrite	--	--	--	--	--	--	--	--	--	tr	--	--	--
Chlorite	--	tr	--	15	tr	2	--	--	--	7	tr	--	tr
Diopside	tr	--	--	--	--	--	--	--	--	--	3	--	--
Epidote	--	--	--	1	--	--	--	4	--	--	--	1	3
Fluorite	--	--	--	tr	--	--	--	--	--	1	--	--	--
Galena	tr	tr	--	--	--	10	--	--	--	--	--	--	--
Garnet	40	4	1	1	--	--	--	6	--	1	10	2	6
Goethite	--	--	--	--	--	50	35	--	--	--	--	--	--
Gold	tr	--	--	--	--	--	--	--	--	--	tr	--	--
Hematite	--	9	--	tr	--	--	40	1	5	2	5	15	--
Ilmenite	12	--	10	--	--	5	22	5	75	60	50	--	--
Jamesonite	--	--	--	--	--	--	--	--	--	tr	--	--	--
Leucoxene	--	--	--	--	--	10	--	--	--	--	--	--	--
Limonite	4	--	--	tr	--	--	--	3	--	tr	--	--	10
Magnetite	30	--	--	--	--	--	--	tr	--	--	15	--	--
Malachite	--	tr	tr	--	--	--	--	--	--	--	--	--	--
Molybdenite	--	tr	--	--	--	--	--	--	--	--	--	--	--
Monazite	--	--	10	--	--	--	--	--	--	4	tr	--	--
Muscovite	2	--	--	--	--	--	--	--	10	--	tr	--	tr
Pyrite	2	1	--	tr	--	1	1	--	4	1	1	2	1
Pyrochlore-microlite	--	--	--	--	--	--	--	--	--	--	tr	--	--
Pyrrhotite	--	50	35	7	30	tr	--	--	--	--	--	40	55
Rock forming minerals ^{3/}	--	--	--	--	--	--	--	--	--	--	--	15	--
Rutile	1	--	--	--	--	5	--	--	4	--	--	--	--
Scheelite	2	tr	tr	--	--	--	--	tr	--	tr	1	tr	tr
Schist fragments	3	--	--	--	--	--	--	70	--	--	--	--	--
Sphalerite	--	--	--	--	--	--	--	--	--	1	--	--	--
Sphene	--	--	--	--	tr	2	--	--	--	--	1	--	tr
Spinel	--	--	--	--	--	--	--	2	--	--	1	--	--
Topaz	--	20	5	--	--	--	--	--	--	--	tr	--	--
Tourmaline	tr	--	--	--	--	--	--	2	--	--	1	--	--
Uranothorianite	--	--	--	--	--	--	--	--	--	--	tr	--	--
Wolframite	1	--	--	--	--	--	--	tr	--	--	5	--	--
Xerotime	tr	--	--	--	--	--	--	--	--	--	--	--	--
Zircon	tr	--	15	6	5	2	1	1	2	10	1	tr	5

^{1/} by W. S. West and J. J. Matzko^{2/} tr - trace^{3/} mostly quartz and feldspar

Table 9.--Uranium-bearing minerals in samples from the Miller House-Circle Hot Springs area, Alaska

Locality and type of sample	Uranium-bearing minerals										
	Allanite	Apatite	Fluorite	Garnet	Limonite	Malachite	Scheelite	Sphene	Topaz	Uranothorianite	Zircon
Independence Creek: Placer-----	X	-	-	X	X	-	X	-	-	-	X
Mammoth Creek: Granite-----	X	-	-	X	-	X	X	-	X	-	-
Bedrock Creek: Granite-----	-	-	-	X	-	X	X	-	X	-	X
Boulder Creek: Granite-----	X	-	X	X	X	-	-	X	-	-	X
Deadwood Creek: Granite-----	-	-	X ^{1/}	-	-	-	-	X	-	-	X
Placer ^{2/} -----	-	-	-	-	-	-	X	-	-	-	-
Discovery Creek: Placer-----	-	-	-	X	-	-	X	-	-	-	X
Ketchum Creek: Placer ^{2/} -----	X	-	-	X	-	-	X	X	X	-	-
Hot Springs Creek: Placer ^{2/} -----	X	-	-	-	-	-	-	X	-	-	-
Portage Creek: Granite-----	-	X	X	X	X	-	X	-	-	-	X
Placer-----	X	-	-	X	-	-	X	X	X	X	X
Half Dollar Creek: Granite-----	X	-	-	X	X	-	X	X	-	-	X

^{1/} Mineral present in sample collected by White and Tolbert in 1949

^{2/} Samples collected by White and Tolbert in 1949

equivalent uranium. The uranothorinite appears to be responsible for most of the radioactivity of the sample. An assay made on part of this sluice-box concentrate at the Assay Office of the Territorial Department of Mines, Fairbanks, Alaska, gave 4.65 percent tungsten trioxide (WO_3). Most of the tungsten is in the form of wolframite with only minor amounts of scheelite.

Although no quantitative analyses for uranium and thorium were made on the samples from the Miller House-Circle Hot Springs area, mineralogic studies indicate that most of the radioactivity of these samples is apparently due to uranium rather than thorium.

CONCLUSIONS

No lode or placer deposits in the Hope Creek and Miller House-Circle Hot Springs areas were found to contain sufficient quantities of uranium-bearing minerals to be of commercial value.

A wide variety of radioactive minerals were found to occur in granite. Certain uranium-bearing minerals in the Miller House-Circle Hot Springs area, which are also believed to contain uranium in the Hope Creek area, appear to be primary accessory minerals in the granite. Other minerals, like fluorite, topaz, and several metallic sulfides in both areas, and cassiterite and wolframite in the Miller House-Circle Hot Springs area, were probably formed as a result of pneumatolytic action after the crystallization of the magma or during the late stages of crystallization. As some of the minerals in the latter group are known to be uranium-bearing, uranium, in part, seems to have been introduced by hydrothermal solutions during the process of pneumatolysis.

On the basis of present information the Hope Creek area does not appear to be favorable for the occurrence of high-grade uranium ores. It should be pointed out, however, that the work done in 1952 consisted only of a brief reconnaissance, primarily to locate and test reported fluorite occurrences. Prospectors searching the area for other metals should keep in mind the possible association of uranium with mineralized zones containing hematite of hydrothermal origin, with minerals containing silver, cobalt, nickel, bismuth, and fluorine, and to a lesser extent with lode deposits containing copper, tin, lead, molybdenum, and gold.

The investigation conducted by the Geological Survey in 1949 (Wedow, White, and others, 1954) and in 1952 (this report) show that there is little hope of discovering commercial concentrations of uranium in the Miller House-Circle Hot Springs area by established methods of radioactivity traversing with portable survey meters, primarily because of the widespread cover of vegetation, soil, and disintegrated bedrock. On the other hand, this area, particularly the watershed of Portage Creek, cannot be ruled as unfavorable for the occurrence of uranium in lode deposits because of the relatively high uranium content of water and the presence of uranothorianite in concentrates from Portage Creek. (See tables 7 and 8.) Prospectors interested in the uranium possibilities of the Miller House-Circle Hot Springs area will probably find that geochemical methods of prospecting, such as water, soil, and vegetation sampling, would be the best techniques to use in the search for uraniferous lodes in the area. Also, the testing of heavy-mineral concentrates from gravels, slope wash, and disintegrated bedrock for radioactivity, in a fashion similar to that used by West (1953) in the Darby Mountains of the Seward Peninsula, would aid considerably in localizing the occurrence of the uranothorianite on Portage Creek.

LITERATURE CITED

- Johnson, B. L., 1910, Occurrence of wolframite and cassiterite in the gold placers of Deadwood Creek, Birch Creek district (Alaska): U. S. Geol. Survey Bull. 442-F, p. 246-250.
- Mertie, J. B., Jr., 1937, The Yukon-Tanana region, Alaska: U. S. Geol. Survey Bull. 872.
- _____, 1938, Gold placers of the Fortymile, Eagle, and Circle districts, Alaska: U. S. Geol. Survey Bull. 897-C, p. 133-261.
- Nakanishi, M., 1951, Uranium content of sea water: Chem. Soc. Japan, v. 24, no. 1, p. 36.
- Pietsch, Audrey, and Grimaldi, F. S., 1954, The fluorimetric determination of uranium in nonsaline and saline waters: U. S. Geol. Survey Bull. 1006, pt. 17.
- Prindle, L. M., 1910, Sketch of the geology of the northeastern part of the Fairbanks quadrangle (Alaska): U. S. Geol. Survey Bull. 442-F, p. 203-209.
- _____, 1913, A geologic reconnaissance of the Fairbanks quadrangle, Alaska: U. S. Geol. Survey Bull. 525.
- _____, 1913a, A geologic reconnaissance of the Circle quadrangle, Alaska: U. S. Geol. Survey Bull. 538.
- Wedow, Helmuth, Jr., White, M. G., and others, 1954, Reconnaissance for radioactive deposits in east-central Alaska, 1949: U. S. Geol. Survey Circ. 335.
- West, W. S., 1953, Reconnaissance for radioactive deposits in the Darby Mountains, Seward Peninsula, Alaska, 1948: U. S. Geol. Survey Circ. 300.

CHAPTER C.--KOYUKUK-CHANDALAR REGION

By Arthur E. Nelson

INTRODUCTION

The presence of radioactive materials is known in two localities in the Koyukuk-Chandalar region (Wedow and others, 1951; White, 1952). Accordingly, radioactivity reconnaissance investigations were undertaken in the Chandalar mining district and at the Gold Bench mining area (fig. 1). Airborne radioactivity traverses were conducted over a part of the region while flying the party into the region and returning them to Fairbanks.

Radioactivity traverses and outcrop tests were made in and around the mines and prospects, mineralized zones, and the country rock at the two localities, and samples were collected for equivalent-uranium determinations and mineralogic studies. Stream gravels were sampled to determine if radioactive minerals are being liberated from the bedrock of areas drained by the streams.

A standard portable survey meter was modified to accept interchangeable 6-inch beta-gamma and 2- by 20-inch gamma probes; these respective probes were used for individual outcrop tests and normal ground traverses. For airborne traverses a standard survey meter coupled with six 2- by 40-inch gamma probes, connected in parallel, was mounted in a light aircraft (Wedow, 1951).

The field work in this area was conducted in late August 1952, by Arthur E. Nelson, geologist, and Richard S. Smith, geologic field assistant. This work was done on behalf of the Division of Raw Materials of the U. S. Atomic Energy Commission.

GEOLOGY

The Koyukuk-Chandalar region (fig. 1) lies in the southern foothills of the Endicott Mountains which are a part of the Brooks Range. The region is drained by the south-flowing Koyukuk River on the west and the Chandalar River on the east, both of which empty into the Yukon River. To the south the area is bounded by the Koyukuk flats and the Hodzana highlands; to the north, by the crest of the Endicott Mountains.

Maddren (1913) gives an account of the geology of the Koyukuk-Chandalar region, and Mertie (1925) gives a detailed account of the Chandalar mining district; hence, only a summary of the main geologic features is presented herein.

The rocks of the region consist of a series of sedimentary strata which have been folded and faulted and range from pre-Cambrian or pre-Ordovician to Cretaceous in age and igneous rocks of Paleozoic and Mesozoic age. Generally, the oldest sedimentary rocks occur in the southernmost extremities of the district and are overlain by successively younger sedimentary beds to the north.

The southernmost belt of sedimentary rocks, consisting chiefly of quartzite and quartzitic schist of pre-Cambrian or pre-Ordovician age, is folded along a general east-west axis and has a northerly dip. Overlying the quartzite and quartzitic schist is a sedimentary sequence of Paleozoic age now metamorphosed to schist and phyllite. These latter rocks have the greatest areal distribution in the region and have the same general attitude as the underlying quartzite and quartzitic schist. Silurian limestone overlies the metamorphic rocks of Paleozoic age and is in turn overlain by Devonian and Mississippian slates, sandstones, and limestones. In the southern part of the district Cretaceous conglomerates, sandstones, and shales lie unconformably on the


older rocks. The igneous rocks in the region consist of late Silurian or early Devonian granitic gneiss and a granodiorite of possible Mesozoic age. Greenstones of upper Paleozoic age occur as dikes, sills, and flows and are associated with all rocks of Paleozoic and pre-Paleozoic (?) age.

AREAS INVESTIGATED

No significant radioactivity anomalies were noted during any of the airborne traverses conducted in the Koyukuk-Chandalar region. The airborne traverses were made over many mineralized zones located in and around granitic masses which were not easily accessible otherwise. The majority of these areas are located in the region drained by the North Fork of the Koyukuk River. In general that area drained by the South Fork of the Koyukuk River has a heavy cover of overburden, and it was impossible to get effective radioactivity coverage; thus, little traversing was done there.

Chandalar mining district

The Chandalar mining district (fig. 5) is located about 6 miles east-northeast of the head of Lake Chandalar and comprises an area of about 100 square miles. Prospecting for placer- and lode-gold deposits has been carried on intermittently in this district since the turn of the century. Placer mines have been located on four creeks in the district, including Little Squaw, Big Squaw, and Tobin Creeks. Development work on numerous quartz veins in the district consists of adits, shafts, and open-cuts, most of which are now inaccessible. Hence, all that can be seen of the geological relationships is at the surface outcrops.



0 1 2 3 4 5 Miles

Scale

EXPLANATION

- | | | | |
|-------|------------------------|-----|-------------|
| | Radioactivity traverse | — — | Trail |
| ○ | Rock sample | ● | Concentrate |
| ⌘ | Lode mine | × | Placer mine |
| ■ | Building | | |

FIGURE 5.--SKETCH MAP OF THE CHANDALAR MINING DISTRICT SHOWING SAMPLE LOCATIONS

Mining activity in recent years has been somewhat limited; in 1952 only five men were engaged in either prospecting or mining operations. One man was doing some prospecting on a very limited scale, and two men were developing a quartz vein on the divide between Big and Little Squaw Creeks. Sluicing at a small placer mine on Tobin Creek and at another on Big Creek, each a one-man operation, was considerably hampered because of a limited water supply.

Sedimentary and igneous rocks of Paleozoic age were the only formations seen in the district besides the unconsolidated stream and glacial deposits. The sedimentary rocks are primarily mica schist with minor amounts of chlorite schist. The schist is cut by greenstone dikes and locally by small bodies of granitic gneiss. Numerous gold-bearing quartz veins occur in the schist and are apparently confined to four main zones, all of which have a westerly trend, but some dip to the north and others to the south. Locally minor veins are subsidiary to the more persistent ones. These subsidiary veins have different trends which are probably the result of local stresses superimposed upon regional structure.

Radioactivity traverses were made across and along mineralized zones, ore dumps, and adits; specific outcrop tests were made and a general traverse conducted in the main drainage systems of the district. At no point along any of the traverses was a significant anomaly noted. Samples were collected from these areas for further study in the laboratory. The results of laboratory studies of the samples are summarized in tables 10 and 11.

Laboratory studies found traces of monazite, zircon, and uranothorianite in certain stream samples collected during the summer field season of 1952, as indicated in table 10. A few grains of pyrite and galena and one grain of molybdenite were

Table 10.--Radioactivity ^{1/} of stream and placer concentrate samples from the Koyukuk-Chandalar region, Alaska

Sample number	Location	Type of sample	eU ^{2/} (percent)	eU (percent) I. H. ^{3/}	eU (percent) of Magnetic fractions of I. H.							Radioactive minerals	
					0.1 amp	0.4 amp	0.5 amp	0.7 amp	1.0 amp	1.6 amps mag.	1.6 amps non-mag.		
4632	Tobin Creek	stream	<0.001	0.01	X ^{4/}	X	0.10 (monazite)						Monazite in trace amounts
4635	Little Squaw Creek	stream	<.001	.002	X		X	0.005 (monazite)					Monazite - only a few grains
4637	East fork of Big Squaw Creek	stream	<.001	<.001	X								
4638	Middle fork, Big Squaw Creek	stream	.001	.018	X		X (monazite)		3.2 (uranothorianite)		X (zircon)		Uranothorianite in trace amounts with a few grains of monazite and zircon
4639	West fork, Big Squaw Creek	stream	<.001	.003	X								
4640	Unnamed stream west of Big Squaw Creek	stream	<.001	.005	X		X						
4641	Gold Bench, South Fork of the Koyukuk River	stream	<.001	.015	X	X		X	.05 (sphene and thorite(?))	X	0.08 (zircon)		Sphene, zircon, and thorite(?) in trace amounts
4642	-----do-----	placer	.18		X								Uranothorianite with trace amounts of sphene, zircon, and monazite

^{1/} equivalent-uranium analyses and mineral identifications by J. J. Matzko and A. E. Nelson

^{2/} percent equivalent uranium

^{3/} mineral fraction heavier than methylene iodide (sp. gr. 3.3)

^{4/} indicates that magnetic fraction contains less than 0.003 percent equivalent uranium

Table 11.--Radioactivity^{1/} of rock and ore samples from the Koyukuk-Chandalar region,
Alaska

Sample number	Location	Type of sample	Equivalent uranium (percent)
4631	Mine at the head of Tobin Creek	ore dump	≤ 0.001
4633	Valley of upper Tobin Creek	igneous float	.001
4634	Mine on divide between Big Creek and Little Squaw Creek	vein	< .001
4636	Mine on divide between Little Squaw and Big Squaw Creeks	ore dump	.001

^{1/} Equivalent-uranium analyses made in Geological Survey laboratory at College, Alaska.

found in the sample containing the uranothorianite. This suggests the possibility of sulfide minerals occurring in veins in the drainage basin of Big Squaw Creek which may be the probable source of the uranothorianite.

Gold Bench

Gold Bench, a high-level deposit of stream gravels, is situated on the northwest side of the South Fork of the Koyukuk River (fig. 1). Placer gold has been mined from these gravels since about 1899, but no mines were being operated in 1952. The gold-bearing gravels consist mostly of schist fragments with varying amounts of quartz, chert, and igneous rocks and overlie a thick sequence of wash deposits. The size of the unconsolidated sediments (gravel and wash) ranges from coarse- to fine-grained. The deposits are crudely stratified and rise to a height of about 200-300 feet on the south side of the river. Little is known of the geology in the vicinity of Gold Bench, and the actual source of the gravels is unknown at the present time. The following minerals have been observed by White (1952, p. 11) in concentrates from Gold Bench: magnetite, garnet, hematite, zircon, olivine, epidote, sphene, pyrite, scheelite, galena, chalcopyrite, rutile, cinnabar, cassiterite, bismuthinite(?) and thorianite(?).

During the 1952 investigations, a radioactivity traverse of the entire mining area of Gold Bench was made, the stream gravels sampled, and a placer concentrate obtained. The traverse indicated no radioactivity of any significance above the normal background count. However, a placer concentrate obtained contains 0.18 percent equivalent uranium. The results of the laboratory studies on the radioactivity of the samples are summarized in table 10.

CONCLUSIONS

The results of radioactivity investigations in the Koyukuk-Chandalar region during the summer of 1952 showed that no high-grade radioactive deposits are associated with certain metalliferous lode deposits in the Chandalar mining district which were previously deemed favorable for the occurrence of uranium ores.

Trace amounts of monazite have been found in samples from Little Squaw, Big Squaw, and Tobin Creeks in the Chandalar mining district, but traverses in the area did not locate the bedrock source of the monazite. However, Mertie (1925, p. 263) has suggested the source of the monazite to be some highly acid granitic rock, possibly of pegmatitic character. The results of investigations in the Chandalar mining district indicate that there is little possibility of finding commercial quantities of monazite in the stream gravels.

Traces of uranothorianite have been found in the stream gravels of the middle fork of Big Squaw Creek in the Chandalar mining district. The valley of Big Squaw Creek is about 5 miles long and approximately 1.5 miles from its mouth divides into three branches; a west fork, a middle fork, and an east fork. Stream-gravel samples were taken from each fork, but only the sample from the middle fork contained uranothorianite.

The traverses made in the district include one on a portion of the ridge at the head of the valley of Big Squaw Creek; this traverse, however, detected no anomalous radioactivity. Likewise, no prominent zones of oxidation or alteration were noted in the valley of Big Squaw Creek to denote the presence of sulfide-bearing mineral deposits that could be the source of the uranothorianite and the other metallic minerals occurring in the placer concentrates. Nevertheless, it is highly probable that such a

mineral deposit does exist but may not be very extensive as only trace amounts of the metallic minerals were found. It is thought that these minerals were not deposited by glacial action because the upper valley of Big Squaw Creek does not appear to have been glaciated nor do the tops of the mountains at the head of Big Squaw Creek seem to have been overrun by glaciers as has been pointed out by Mertie (1925).

Thus, the traces of uranothorianite and metallic sulfides in concentrates from the placers of the valley of Big Squaw Creek suggest that this locality may be favorable to prospect for a uraniferous lode deposit. It should be stressed, however, that the discovery of the bedrock source of the uranothorianite at this locality would not necessarily lead to commercial production of uraniferous ores. Such items as size, extent, and grade of a deposit, insofar as they can be determined by surficial investigations, as well as the remoteness and, hence, accessibility of the district are all factors requiring much consideration and study prior to exploration and development.

Uraniferous thorianite has been noted in placer concentrates taken from Gold Bench previous to 1952 (White, 1952, p. 8-12). A placer concentrate obtained during the summer of 1952 contains 0.18 percent equivalent uranium which is due primarily to uranothorianite and, to a lesser extent, to monazite.

Maddren (1913, p. 106) in discussing the origin of the placer gold occurring at Gold Bench, and hence, presumably, of the uranothorianite as well, suggests that the mountains on the south side of the South Fork of the Koyukuk River are the probable source area because the rocks there are known to contain gold. These mountains are drained by tributaries of the South Fork and are formed in part of igneous rocks which may have genetically associated metalliferous lode deposits that could be the source of the radioactive and metallic minerals in the placers at Gold Bench. Very likely, a reconnaissance consisting of the radioactivity testing of concentrates from the placer

gravels of minor creeks and tributaries and slope wash in the mountains south of the South Fork could be used successfully to locate the source of the uranothorianite. Such a reconnaissance technique is similar to that used in 1947 by West and Matzko (Gault, Killeen, West, and others, 1953, p. 21-27) and in 1948 by West (1953) on the eastern part of the Seward Peninsula, Alaska, where soil conditions and sampling problems are similar in nature to those in the vicinity of the South Fork of the Koyukuk River.

LITERATURE CITED

- Gault, H. R., Killeen, P. L., West, W. S., and others, 1953, Reconnaissance for radioactive deposits in the northeastern part of the Seward Peninsula, Alaska, 1945-47 and 1951: U. S. Geol. Survey Circ. 250.
- Maddren, A. G., 1913, The Koyukuk-Chandalar region, Alaska: U. S. Geol. Survey Bull. 532.
- Mertie, J. B., Jr., 1925, Geology and gold placers of the Chandalar district, Alaska: U. S. Geol. Survey Bull. 773-E, p. 215-263.
- West, W. S., 1953, Reconnaissance for radioactive deposits in the Darby Mountains, Seward Peninsula, Alaska, 1948: U. S. Geol. Survey Circ. 300.
- White, M. G., 1952, Radioactivity of selected rocks and placer concentrates from northeastern Alaska: U. S. Geol. Survey Circ. 195.

UNPUBLISHED REPORTS

- Wedow, Helmut, Jr., 1951, Adaptation of portable survey meters for airborne reconnaissance with light planes in Alaska: U. S. Geol. Survey Trace Elements Memo. Rept. 323. (In open files--released through Technical Information Service, U. S. Atomic Energy Commission, Oak Ridge, Tennessee.)
- Wedow, Helmut, Jr., and others, 1951, Interim report on an appraisal of the uranium possibilities of Alaska: U. S. Geol. Survey Trace Elements Memo. Rept. 235. (In open files of the U. S. Geol. Survey.)

CHAPTER D.--FORTY MILE DISTRICT

By John J. Matzko

INTRODUCTION


A sample of fresh fluorite ranging from colorless to blue, green, and purple was received by the Territorial Department of Mines in the summer of 1952 from Mr. George King, Boundary, Alaska. Later, in discussions with Mr. King, the writer learned that the sample came from Mr. George Robinson of Jack Wade on Wade Creek; also, that a Mr. McDonald, who was prospecting in the Fortymile district, had uncovered some fluorite along the highway east of Chicken, Alaska (figs. 1 and 6).

A party consisting of John J. Matzko, geologist, and Fred Freitag, field assistant, investigated these occurrences during the latter part of August 1952. This work was done on behalf of the Division of Raw Materials of the U. S. Atomic Energy Commission.

AREAS INVESTIGATED

Wade Creek

Wade Creek heads against Steel Dome and flows almost due southwest to join Walker Fork. The bedrock in Wade Creek includes several varieties of schist and some thin-bedded ferruginous limestones, all of which are a part of the Birch Creek schist of pre-Cambrian age (Prindle, 1909; Mertie, 1938). Several small bodies of granitic rocks intrude schist in the valley. A basaltic dike is reported to occur near the head of the creek (Prindle, 1909, p. 36).



Base from Eagle quadrangle, Alaska Reconnaissance Topographic Series

SCALE 1:250,000



Mr. Robinson, who had given the fluorite sample to Mr. King for identification, was working a small gold-placer deposit on Wade Creek. Robinson reported that he had found the fluorite in a prospect pit which he had dug on a fourth left tributary above the mouth of Wade Creek (fig. 6) and that the pit had since caved.

A concentrate of stream-gravel (sample 4626, fig. 6) from Wade Creek taken below the mouth of the fluorite-bearing tributary has an equivalent uranium content of 0.001 percent.

Walker Fork

Mr. George King reported to the writer that a Mr. McDonald prospecting in the Fortymile district, Eagle quadrangle, had uncovered some fluorite along the highway east of Chicken. The fluorite monument, which McDonald had set up to mark the location, was found east of Chicken and about 2 miles west of the Walker Fork bridge on the south side of the road (fig. 6). Only a few extra pieces of weathered fluorite besides those of the marker were found. A traverse made with a Geiger counter did not locate any anomalous radioactivity.

A channel sample, 4627, taken across the zone of decomposed rock contains 0.003 percent equivalent uranium as determined in the laboratory. Mineralogic examination of the sample indicates that goethite and hematite are the major constituents; colorless fluorite constitutes about 5 percent of the sample; and iron-stained mica, pyrite, and rutile occur in minor amounts.

Bedrock outcrops alongside the road west of the bridge across Walker Fork and on either side of the sampled fluorite zone consist of fresh-appearing granodioritic rock. Details of the geology of the area have been discussed by Prindle (1909) and Mertie (1938).

CONCLUSIONS

Radioactive minerals occur only in trace amounts in the weathered fluorite zone west of the Walker Fork bridge. The source of the fluorite, which occurs in the gravels of a tributary of Wade Creek, was not determined, and no radioactive minerals were found in a stream sample concentrate taken below the mouth of the tributary.

LITERATURE CITED

- Mertie, J. B., Jr., 1938, Gold placers of the Fortymile, Eagle, and Circle districts, Alaska: U. S. Geol. Survey Bull. 897-C, p. 133-261.
- Prindle, L. M., 1909, The Fortymile quadrangle, Yukon-Tanana region, Alaska: U. S. Geol. Survey Bull. 375.