


(200)
T67r
No. 581

Uranium in carbonaceous rocks in the Townsend and Helena Valleys, Montana

By G. E. Becraft



Trace Elements Investigations Report 581

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

Geology and Mineralogy

This document consists of 24 pages.
Series A

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

URANIUM IN CARBONACEOUS ROCKS IN THE TOWNSEND
AND HELENA VALLEYS, MONTANA *

By

George E. Becraft

April 1956

Trace Elements Investigations Report 581

This preliminary report is distributed
without editorial and technical review
for conformity with official standards
and nomenclature. It is not for public
inspection or quotation.

*This report concerns work done on behalf of the Division
of Raw Materials of the U. S. Atomic Energy Commission.

USGS - TEI- 581

GEOLOGY AND MINERALOGY

<u>Distribution (Series A)</u>	<u>No. of copies</u>
Atomic Energy Commission, Washington	2
Division of Raw Materials, Albuquerque	1
Division of Raw Materials, Austin	1
Division of Raw Materials, Butte	1
Division of Raw Materials, Casper	1
Division of Raw Materials, Denver	1
Division of Raw Materials, Ishpeming	1
Division of Raw Materials, Phoenix	1
Division of Raw Materials, Rapid City	1
Division of Raw Materials, St. George	1
Division of Raw Materials, Salt Lake City	1
Division of Raw Materials, Washington	3
Exploration Division, Grand Junction Operations Office	1
Grand Junction Operations Office	1
Technical Information Extension, Oak Ridge	6
U. S. Geological Survey:	
Fuels Branch, Washington	1
Geochemistry and Petrology Branch, Washington	1
Geophysics Branch, Washington	1
Mineral Classification Branch, Washington	1
Mineral Deposits Branch, Washington	2
P. C. Bateman, Menlo Park	1
A. L. Brokaw, Grand Junction	1
N. M. Denson, Denver	1
V. L. Freeman, College	1
R. L. Griggs, Albuquerque	1
M. R. Klepper, Spokane	1
A. H. Koschmann, Denver	1
W. D. R. Keffer, Laramie	1
L. R. Page, Washington	1
Q. D. Singewald, Beltsville	1
A. E. Weissenborn, Spokane	2
TEPCO, Denver	2
TEPCO, RPS, Washington, (including master)	2
	<u>45</u>

CONTENTS

	Page
Abstract	4
Introduction	4
General geology	7
Geology and uranium deposits of areas examined in detail	8
Area northeast of Winston	8
Geology	8
Uranium deposits	12
Locality A	13
Locality B	13
Locality C	14
Locality D	15
Locality E	15
Locality F	15
Locality G	16
Other localities	16
Area west of Hauser Lake	18
Area in T. 3 N., R. 1 W.	20
Area in T. 2 N., R. 1 W.	20
Origin of the uranium deposits	22
Suggestions for prospecting	23
Literature cited	24

ILLUSTRATIONS

Figure 1. Index map showing the Townsend and Helena Valleys and the locations of figures 2, 3, 4, and 5	6
2. Geologic map showing the radioactivity anomalies northeast of Winston, Broadwater County, Montana	9
3. Geologic map showing radioactivity anomalies west of Hauser Lake in the Helena Valley, Montana	19
4. Radioactivity anomalies in T. 3 N., R. 1 W., near the southern end of the Townsend Valley, Montana	21
5. Radioactivity anomalies in T. 2 N., R. 1 W., near the southern end of the Townsend Valley, Montana	21

TABLE

Table 1. Equivalent uranium and uranium analyses, in percent, from localities shown on figure 2	17
---	----

URANIUM IN CARBONACEOUS ROCKS IN THE TOWNSEND
AND HELENA VALLEYS, MONTANA

By George E. Becraft

ABSTRACT

Uranium-bearing carbonaceous shale and lignite beds are exposed in five areas in the Townsend and Helena Valleys in western Montana. The greatest number of exposures is in an area of several square miles northeast of Winston in the Townsend Valley. The uranium-bearing beds are in the lower part of a Tertiary unit that consists largely of thin-bedded, white to buff, pure and impure tuffs, locally altered to bentonite.

The uranium occurrences, none of which appear to be commercial, have three characteristics in common: (1) they are in and adjacent to carbonaceous shale or lignite interbedded with light-gray or white, fine-grained tuffs and lapilli tuffs, (2) the stratigraphic section in the vicinity of the deposits includes bentonite and partly bentonized tuff, and (3) the distribution of the uranium in the favorable beds is erratic. The uranium was probably leached from the tuffs and lapilli tuffs by meteoric water during bentonization and was concentrated in the carbonaceous shale and lignite.

Similar Tertiary rocks are present in many of the major valleys in Western Montana and probably warrant prospecting for uranium. Areas containing white, fine-grained tuff or lapilli tuff, bentonite, and coal, or carbonaceous shale would be particularly favorable for prospecting.

INTRODUCTION

Uranium occurrences were discovered in Tertiary sedimentary rocks, consisting largely of volcanic debris, northeast of Winston, Mont., in the Townsend Valley early in 1955. Subsequently, uranium was discovered in similar rocks near the southern end of the Townsend Valley and near the eastern end of the Helena Valley.

The Townsend and Helena Valleys are wide basins in western Montana near the headwaters of the Missouri River (fig. 1). U. S. Highway 10 N traverses the length of both valleys. Helena is near the southwestern margin of the Helena Valley, and Townsend is near the center of the Townsend Valley. The Missouri River flows northward through the Townsend Valley, and Prickly Pear Creek flows northward across the Helena Valley and joins the Missouri River near the northeastern margin of the valley.

Earlier work in the Townsend Valley includes a report on the geology and groundwater by Pardee (1925); a report on the geology of the Canyon Ferry quadrangle, which includes the northern end of the Townsend Valley, by Mertie, Fischer, and Hobbs, (1951); and a report by Freeman, Ruppel, and Klepper (in press) which includes detailed maps of most of the western part of the Townsend Valley.

Brief descriptions of the geology of the Helena Valley are included in reports by Knopf, (1913); Pardee and Schrader, (1933); and Lorenz and Swenson, (1951). Hail and Gill, (1953) briefly describe slight radioactivity associated with carbonaceous shales in the Helena Valley.

During May and June 1955, the writer examined several areas containing uranium minerals and radioactivity anomalies in the Tertiary sedimentary rocks in the Townsend and Helena Valleys. Much of this time was spent making a detailed map and study of the uranium occurrences northeast of Winston. Extensive radiometric surveys were made with a scintillation counter in selected areas and with a carborne scintillation counter over larger areas in both valleys. Paul E. Myers assisted the writer with much of the radiometric surveying. This examination is part of a study of the uranium deposits in and near the Boulder batholith that is being carried out by the U. S. Geological Survey on behalf of the Division of Raw Materials of the U. S. Atomic Energy Commission.

The writer wishes to acknowledge the kind permission granted by the ranch owners in the vicinity of Beaver Creek to cross their ranches at a time inconvenient to them.




FIGURE 1.—Index map showing the Townsend and Helena Valleys and the locations of Figures 2, 3, 4 and 5.

GENERAL GEOLOGY

Pardee (1925, p. 1) defines the Townsend Valley as "the widened part of the valley of Missouri River between Canyon Ferry and Toston and a broad lowland west of Toston" It "is about 60 miles long and 600 square miles in extent." In a brief summary of the geology of the area, he states (1925, p. 2-3) that "The rocks consist of two groups separated by a major unconformity. The older group includes a great thickness of quartzite, shale, and limestone that range in age from Precambrian to Cretaceous and have been severely deformed and deeply eroded. Associated with these beds are intrusive and extrusive igneous rocks, chiefly of Cretaceous or early Tertiary age. These rocks compose the mountains and in the valley form the bedrock upon which the younger rocks lie.

"The younger group, which occupies most of the valley, includes beds of Oligocene, Miocene, Pleistocene, and Recent age."

All of the known radioactivity anomalies in the Townsend Valley are in rocks of Oligocene age. Northeast of Winston, the uranium occurrences are in the lower part of the Oligocene rocks, and the anomalies near the southeastern margin of the valley are in similar and probably correlative rocks.

Helena Valley is another wide basin, separated from Townsend Valley by a broad, low divide northwest of Winston. The valley extends northwestward from the divide about 20 miles and is about 10 miles wide at its widest part. The geology is similar to the geology of the Townsend Valley except that the Quaternary deposits are much more extensive, and as a result the Tertiary rocks in the valley are not exposed except in a few small areas. A few weak radioactivity anomalies occur in rocks similar to the Oligocene rocks northeast of Winston.

GEOLOGY AND URANIUM DEPOSITS OF AREAS EXAMINED IN DETAIL

Area northeast of Winston

Geology

The area northeast of Winston (fig. 2) is near Canyon Ferry Lake in the Canyon Ferry quadrangle and extends north and south of Beaver Creek. Mertie divided the Tertiary rocks in the Canyon Ferry quadrangle into 5 units (Mertie, Fischer, and Hobbs, 1951, p. 30). Three of these units are exposed in the area of figure 2.


Unit 1 consists of conglomerate, red shale, interbedded dark-gray, partly bentonized tuff, and bentonite. It is poorly exposed in the area. Mertie (Mertie, Fischer, and Hobbs, 1951, p. 31-32) describes unit 1 as follows:

"The boundaries of this unit are not well-defined. The beds include some conglomerate, which disintegrates by weathering so that the resulting material is not easy to distinguish from the terrace gravels. Along its eastern boundary, the sediments grade into those of unit 2, so that this boundary is a matter of interpretation. The boundaries shown are therefore necessarily somewhat inexact.

"The sediments of unit 1 are nowhere continuously exposed. A few outcroppings show that the unit consists of conglomerate and red shale, the latter including some bentonitic horizons, but the proportions of conglomerate to shale cannot be determined. At one site along the north side of Beaver Creek, the conglomerate is in place, and consists of well-rounded gravels, cobbles, and boulders, as much as 3 feet in diameter, with an average size of about 6 inches. They are mainly basic lava, with a small amount of quartzite. Some of the cobbles are exfoliated by weathering. The matrix consists of a yellow to reddish grit, made up of small well-rounded grains of limestone, and a small amount of other kinds of country rock. About a mile north of Beaver Creek, the red shale is exposed in an elbow-shaped bend of a small gulch. These beds are banded, tuffaceous, and bentonitic, and are composed mainly of grains as much as 0.1 millimeter in size, averaging perhaps 0.025 millimeter. The reddish-brown grains, originally glass, are altered to montmorillonite and kaolin. Angular grains of quartz, sericite, chloritized biotite, and altered iron ores are the other constituents. In the next gulch to the south, about three-fourths of a mile north of Beaver Creek, 10 feet of reddish-brown conglomerate and grit are exposed, the matrix of which is composed mainly of grains of limestone."

The author has slightly modified the contact between units 1 and 2 as shown by Mertie, Fischer, and Hobbs (1951, pl. 2). One modification is in the SE 1/4 sec. 18 (fig. 2) where Mertie, Fischer, and Hobbs included in unit 2 several thick beds of conglomerate, which are made up of cobbles and pebbles of pre-Tertiary rocks, and interbedded bentonite and partly bentonized, dark-gray tuffs. These rocks are typical of unit 1 and are therefore included in unit 1 by the author. Another modification was made near

R. 1 E.



EXPLANATION

- Qal Alluvium
- Qoq Old gravels and sand
- Tu3 Unit 3
Massively bedded, buff, impure tuffs
- Tu2 Unit 2, showing areas of bentonite
Thin-bedded, white to buff, pure and impure tuffs including areas of bentonite and thin beds of dark brown to black carbonaceous shale.
- Tu1 Unit 1, showing areas of bentonite
Conglomerate, red shale, and dark gray, impure tuffs in places bentonized.

Quaternary
TERTIARY

Oligocene

- Contact, dashed where approximately located
- Concealed contact
- |--- Fault, showing dip, dotted where concealed
- |--- Vertical fault
- 13 \ Strike and dip of beds
- Horizontal bedding
- A ● Radioactivity anomalies referred to in report

SCALE 1:24000
1000 0 1000 FEET
CONTOUR INTERVAL 40 FEET

Figure 2 - Geologic Map showing radioactivity anomalies northeast of Winston, Broadwater County, Montana

Geology by C.E. Beckett, 1955

46° 30'

113° 35'

the southern margin of the Canyon Ferry quadrangle where several beds of reddish-brown conglomerate interbedded with bentonite and partly bentonized tuff are exposed along an abandoned road half a mile from the lake. These beds were included in unit 2 by Mertie, Fischer, and Hobbs but are included in unit 1 by the present author.

In general the beds of unit 1 strike northwest and dip 15° - 25° NE.; however, along the abandoned road near the south margin of figure 2, the beds strike north and dip about 40° E. South of the road for a short distance, the beds strike northeast and dip 20° - 30° SE. These differences in attitudes may indicate a concealed east-trending fault north of the abandoned road; an anticline plunging to the east; or a topographic high in the older rocks at the time of deposition of the Tertiary rocks; but, because of lack of exposures, the reason for the differences could not be determined.

Rocks of unit 2 underlie most of the area of figure 2 and are exposed north of Beaver Creek for a distance of 2-3/4 miles from the contact of unit 1 to Canyon Ferry Lake. Previous workers separated unit 2 into three parts (Mertie, Fischer, and Hobbs, 1951, p. 33-35), but in this report the lower two-thirds is considered as a single part which differs markedly from the upper one-third. The original division was based on the presence of more and thicker beds of bentonite in the lower third than in the middle third of the unit. However, the amount of bentonite present in the section differs considerably more along the strike of the unit than across the strike. Although most of the tuffs in the lower two-thirds of unit 2 are bentonized to a slight degree, the almost completely bentonized tuffs--referred to in this report as bentonites--are more common near the south margin of figure 2 and become less common northwestward along the strike of unit 2. In section 18 only a few thin beds are bentonized to the degree common south of Beaver Creek.

The lower part of unit 2 is largely unaltered or only slightly altered tuff and lapilli tuff, which make up about 60 percent of the section, and bentonite, which makes up about 20 percent. The remainder of the section is tuffaceous sandstone, conglomerate, and breccia and includes many thin beds of carbonaceous shale and chert. The tuffs, lapilli tuffs, and tuffaceous sandstones are predominantly white or light gray but also include gray, cream, yellow, and buff beds. The carbonaceous shale beds range from light brown to black. The lithology commonly changes along strike which makes correlation of sections very difficult.

The tuffs and lapilli tuffs consist of angular fragments of quartz, plagioclase, orthoclase, and chert and angular to subrounded fragments of pumice in a matrix of fine glassy material. The pumice fragments and matrix are largely kaolinized or bentonized. The mineral fragments make up from 10 to 50 percent of the rock.

The sandstones consist largely of subangular to subrounded grains of quartz, plagioclase, orthoclase, and chert and commonly contain rounded grains of pumice largely altered to clay. In the coarser sandstones occasional rounded grains of rhyolite are present.

The carbonaceous shales are from a few inches to 18 inches thick. Many of the shale beds contain only a small percentage of carbonaceous material, but some consist almost entirely of carbonized plant fragments.

The chert beds, which may represent tuff beds replaced by silica, are mostly less than a foot thick and are associated with bentonite or partly bentonized tuff. In places the chert is opalized.

The upper part of unit 2 differs markedly from the lower part. It consists largely of buff to gray, fine-grained tuff, very little of which shows any bentonization. Beds of limestone are common and no beds of carbonaceous shale were recognized. Mertie, Fischer, and Hobbs (1951, p. 35) report that two beds of diatomaceous earth occur in the upper part of the sequence.

In general the beds of unit 2 strike northwest and dip northeast. The dip is usually greater in the lower part of the unit than in the upper part. This difference in average dip may indicate deposition in a subsiding basin. Following deposition of the lower beds but before deposition of the upper beds, the basin may have subsided differentially, the greatest subsidence being east of the mapped area.

In places gentle folds are superimposed on the regional structure causing local differences in the attitudes of the beds.

North of Beaver Creek, several normal faults cut the lower part of unit 2. Commonly the east block moved down but in a few places the west block apparently moved down. In the southeast corner of section 18, the contact between unit 1 and unit 2 is offset by a normal fault that strikes about north and dips 60°E. Most of the apparent offset, about 1,200 feet, is probably the result of erosion rather than strike-slip displacement. Banded chalcedony was deposited along some of the faults probably by meteoric water because no evidence of hydrothermal alteration of the rocks along the faults was observed.

In section 7 in the northwestern corner of figure 2, tuffs that resemble those mapped as unit 3 near Canyon Ferry (Mertie, Fischer, and Hobbs, 1951) are in fault contact with unit 2 (fig. 2). The tuffs are fine-grained, buff colored, and massive.

The Quaternary deposits shown on figure 2 are of two ages-- gravel and sand deposits, principally on terraces cut on the tilted Tertiary rocks, and recent alluvium in the stream valleys. South of Beaver Creek the older deposits consist largely of volcanic rocks probably derived from the Elkhorn Mountains several miles west of the area. North of Beaver Creek, debris from the Spokane Hills, immediately west of the area, is mixed with the gravel. These older deposits are in places at least 10 feet thick and cover very broad areas (fig. 2). Because of this gravel cover and the alluvium in the valleys, several beds of radioactive carbonaceous shale could be traced for only short distances.

Uranium deposits

Radioactivity anomalies were detected at about 20 localities northeast of Winston (figs. 1 and 2). All of the anomalies are in or adjacent to carbonaceous shales in the lower part of Tertiary unit 2. The shale beds are lenticular and no single bed could be traced for more than a few hundred feet. Secondary uranium minerals were identified from five of the localities. No primary uranium minerals were recognized. Almost all of the carbonaceous shale in unit 2 is slightly radioactive, but only the localities where the radioactivity is several times background are shown on figure 2. Several beds of white tuff and white lapilli tuff were also slightly more radioactive than the surrounding rocks. Additional radioactivity anomalies undoubtedly exist under the extensive terrace gravel deposits but could not be detected by scintillation counter traverses because of the depth of the gravel cover.

The radioactivity anomalies shown on figure 2 are described below.

Locality A. -- Locality A is in the SE 1/4 sec. 20 near the top of the steep north slope of the broad terrace north of Beaver Creek. Uranium occurs in a poorly indurated, carbonaceous shale and in a fine-grained, partly bentonized, gray tuff overlying the shale. Locally, lenses of sandstone in the overlying tuff also contain uranium. The shale strikes N. 15° W. and dips 20° NE, and ranges in thickness from less than 6 inches to 18 inches. The bed is exposed for a distance of about 200 feet in a bulldozed cut. The radioactivity is not distributed evenly throughout the shale and is strongest over a strike distance of about 30 feet near the center of the exposed area. Results of drilling on the terrace indicate that the uraniferous bed extends at least a few hundred feet south of the exposed area.

A yellowish-green, micaceous mineral that coats fractures in the in the shale and overlying tuff and replaces plant fragments in the shale was identified optically, spectrographically, and by X-ray diffraction pattern as metatorbernite by Jerome Stone of the U. S. Geological Survey. The mineral fluoresces bright green. No other uranium minerals were detected at this locality.

An attempt to strip mine the uranium at this locality was made in June 1955. Because of the dip of the uranium-bearing bed, a considerable amount of overburden had to be removed. The overburden was stripped to the carbonaceous shale which at this point was about 12 inches thick. Care was taken to mine only the carbonaceous shale.

Locality B. -- Locality B is on the west side of a small knoll in the northeast corner of section 20 (fig. 2). The uranium is in a 12-inch thick bed of light reddish-brown carbonaceous shale and an underlying light greenish-gray chert bed about 6 inches thick. Overlying the carbonaceous shale is a partly bentonized lapilli tuff several feet thick. Underlying the chert bed is a 6-foot zone of alternating chert and white tuffs. Underlying this sequence is a thick, partly bentonized tuff. The beds strike N. 18° W. and dip 22° NE.

The carbonaceous shale is exposed for a strike length of about 200 feet. It lenses out toward the southeast and could not be located on the slope across a small valley to the northwest. The bed is cut off by a northeast-trending fault about 150 feet down dip from the highest exposure.

Carnotite, identified by Jerome Stone, U. S. Geological Survey, coats fractures and has partly replaced plant fragments in the carbonaceous shale. In several places carnotite coats fractures in the underlying chert also. A selected sample of the carbonaceous shale assayed 0.28 percent eU, 0.70 percent U, and 0.2 percent V /. A selected sample of carnotite-coated chert assayed 0.019 percent eU, 0.022 percent U, and 0.01 percent V. A bulk sample from a small ore pile assayed 0.062 percent eU. The bulk sample was ground and separated into size fractions and each fraction was then assayed for equivalent uranium with the following results:

+ 20 mesh	0.028 percent eU
+ 40 mesh	0.032 percent eU
+ 60 mesh	0.044 percent eU
+ 80 mesh	0.072 percent eU
+ 100 mesh	0.064 percent eU
+ 140 mesh	0.074 percent eU
- 140 mesh	0.11 percent eU

By weight, one-half of the sample is finer than 60 mesh. These data indicate that the bulk of the uranium is in the finer fractions, and the uranium might be concentrated by grinding and screening.

Locality C. -- Locality C is on a low, rounded spur near the bottom of a broad valley in section 18 (fig. 2). The uranium is in a brown carbonaceous shale bed about eight inches thick. Overlying this shale is a thin soil cover. Underlying the shale is a 4-inch bed of white, fine-grained tuff which is underlain by another 8-inch carbonaceous shale bed. These carbonaceous shale beds are more indurated than those at Localities A and B.

The beds are exposed for a total distance of about 50 feet in two shallow bulldozer cuts and elsewhere are covered by colluvium. The attitude of the beds is N. 80° W., 7° NE. Only the upper shale bed is strongly radioactive, but locally the lower shale bed has slight radioactivity. A selected sample of the upper shale assayed 0.051 percent eU, 0.10 percent U, and 0.03 percent V. No uranium minerals were detected.

/ All vanadium analyses in this report were made by K. E. Valentine, U. S. Geological Survey.

Locality D. -- Locality D is an outcrop at the northwest end of a low mound near the center of a broad valley in section 18 (fig. 2). The uranium is in black, well-indurated carbonaceous shale about 6 inches thick overlying a 12-inch dark-gray chert bed and underlying a lighter-gray chert bed at least 8 inches thick. The beds strike about N. 40° W. and dip about 3° NE. and are exposed for a distance of about 30 feet. Carnotite occurs in the black shale and coats fractures in the underlying chert. A selected sample of the black shale assayed 0.11 percent eU, 0.20 percent U, and 0.3 percent V.

About 500 feet southeast of this outcrop, a radioactive greenish-gray, partly bentonized lapilli tuff bed about 4 feet thick is exposed in a bulldozer cut. The bed is overlain and underlain by gray, partly bentonized tuff. The bed is exposed for a distance of about 100 feet and has an attitude of N. 20° W., 25° NE. The radioactive material is distributed erratically in the bed. A selected sample of the radioactive material assayed 0.069 percent eU.

Locality E. -- Locality E is in a shallow bulldozer cut on the north side of a small ridge near the center of the western margin of section 20 (fig. 2). The uranium is in a carbonaceous shale exposed over an area of only a few square feet near the bottom of the south wall of the bulldozer cut. The rock surrounding the carbonaceous material is partly bentonized tuff and bentonite. In the vicinity of the anomaly the rocks are cut by several small, highly irregular faults and about 40 feet east of the anomaly by a larger fault. The details of the structure are difficult to determine because of poor exposures and flowage of the bentonite. No uranium minerals were recognized, but a selected sample of the highly carbonaceous material assayed 0.12 percent eU, 0.23 percent U, and 0.02 percent V. A composite sample of the bentonite in the vicinity contained 0.002 percent eU but no uranium.

Locality F. -- Locality F is about 250 feet north of Locality E in section 20 (fig. 2). A sequence about 6 feet thick of carbonaceous beds and fine-grained tuff beds is exposed in a road cut. The carbonaceous beds range from lignite to shales containing only a small amount of plant fragments. The beds are from 2 inches to 12 inches thick and strike N. 15° W. and dip 25° NE. The sequence is overlain by bentonite and underlain by partly bentonized tuff. A bentonitic mud flow covers the beds to the west

of the road cut, and the beds are cut off near the east end of the cut by a fault which strikes N. 10° W. and dips 62° SW. East of the fault the rock is white sandy tuff made up of 30 percent to 50 percent of rounded pumice fragments and about 10 percent rounded grains of chert in a fine-grained glassy matrix.

Several of the carbonaceous beds are radioactive. A sample from a 2-inch bed of lignite about 12 inches above the base of the sequence assayed 0.015 percent eU and 0.022 percent U. A sample selected from the most radioactive bed in the sequence--a 6-inch bed of reddish-brown carbonaceous shale about 4 feet above the base--assayed 0.051 percent eU and 0.021 percent U. A bulk sample of all the carbonaceous beds assayed 0.008 percent eU and 0.009 percent U.

Locality G. -- Locality G is a large area south of Beaver Creek in section 28 (fig. 2). The rock in this area is predominantly bentonite with smaller amounts of partly bentonized tuffs and lapilli tuffs. Three beds of carbonaceous shale in the area of bentonite are radioactive. The lowest shale bed in the section is 3 inches thick and is underlain by a 6-inch chert layer which is underlain by bentonite. The bentonite appears to have originally been a lapilli tuff consisting largely of ash and rounded fragments of pumice about a quarter of an inch in diameter. Overlying the shale is a sandy tuff, about 2 feet thick, that consists almost entirely of rounded pumice fragments altered to bentonite in a matrix of glassy material largely altered to clay. About 50 feet higher in the section is a 4-inch carbonaceous shale and about 2 feet higher is another carbonaceous shale about 2 inches thick. The beds between the shales have been almost completely altered to bentonite. These beds strike N. 30° - 35° W. and dip 25° - 40° NE. The carbonaceous beds are exposed for a distance of about 250 feet. Metatorbernite is locally along fractures in the lowest shale bed. Samples selected from two of the beds assayed 0.36 and 0.054 percent eU.

Nonradioactive, thoroughly silicified plant fragments and logs are associated with the bentonite at several places.

Other localities. -- The other localities shown on figure 2 are similar to those described above, but the uranium content is generally considerably lower; therefore the information on these localities is presented on table 1 along with the analytical data given above for comparison.

Table 1. --Equivalent uranium and uranium analyses, in percent,
from localities shown on figure 2.

Locality	eU	U	Description	Thickness of bed (inches)
B	0.28*	0.70	carbonaceous shale containing carnotite	12
B	.019*	.022	chert with carnotite coated fractures	6
B	.062	***	carbonaceous shale and chert	<u>1/</u>
C	.051*	.10	carbonaceous shale	8
C	.23	***	carbonaceous shale	--
C	.24	***	carbonaceous shale	--
D	.11*	.20	carbonaceous shale containing carnotite	6
D	.069	***	partly bentonized lapilli tuff	12
D	.034	***	carbonaceous shale containing carnotite	6
E	.12*	.23	carbonaceous shale	--
E	.20	***	carbonaceous shale	--
E	.002**	0	bentonite	<u>1/</u>
F	.015**	.022	carbonaceous shale	2
F	.051**	.021	carbonaceous shale	6
F	.008**	.009	carbonaceous shale	<u>1/</u>
G	.36	***	carbonaceous shale containing metatorbenite	3
G	.054	***	carbonaceous shale	4
H	.007**	.004	partly bentonized fine-grained tuff	12
H	.013**	.009	fine-grained tuff	--
J	.004**	0	tuffaceous sandstone	36
K	.004**	0	coarse, partly bentonized sandstone	--
L	.013**	.019	silicified wood	<u>2/</u>
M	.038**	.071	brown carbonaceous shale	7
N	.005**	.003	white, coarse, sandy tuff	48
P	.008**	.006	gray tuff with plant fragments	48
Q	.009**	.004	tuffaceous sandstone	24
Q	.018**	.036	carbonaceous shale containing carnotite	18
Q	.001**	0	light-gray chert	12
R	.007**	.005	carbonaceous shale	8
S	.011**	.012	carbonaceous shale	6
T	.009	***	brown chert	4
U	.010	***	carbonaceous shale	6

* Analysts: Benjamin A. McCall and Roosevelt Moore

** Analysts: C. Angelo and M. Finch

*** Not analyzed for uranium

1/ bulk sample


2/ fossil tree trunk fragment

Area west of Hauser Lake


The area shown on figure 3 is on the drainage divide between Hauser Lake and Lake Helena in the Helena Valley (fig. 1). Most of the area is covered by Quaternary gravel and soil, much of which is cultivated. Tertiary rocks are exposed in three small areas (fig. 3). The westernmost exposure of Tertiary rocks in section 29 (fig. 3) consists of conglomerate, tuffaceous conglomerate, tuffaceous sandstone, and bentonite. The conglomerate is made up of pebbles and cobbles of dark volcanic rock, quartzite, and limestone. The rock is not well indurated and tends to crop out only as layers of loose gravel. Most of the conglomerate beds are broadly lenticular and crossbedded. The tuffaceous conglomerate consists of pebbles similar to those in the conglomerate in a matrix of ash now largely altered to clay. The amount of pebbles differs considerably from one bed to another. The conglomerate and tuffaceous conglomerate make up about one-half of the nonbentonitic rocks in the section. The remainder are light-gray, tuffaceous sandstone, which consists of rounded grains of chert and fragments of pumice in a glassy matrix now largely altered to clay, and gray, partly bentonized tuffs. The bentonite is in beds from 4 feet to 12 feet thick. It appears to have been formed from alteration of fine-grained tuff. The bentonite is largely gray but contains a few beds that are purplish-brown as a result of carbonaceous material in the beds. The attitude of the beds in this vicinity is N. 45° W., 8° SW.

In the vicinity of locality D (fig. 3), the section contains less nonvolcanic material than it does farther north. About 20 percent of the section is conglomerate or sandy conglomerate, about 30 percent is bentonite, and the remainder is largely partly bentonized tuff and lapilli tuff. The rocks are well bedded, and most beds are less than one foot thick. Four beds of purplish-brown carbonaceous shale ranging in thickness from one inch to four inches are interbedded with light-gray, partly bentonized tuff near the top of the exposed section. All of the carbonaceous shale is slightly radioactive. A selected sample from the lowest bed assayed 0.011 percent eU.

In the vicinity of locality C (fig. 3), the rocks are predominantly light-gray sandstone and greenish-gray bentonite. Some beds contain many rounded pebbles of older rocks, and a few broadly lenticular beds of conglomerate are interbedded with both the bentonite and the sandstone. Most of the sandstones and conglomerates are crossbedded. A one-inch bed of purplish-gray carbonaceous shale at locality C is slightly radioactive, but it is not as radioactive as the shale at locality D.



SCALE 1:24,000



EXPLANATION

- Qq
Quaternary
Undivided alluvium and older gravel
- TUN
Tertiary
Undivided tuff, sandstone, conglomerate,
bentonite and carbonaceous shale
- Contact
- ∟∟
strike and dip of beds
- B
Radioactivity anomaly

FIGURE 3.—Geologic map showing radioactivity anomalies west of Hauser Lake in the Helena Valley, Montana.

The Tertiary rocks in the vicinity of localities A and B (fig. 3) are partly bentonized fine-grained tuff and bentonite. Interbedded with these are a few beds of light-gray, partly bentonized lapilli tuff. The lapilli are subrounded fragments about 1/4 inch in diameter of light-gray, fine-grained tuff.

At locality B are four slightly radioactive beds of light purplish-gray thin-bedded carbonaceous shale from 3 inches to 1 foot thick. Two thin, slightly radioactive beds of carbonaceous shale crop out at locality A. The shale at localities A and B is less radioactive than the shale at locality C.

Area in T. 3 N., R. 1 W.

The area shown in figure 4 is near the south end of the Townsend Valley (fig. 1). Most of the area is under cultivation and outcrops are poor. At locality A (fig. 4), a four-foot bed of white lapilli tuff is exposed on the southwest side of a small valley. The rock consists of large rounded pumice lapilli in a fine-grained matrix. It is crossbedded and overlies bentonite. A sample across three feet of tuff assayed 0.004 percent eU and 0.002 percent U. A similar, white, slightly radioactive lapillituff is exposed along a low bluff at locality C.

At locality B, a 5-inch bed of light purplish-gray carbonaceous shale is interbedded with gray, partly bentonized tuff. Locally, the shale is partly replaced by chert. A selected sample of the carbonaceous shale assayed 0.006 percent eU and 0.018 percent U. At locality D, a radioactivity anomaly was detected with a carbone scintillation counter near the base of a small knoll. Outcrops are poor but float indicates that the knoll is capped by a bed of white lapilli tuff similar to that at locality A. The source of radioactivity was not found.

Area in T. 2N., R. 1 W.

The area shown in figure 5 is about 5 miles south of the area of figure 4 at the southern end of the Townsend Valley (fig. 1). Almost all of the area is under cultivation and outcrops are extremely poor. The large amounts of bentonite in the soil and on the graded roads and the few outcrops of tuffaceous rocks indicate that the area is probably entirely underlain by Tertiary rocks similar to those northeast of Winston.




FIGURE 4.—Radioactivity anomalies in T.3N., R.1W. near the southern end of the Townsend Valley, Montana.




FIGURE 5.—Radioactivity anomalies in T.2N., R.1W. near the southern end of the Townsend Valley, Montana

At locality A (fig. 5), a slightly radioactive bed of reddish-brown carbonaceous shale 10 inches thick is exposed on both sides of a graded country road. The carbonaceous shale is interbedded with partly bentonized, fine-grained tuff, sandstone, and bentonite. About 6 feet above the shale is a thin bed of limestone.

At locality B, a slight radioactivity anomaly was detected with a carborne scintillation counter. The source of the radioactivity could not be located definitely, but the soil contains a small amount of carbonaceous shale which might be the source.

ORIGIN OF THE URANIUM DEPOSITS

In the Townsend and Helena Valleys, the uranium deposits have several characteristics in common: (1) they are in and adjacent to carbonaceous shale or lignite interbedded with light-gray or white fine-grained tuffs and lapilli tuffs, (2) the stratigraphic section in the vicinity of the deposits includes bentonite and partly bentonized tuff, (3) the distribution of the uranium in the favorable beds is erratic. The source of the uranium was probably the light-gray and white fine-grained tuffs and lapilli tuffs, samples of which contain 0.002 percent to 0.009 percent uranium. A bulk sample of the bentonite, which was formed by alteration largely from similar tuffs and lapilli tuffs, contained no uranium, and no radioactivity was detected in the bentonite during radiometric traversing. These data suggest an epigenetic origin for most of the uranium in the carbonaceous shale and lignite. The uranium was probably leached from the tuffs and lapilli tuffs by meteoric water during bentonization and was concentrated in the carbonaceous shale and lignite, essentially in the manner suggested by Denson (1952) for other deposits of uranium in coal, lignite, and carbonaceous shale in the Rocky Mountain region.

Faulting is common in the Tertiary rocks, and northeast of Winston the development of banded chalcedony along some faults indicates that meteoric water that had leached silica from the surrounding rocks moved along the fault. However, no direct relationship of the uranium deposits to faults was found, nor was any other structural control apparent.

SUGGESTIONS FOR PROSPECTING

Tertiary rocks similar to the rocks described in this report are widespread in many of the major valleys in western Montana. W. J. Hall, Jr., and J. R. Gill (1953) examined many of the known occurrences of coal, lignite, and carbonaceous shale in these valleys, but large areas of Tertiary rocks are still unexplored. Areas of white, fine-grained tuff or lapilli tuff, areas of bentonite, and areas containing coal, lignite, or carbonaceous shale warrant detailed prospecting with a scintillation counter. Areas of white rocks or bentonite can be detected easily on aerial photographs or from the air, thus actual prospecting should be preceded by aerial reconnaissance or a careful study of aerial photographs. Then traverses using carborne scintillation counters could be planned for the most promising areas and traverses on foot could be made where necessary.

LITERATURE CITED

- Denson, N. M., 1952, Summary of uranium-bearing coal, lignite, and carbonaceous shale investigations in the Rocky Mountain region during 1951, with description of deposits by G. O. Bachman, J. R. Gill, W. J. Hail, Jr., J. D. Love, Harold Masursky, N. M. Denson, G. W. Moore, G. N. Pippingos, J. D. Vine, and H. D. Zeller: U. S. Geol. Survey TEM-341A, 44 p., issued by U. S. Atomic Energy Comm. Tech. Inf. Service, Oak Ridge, Tenn.
- Freeman, V. L., Ruppel, E. T., and Klepper, M. R., in preparation, Geology of part of the Townsend Valley, Broadwater and Jefferson Counties, Montana: U. S. Geol. Survey Bull. 1054.
- Hail, W. J., Jr., and Gill, J. R., 1953, Results of reconnaissance for uraniferous coal, lignite, and carbonaceous shale in western Montana: U. S. Geol. Survey Circ. 251, 9 p.
- Knopf, Adolph, 1913, Ore deposits of the Helena mining region, Montana: U. S. Geol. Survey Bull. 527, 143 p.
- Lorenz, H. W., and Swenson, F. A., 1951, Geology and ground-water resources of the Helena Valley, Montana: U. S. Geol. Survey Circ. 83, 68 p.
- Mertie, J. B., Jr., Fischer, R. P., and Hobbs, S. W., 1951, Geology of the Canyon Ferry quadrangle, Montana: U. S. Geol. Survey Bull. 972, 97 p.
- Pardee, J. T. 1925, Geology and ground-water resources of Townsend Valley, Montana: U. S. Geol. Survey Water Supply Paper 539, 61 p.
- Pardee, J. T., and Schrader, F. C., 1933, Metalliferous deposits of the greater Helena Mining region, Montana: U. S. Geol. Survey Bull. 842, 318 p.