

(200)
T67N

UNITED STATES GEOLOGICAL SURVEY


TEI-144

**HYDROTHERMAL URANOTHORITE IN FLUORITE
BRECCIAS FROM THE BLUE JAY MINE, JAMESTOWN,
BOULDER COUNTY, COLORADO**

By
George Phair
Kiyoko Onoda

This preliminary report is released without editorial and technical review for conformity with official standards and nomenclature, to make the information available to interested organizations and to stimulate the search for uranium deposits.

March 1951



29563

Prepared by the Geological Survey for the
UNITED STATES ATOMIC ENERGY COMMISSION
Technical Information Service, Oak Ridge, Tennessee

GEOLOGY AND MINERALOGY

**Reproduced direct from copy
as submitted to this office.**

HYDROTHERMAL URANOTHORITE IN FLUORITE BRECCIAS FROM THE
BLUE JAY MINE, JAMESTOWN, BOULDER COUNTY, COLORADO

by

George Phair and Kiyoko Onoda

ABSTRACT

In the course of thin-section and heavy-mineral studies of fluorite breccias from Jamestown, Colo., the mineral thorite, variety uranothorite, was identified on the basis of chemical, spectrographic, optical, and X-ray data. This uranothorite is compared with uranothorite from South Westland, New Zealand, described by Hutton (1950), and with similar material from an Alaskan placer sample studied by the writers. These uranothorites are in turn contrasted with the uranothorites described in the older literature all of which represent hydrated material. Thorium halos in fluorite are described, probably for the first time. The uranothorite associated with the fluorite breccias is clearly of hydrothermal origin--unlike all previously described thorites and uranothorites with one very doubtful exception.

INTRODUCTION

The Blue Jay mine is one of the larger fluorite producers in the Jamestown district, Boulder County, Colo. G. Phair visited the second level of this mine in September 1949, guided by R. U. King and H. C. Granger of the U. S. Geological Survey's Denver office, and collected some 30 pounds of the breccia ore for mineralogic and chemical studies of the radioactive constituents. A preliminary binocular study of the heavy-mineral contents of this ore revealed scattered grains of an unknown glassy green mineral. Qualitative chemical and spectrographic tests showed that the mineral resembled uranothorite in chemical composition. Although the uranothorite (?) accounted for only a small part of the total radioactivity of this ore, its occurrence in a mineral assemblage which is definitely of hydrothermal origin seemed unique. Largely for this reason the mineral was considered of sufficient interest to justify the numerous separations and the tedious hand-picking of grains for its positive identification. Moreover, the fact that a green mineral had been found in a placer sample from the Ruby district, Alaska, emphasized the need for the specific determination. This mineral had been identified only tentatively as thorite, variety uranothorite, by Onoda because its physical properties departed widely from those of the dozen or so thorites described in the literature prior to 1950.

Acknowledgments

This investigation is part of the program on radioactive raw materials undertaken by the U. S. Geological Survey on behalf of the U. S. Atomic Energy Commission. We are indebted to J. C. Rabbitt, E. S. Larsen, Jr., and T. Botinelly, all of the Geological Survey, for critical review of the manuscript, and to other members of the staff, particularly to J. Berman and E. Dwornik, for helpful discussions of phases of the "thorite" problem of mutual interest. The many chemists and spectrographers of the Survey who obtained the analytical data upon which this report is largely based are cited in the text. We should like also to thank Max White of the Geological Survey for the use of mineralogic material and Professor Clifford Frondel of Harvard University for the reference to Gordon's description of doubtful uranothorite from Llallagua, Bolivia.

METHODS OF SEPARATION

To obtain enough uranothorite for analyses, the parts of the fluorite breccia selected for high radioactivity were crushed to -120, +200 mesh and the heavy minerals separated in methylene iodide. These heavy minerals consisted chiefly of deep-purple fluorite containing heavy impurities; impure carbonates, pyrite, galena, sphalerite, iron oxides, uraninite, and uranothorite. The light fractions consisted chiefly of light-purple fluorite, quartz, carbonate, clay minerals, and altered feldspar, and in bulk samples the light fractions showed negligible radioactivity. Some slight additional concentration was obtained by passing the heavy-mineral fractions through the Frantz

isodynamic separator; the uranothorite tended to accumulate in the moderately magnetic fraction. The desired material was hand-picked from these concentrates under the binocular microscope, and this sample, in turn, was hand-picked a second time for reasons to be described.

IDENTIFICATION OF THE URANOTHORITE

Preliminary optical studies had shown that the green mineral in the fluorite breccias, like that in the Alaskan placer sample, was completely metamict (isotropic) with $n = 1.84$. The specific gravity of several crystals of the Alaskan mineral determined on the Berman microbalance was found to be 6.5 ± 0.2 . It was impossible to reconcile the green color, the high refractive index, and the high density of these minerals with the properties of thorite and uranothorite reported in the literature prior to 1950 ($n = 1.68$ to 1.72 , sp. gr. 4.1 to 5.2). Nevertheless both unknown minerals had the typical thorite form, tetragonal with the prism (110) terminated by the pyramid (111). Like most completely metamict minerals, prior to ignition they gave the X-ray pattern of an amorphous solid. Unlike many completely metamict minerals the isotropic thorites do not revert to their original crystalline condition after being heated. Instead, as noted by Robinson (1950), uranothorites after ignition at 800°C . for half an hour in an inert atmosphere give an anomalous face-centered cubic powder pattern; neither of the two unknown green minerals was an exception to this rule. According to J. Berman of the Geological

Survey, who is at present engaged in X-ray diffraction studies of the different members of the thorite group, our uranothorites after ignition give powder patterns nearly identical to that of a thorianite having $a_0 = 5.60 \text{ \AA}$ and similar to that given by synthetic ThO_2 prepared by igniting $\text{Th}(\text{NO}_3)_4$ in a stream of nitrogen. The patterns of heated thorianite and of our heated uranothorites are nearly identical with respect to the relative intensities and spacings of the major lines; they differ in only one particular, namely, several weak lines present in the thorianite pattern are absent from the less intense uranothorite pattern.

The preliminary qualitative spectrographic analyses were made on a few nearly pure grains of each of the two green minerals. The additional separations yielded, after a single hand-picking, a sample of the Jamestown mineral large enough for the semiquantitative work planned, but because of mineral intergrowths, its over-all purity was lower than was thought desirable. In contrast to the Alaskan mineral, pure samples of which could be obtained with little difficulty, the hand-picked Jamestown material consisted largely of grains of the green mineral more or less intergrown with a brown alteration product of much lower refractive index. Grains containing other mineral intergrowths were less abundant but common. The bulk sample was subdivided by a second hand-picking as follows:

1. Mixed green and brown phases free of other impurities, about 30 mg
2. Green mineral alone plus impurities, about 5 mg
3. Brown mineral alone plus impurities, about 5 mg

All mixed green and brown grains containing other impurities were discarded.

The small size of the samples of the separate brown and green phases (2 and 3 above) permitted only a qualitative spectrographic analysis of each plus a quantitative chemical analysis for uranium. The results of these analyses showed that both the green mineral and its brown alteration product were thorium minerals of similar composition (table 1). The presence of much intergrown sphalerite in both samples accounts for: (1) the high zinc content, (2) the high refractive index (over 2.00) of some of the intergrown material, and (3) the weak sphalerite powder pattern given by the less pure grains of the green and of the brown minerals before ignition. The brown alteration product contained much less uranium (4.2 percent) than did the green (20.0 percent). It was weakly birefringent with a mean index of refraction close to 1.64 in white light. After ignition for half an hour in the flame of a blast lamp, it showed a large increase in mean refractive index ($\Delta n = +0.14$) which was probably caused by a loss of water, and it turned greenish brown. In its mean index, color, high thorium and silicon contents, low uranium content, probable high water content, and in its association with other thorium minerals the brown mineral fits the descriptions of the simple hydrated species (?), hydrothorite, $\text{ThSiO}_4 \cdot 4\text{H}_2\text{O}$ (?).

The analytical data obtained on the sample of mixed brown and green thorium minerals free of other impurities are shown in table 1. Also shown are the corresponding data for (1) the Alaskan green mineral,

Table 1.--Chemical compositions of several uranothorites

| | Ranges of 4 earlier anal. <u>1/</u> in wt. percent | South Westland, N.Z. (Hutton 1950) <u>1/</u> | Placer, Ruby, Alaska Phair and Onoda <u>2/</u> | Green and brown minerals in fluorite Phair and Onoda <u>2/</u> |
|-------------------|--|---|---|--|
| Th | 40.6-56.6 c <u>3/</u> | 55.1 c | 56.6 c | >>10.0 s <u>4/</u> |
| U | 6.7-14.6 c | 10.1 c | 8.2 c | 11.8 c |
| Si | 6.1-9.1 c | 9.1 c | >10.0 s | >10.0 s |
| Fe | 0.9-5.3 c | 1.6 c | 0.1-1 s | 1-10 s |
| Ce | 0.07-1.18 c | 0.9 c | close to 0.1 s | < 0.1 s |
| Ca | 0.7-3.1 c | 0.2 c | close to 0.1 s | 1-10 s |
| Pb | 0.4-1.5 c | 0.5 c | not detected s | 0.1-1 s |
| H ₂ O+ | 9.24-11.31 c | 1.93 c | very low by qual. test | very low by qual. test |
| H ₂ O- | | | | |

Green mineral plus inclusions

U 20.0 c

Th, Zn major s (qual.)

Brown mineral plus inclusions

U 4.2 c

Th, Si, Zn major s (qual.)

1/ Analyses listed by Hutton 1950, converted to percentages of elements.

2/ Analyses by following members of U. S. Geological Survey: chemical analyses by F. S. Grimaldi, Mary H. Fletcher, Irving May, and Harry Levine; semi-quantitative spectrographic analyses by C. L. Waring; qualitative spectrographic analyses by Jules Stich.

3/ c indicates data obtained by chemical analysis.

4/ s indicates data obtained by semi-quantitative spectrographic analysis except where followed by (qual.) indicating that constituents were determined only as major, minor, or trace.

column 3; (2) definite uranothorite of similar composition described by Hutton (1950) column 2; (3) the ranges in composition shown by four analyses of other uranothorites listed in the older literature. The results of the analyses indicated that the green Alaskan mineral and the green Jamestown mineral (together with its hydrated brown alteration product) were chemically very similar. Each contained only the constituents typical of uranothorite in amounts within the ranges shown by chemical analyses of uranothorites from other localities. As might be expected the uranium content of the bulk sample of the Jamestown material (11.8 percent) was intermediate between the uranium contents of its separate green and brown mineral constituents. The percentage of uranium (20.0 percent) in the green Jamestown mineral is high even for the uranium-rich variety, uranothorite, but not excessively so. Uranothorites containing as much as 17 percent uranium (20.0 percent U_3O_8) are listed in the table of Canadian radioactive minerals compiled by A. H. Lang (1950).

The publication in July 1950 of Hutton's description of a definite nonmetamict placer uranothorite from South Westland, New Zealand, which had a maximum index of refraction, color, and specific gravity similar to those shown by the Alaskan and Jamestown thorium minerals, provided a useful standard of comparison and explained the differences in physical properties between these and the uranothorites previously described. Hutton pointed out that the older descriptions of thorite and uranothorite were based upon hydrated metamict material which contained on the average about 10 percent water. It is not surprising

that such material had low indices (generally in the range 1.68 to 1.72) and low specific gravities (generally in the range 5.19 to 5.40). Larsen (1921) reported that several Norwegian thorites showed a marked increase in index of refraction after ignition for half an hour over a blast lamp with resulting indices as high as 1.85. Neither the green uranothorite from the Alaskan placer nor that from the Blue Jay mine, Colo., showed an increase in refractive index after ignition for half an hour over a blast lamp; the Alaskan mineral actually showed a slight but probably real decrease in index. Presumably these uranothorites like Hutton's are relatively anhydrous, but unlike Hutton's they are metamict. Recently (February 1951) D. R. George described a green uranothorite from several California placers which has physical properties (color, refractive indices, and specific gravity) almost identical with the two uranothorites described in this paper; moreover, like these, it is completely metamict and is intergrown with a brown alteration product of lower mean index. The physical properties of the four comparable uranothorites are summarized in table 2.

Of 18 thin sections of parts of the fluorite breccia selected for high radioactivity, 6 showed one or more euhedra of green uranothorite, partly replaced by pseudomorphs of yellow-brown hydrothorite. Nearly all of these uranothorite euhedra form inclusions in fluorite. The square basal sections range in length from 0.02 to 0.10 mm, and the prismatic sections are only slightly longer. Within large grains of fluorite, halos that have maximum widths of 41 microns corresponding to the calculated range in fluorite of high-energy alpha particles

Table 2.--Comparison of physical properties of four uranothorites

| | S. Westland, N. Z. (Hutton 1950) | California (George 1951) | Ruby, Alaska (Phair, Onoda) | Blue Jay mine, Colo. (Phair, Onoda) |
|----------------------------|--|-----------------------------|---|---|
| System | tetragonal | tetragonal | tetragonal | tetragonal |
| Specific gravity | 6.7 ± 0.05 | 6.36 | 6.5 ± 0.2 | ---- |
| Color in oils | green (dichroic) | blue green | greenish | green |
| Optical characteristics | uniaxial + | isotropic (metamict) | isotropic (metamict) | isotropic (metamict) |
| Refractive indices | $n_X = 1.815$ $n_Y(\text{max}) = 1.850$ | $n = 1.82 - 1.87$ | $n = 1.86 \pm 0.01$ | $n = 1.86 \pm 0.01$ |
| X-ray diffraction patterns | ---- | ---- | anom. f-c cubic $a = 5.60 \text{ \AA}$ | anom. f-c cubic $a = 5.60 \text{ \AA}$ |

emitted by Po^{212} (Th C') serve as conspicuous indicators of uranothorite (fig. 1). Alpha-sensitive nuclear emulsions applied to uncovered thin sections and developed in the manner described by L. R. Stieff and T. W. Stern (1950) showed, under suitably high magnification, swarms of alpha tracks directed radially outward from the uranothorite cores of the halos. Thorium halos in fluorite, unlike uranium halos, have not been previously reported. These halos, together with the more abundant uranium halos (maximum widths of 0.031 micron) and other color effects induced by radioactivity in the Jamestown fluorites, will be described in greater detail in a forthcoming report.

DISCUSSION

In thin section most of the uranothorite is associated with relatively unshredded fluorite and is almost completely unshredded, retaining its sharp euhedral outlines (fig. 1). The lack of visible crushing indicates that the uranothorite crystals were not introduced into the fluorite mechanically in the course of the brecciation. The close association of the uranothorite with a fluorite-carbonate-sulfide assemblage, and in particular its tendency to form inclusions in the fluorite, is strong evidence of a hydrothermal origin. Apparently all thorites and uranothorites previously described in the literature have been found either as residual concentrates in placers or in situ in granitic igneous rocks, especially pegmatites. The one exception to this rule is a very doubtful occurrence of a single microscopic crystal of thorite (?) noted by Gordon (1944) in a sample from the high-

temperature tin veins of Llallagua, Bolivia. Its tentative identification as thorite was based upon morphological measurements; its interfacial angles were intermediate between thorite and xenotime but closer to thorite.

The temperatures at which the Jamestown uranothorite was deposited may have reached the mesothermal range but were certainly no higher. The Jamestown fluorite deposits are at the northern end of the Front Range mineral belt--a region wherein the ore deposits assume many epithermal characteristics in contrast to the southern end of the belt at which mesothermal conditions prevailed (Lindgren, 1933). Lindgren lists the Jamestown fluorite deposits with mesothermal ores, and Goddard (1946) noted that enargite and tennantite were among the early sulfides in many of the ores. However, the brecciation that accompanied the latest stages of fluorite mineralization left open spaces and could have taken place only under relatively near-surface conditions.

Previously thorite has been considered a rare mineral, but it seems likely that, once the properties of the unaltered mineral become more widely known, it will be found to have a rather widespread distribution not only in "high-temperature" granitic pegmatites and in placers, but also in many hydrothermal veins.

The rare-earth contents of many fluorites are relatively high; samarium, neodymium, and cerium in a fluorite described by Chang (1945) range from 0.1 to 1.0 percent. Heinrich (1948) found that "fluorite and rare-earth minerals are associated in 19 of the 22 pegmatite districts in North America from which fluorite has been reported."

Th^{4+} has many chemical properties in common with the rare earths, and its ionic radius is in the same range. The oxides UO_2 , ThO_2 , CeO_2 , and the fluoride CaF_2 all have the face-centered cubic "fluorite" structure with the unit cell close to 5.50 Å. The preference of the uranothorite and fluorite for the same geochemical environment is, therefore, readily explained by the Goldschmidt rule. It seems probable that thorium minerals as well as rare-earth minerals will prove to be of widespread, but scattered, distribution not only in fluorite-bearing pegmatites, but also in many fluorite-rich hydrothermal deposits.

REFERENCES

- Chang, Ting-Chao, 1945, Chemical studies on Chinese fluorites III: Shanghai Sci. Inst. Jour., new ser., vol. 2, no. 5.
- George, D. R., 1951, Thorite from California, a new occurrence and variety: Am. Mineralogist vol. 36, p. 129.
- Goddard, E. N., 1946, Fluorspar deposits of the Jamestown district, Boulder County, Colo.: Colorado Sci. Soc. Proc. vol. 15, no. 1, p. 19.
- Gordon, S. G., 1944, The mineralogy of the tin mines of Cerro de Llallagua, Bolivia: Acad. Nat. Sci. Philadelphia Proc. vol 96, p. 329.
- Heinrich, E. W., 1948, Fluorite-rare earth mineral pegmatites of Chaffee and Fremont Counties, Colo.: Am. Mineralogist vol. 33, p. 64.
- Hutton, C. O., 1950, Studies of heavy detrital minerals: Geol. Soc. America Bull. 61, no. 7, pp. 677-682.
- Lang, A. H., 1950, Summary account of Canadian uranium deposits, Table I: Canada Geol. Survey.
- Larsen, E. S., Jr., 1921, The microscopic determination of the non-opaque minerals: U. S. Geol. Survey Bull. 679, p. 144.
- Lindgren, W., 1933, Mineral deposits, McGraw-Hill Book Co., p. 635.
- Robinson, S. C., 1950, Canada Geol. Survey Memorandum.
- Stieff, L. R., and Stern, T. W., 1950, The preparation of nuclear-track plates and stripping films for the study of radioactive minerals: U. S. Geol. Survey Trace Elements Investigations Rept. 127.




Figure 1.--Thorium halo surrounding euhedral uranothorite inclusion in fluorite. A basal section of uranothorite forms the center of a thorium halo consisting of a semi-opaque inner "ring" and an outer bleached "ring." The width of the halo is 41 microns as measured from the inner boundary of the darkened zone to the outer boundary of the bleached zone. (Uncrossed nicols.)

END OF DOCUMENT