

(200)
T6712

UNCLASSIFIED

Geology - Mineralogy

This document consists of 5 pages.
Series A.

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

SECOND OCCURRENCE OF BAYLEYITE IN THE UNITED STATES *


By

T. W. Stern and Alice Dowse Weeks

July 1952

Trace Elements Investigations Report 243

*Am. Mineralogist, 1.37 nos 11-12,
p. 1058-1060, 1952.*



* This report concerns work done on behalf of the Division
of Raw Materials of the U. S. Atomic Energy Commission

USGS - TEI Report 243

GEOLOGY - MINERALOGY

<u>Distribution (Series A)</u>	<u>No. of copies</u>
American Cyanamid Company, Watertown	1
Argonne National Laboratory	1
Atomic Energy Commission, Washington	2
Carbide and Carbon Chemicals Company, Y-12 Area	1
Colorado Raw Materials Office (F. H. MacPherson)	1
Division of Raw Materials, Grand Junction	6
Division of Raw Materials, Grants	1
Division of Raw Materials, Denver	1
Division of Raw Materials, Hot Springs	1
Division of Raw Materials, New York	6
Division of Raw Materials, Salt Lake City	1
Division of Raw Materials, Richfield	1
Division of Raw Materials, Butte	1
Division of Raw Materials, Washington	3
Dow Chemical Company, Pittsburg	1
Technical Information Service, Oak Ridge	6
U. S. Geological Survey:	
Mineral Deposits Branch, Washington	1
Geochemistry and Petrology Branch, Washington	10
Geophysics Branch, Washington	1
Alaskan Geology Branch, Washington	1
Fuels Branch, Washington	1
V. E. McKelvey, Washington	1
L. R. Page, Denver	1
R. P. Fischer, Grand Junction	1
A. E. Weissenborn, Spokane	1
J. F. Smith, Jr., Denver	1
N. M. Denson, Denver	1
C. B. Read, Albuquerque	1
A. H. Koschmann, Denver	1
C. A. Anderson, Tucson	1
R. J. Roberts, Salt Lake City	1
TEPCO, Washington	4
(Including master)	

SECOND OCCURRENCE OF BAYLEYITE
IN THE UNITED STATES

by

T. W. Stern and Alice Dowse Weeks

ABSTRACT

The mineral bayleyite was first described in 1951 from a mine in Yavapai County, Ariz. This report describes bayleyite from the Hideout No. 1 mine, San Juan County, Utah, together with its associated minerals and mineralogic properties.

Bayleyite, $\text{Mg}_2\text{UO}_2(\text{CO}_3)_3 \cdot 18\text{H}_2\text{O}$, is one of three new uranium minerals first described from the Hillside mine, Yavapai County, Arizona, by Axelrod, Grimaldi, Milton, and Murata (1951). The minerals occurred at the 300-foot level of the mine and the authors state (1951, p. 2) that "In the mine, bayleyite crystals are sharp and well faceted. In the drier atmosphere of Washington [D. C.] they soon become dull, losing their luster and finally disintegrate to a yellow powder." Branche, Chervet, and Guillemin (1951) who described bayleyite from Morocco, the only foreign locality for this mineral, also noted that the crystals disintegrated rapidly, but they ascribed this property to the humidity of their laboratory.

In July 1951 T. W. Stern collected a yellow efflorescent crust of a highly radioactive mineral from Hideout No. 1 mine on Deer Flats, along the north side of White Canyon, San Juan County, Utah, when the authors visited the mine with L. B. Riley and W. E. Benson also of the U. S. Geological

Survey. Fortunately this material did not dehydrate as the Arizona and Morocco samples had, for when it was studied nine months later it was found to be fully hydrated bayleyite. The Hideout mine is thus the second bayleyite locality in the United States.

The Hideout No. 1 mine, not much more than a prospect in July 1951, is in a copper-uranium deposit in the Shinarump conglomerate (Triassic). The chief minerals observed were chalcocite, bornite, chalcopyrite, brochantite, malachite, and azurite, which replace fossil wood and impregnate coarse arkosic sandstone. Bayleyite and schroeckingerite are the only uranium minerals so far identified from this mine. They occur in the adit less than 100 feet from the cliff face, in contrast to the Arizona specimens which were collected from the 300-foot level. The massive pieces of copper sulfide replacing wood show some radioactivity with the Geiger counter, but no uranium mineral can be seen under a binocular microscope.

The bayleyite specimen from Hideout No. 1 mine is translucent, light yellow, and occurs in thin crusts that break away easily from the underlying rock. The bayleyite consists of microscopic, bladed or prismatic crystals with terminal faces on the upper surface of the crust. The optical properties determined in white light are similar to the type and Moroccan materials, with $\alpha = 1.454 \pm 0.003$, $\beta = 1.492 \pm 0.003$, and $\gamma = 1.502 \pm 0.003$, the extinction angle between X and the c axis is 14° , and Z is parallel to the b axis. It is not possible to obtain an acute bisectrix interference figure because the cross section of the blades is too small. The mineral is faintly pleochroic with pale pink in the X direction and light yellow to greenish yellow in the Y and Z directions.

Bayleyite is water-soluble and it effervesces vigorously and quickly dissolves in cold dilute hydrochloric acid. A semiquantitative spectrographic

analysis, made by C. S. Annell of the Geological Survey, shows:

U	> 10 percent
Mg, Ca (low)	1.0 to 10 percent
Al	0.1 to 1.0 percent
Si, Sr	0.01 to 0.1 percent

The calcium is probably due to a little gypsum and schroeckingerite that occur with the bayleyite. The schroeckingerite is in fine-grained greenish-yellow flakes on the under side of the bayleyite. When activated with long wavelength fluorescent light (3663 Å) the schroeckingerite fluoresces vivid yellow green and the bayleyite fluoresces faintly whitish green. The schroeckingerite gives the superficial appearance of the dehydrated bayleyite described by Axelrod, Grimaldi, Milton, and Murata (1951, p. 2), but an X-ray diffraction powder pattern shows only schroeckingerite lines and no metabayleyite lines. The bayleyite gives an X-ray pattern identical to that of the Arizona material.

This report concerns work done on behalf of the Division of Raw Materials of the U. S. Atomic Energy Commission.

REFERENCES

- Axelrod, J. M., Grimaldi, F. S., Milton, C., and Murata, K. J., 1951, The uranium minerals from the Hillside mine, Yavapai County, Arizona: *Am. Mineralogist*, vol. 36, pp. 1-22.
- Branche, C., Chervet, J., and Guillemin, C., 1951, Nouvelles espèces uranifères françaises: *Soc. franç. minéralogie et cristallographie Bull.*, vol. 74, pp. 458-461.