



HYDROLOGIC STUDIES  
**OFFICE COPY**

*Vowinkel*

STATE OF NEW JERSEY

DEPARTMENT OF CONSERVATION AND ECONOMIC DEVELOPMENT

DIVISION OF WATER POLICY AND SUPPLY

# **RECORDS OF WELLS and GROUND-WATER QUALITY in BURLINGTON COUNTY, NEW JERSEY**

**WATER RESOURCES CIRCULAR No. 7**

Prepared in cooperation with  
United States Department of the Interior  
Geological Survey

1962

**RECORDS OF WELLS AND GROUND-WATER QUALITY  
IN BURLINGTON COUNTY, NEW JERSEY**

**A Preliminary Report  
By  
F. Eugene Rush  
Geologist  
U. S. Geological Survey**

**Prepared by the U. S. Geological Survey  
in cooperation with the  
State of New Jersey**





**State of New Jersey**

**Department of Conservation  
and Economic Development**

**H. Mat Adams, Commissioner**

**Division of Water Policy and Supply**

**George R. Shanklin, Chief Engineer and Acting Director**

**Water Policy and Supply Council**

Kenneth H. Murray, Chairman . . . . .	Tewksbury
William G. Bank . . . . .	Colts Neck
I. Ralph Fox . . . . .	Rumson
Herman A. Klenner . . . . .	Rahway
Thomas J. Mullen . . . . .	Oldwick
August C. Schultes . . . . .	Woodbury
Lillian M. Schwartz . . . . .	Highland Park
David I. Stepacoff . . . . .	Perth Amboy
Vinton N. Thompson . . . . .	Vincentown

**John Wyack, Secretary**





## CONTENTS

	Page
Abstract .....	1
Introduction .....	1
Purpose and scope.....	1
Location and extent .....	2
General geology and hydrology .....	2
Well data .....	2
Chemical character of water .....	4
References .....	5

## ILLUSTRATIONS

Figure 1. Map showing location of selected wells in Burlington County, N. J. ....	Pocket
--	--------

## TABLES

	Page
Table 1. Age, thickness, water-bearing character, and lithology of stratigraphic units in Burlington County, N. J. ....	3
2. Records of selected wells in Burlington County, N. J. ....	6
3. Selected drillers' logs of wells in Burlington County, N. J. ....	14
4. Chemical analyses of water from wells in Burlington County, N. J. ....	103





# **RECORDS OF WELLS AND GROUND-WATER QUALITY IN BURLINGTON COUNTY, NEW JERSEY**

by

**F. Eugene Rush**

## **ABSTRACT**

This basic-data report is limited to the presentation of selected drillers' logs and their geologic correlations, well data, and chemical analyses of water from wells in Burlington County, N. J. The county lies within the Coastal Plain province; it is underlain by unconsolidated sediments of Quaternary, Tertiary, and Cretaceous age and at greater depth by much older consolidated rocks.

## **INTRODUCTION**

### **Purpose and Scope**

The investigation of the ground-water resources of Burlington County, New Jersey is part of a state-wide program of water-resources studies being carried out by the U. S. Geological Survey in cooperation with the New Jersey Department of Conservation and Economic Development, Division of Water Policy and Supply. The investigation has been under the general supervision of Allen Sinnott, District Geologist.

This basic-data report is limited to the presentation of selected drillers' logs, well data, and chemical analyses of water from wells in Burlington County. Most of the drillers' logs and the well data were obtained from the files of the New Jersey Geological Survey, Bureau of Topography and Geology, Division of Research and Development, Department of Conservation and Economic Development. The information is released at this time in advance of an interpretive report because of its value to prospective ground-water users in Burlington County. The data in this report form, in part, the basis for a comprehensive, interpretive report on the ground-water resources of Burlington County, now in preparation.

### Location and Extent

Burlington County is in the southcentral part of New Jersey, extending from the Delaware River on the northwest to Great Bay, an arm of the Atlantic Ocean, on the southeast. The county is bounded by Camden and Atlantic Counties on the southwest and by Mercer, Monmouth, and Ocean Counties on the northeast. (See fig. 1.) Burlington County is the largest county in New Jersey in area, covering 827 square miles. It has a population of 224,499 in the 1960 Federal Census. While the county is principally known for its agriculture, there is considerable manufacturing, particularly along the Delaware River.

### GENERAL GEOLOGY AND HYDROLOGY

Burlington County lies within the Coastal Plain province; it is underlain by layers of sand, silt, and clay of Quaternary, Tertiary, and Cretaceous age. The formation names given to these deposits and the basement crystalline rocks are listed in table 1, which gives their age, thickness, and water-bearing character. The important water-bearing formations in the county are the Raritan and Magothy Formations, the Englishtown Formation, the Wenonah Formation and Mount Laurel Sand, and the Kirkwood Formation and Cohansey Sand.

### WELL DATA

Descriptions of selected wells are given in table 2. The locations of these wells are shown on figure 1. The drillers' logs in this report have been modified, where necessary, to present a more uniform terminology. The drillers' logs and the author's tentative geologic correlations are given in table 3.

Table 1. – Age, thickness, water-bearing character, and lithology of stratigraphic units in Burlington County, N. J.

System	Series	Formation	Thickness (feet)	Water-bearing character	Description
Quaternary	Pleistocene	Cape May Formation	0-40	Variable	Sand and gravel, local clay
		Pensauken Formation	0-40	Variable	Sand and gravel
Tertiary	Pliocene (?)	Cohansey Sand	0-300	Excellent to fair	Sand, medium to coarse, with local ironstone
	Miocene	Kirkwood Formation	0-200	Excellent to fair	Sand, fine, some clay seams
	Eocene	Manasquan Formation	0-100	Poor	Sand, glauconite, clayey
	Paleocene	Vincentown Formation	0-100	Good to poor	Sand, limy, clayey; glauconite
		Hornerstown Sand	0-60	Poor	Glauconite sand, with green clay matrix
Cretaceous		Red Bank Sand	0-40	Fair	Sand and clay with glauconite sand
		Navesink Formation	0-40	Poor	Glauconite sand, with dark gray clay matrix
		Mount Laurel Sand and Wenonah Formation }	0-100	Good to poor	Sand, clayey, some glauconite; silt
		Marshalltown Formation	0-100	Poor	Clay, sandy; glauconite sand
		Englishtown Formation	0-100	Good to poor	Sand and clay
		Woodbury Clay	0-120	Poor	Clay, black, micaceous
		Merchantville Formation	0-80	Poor	Clay, dark gray, glauconite sand
		Magothy Raritan Formations }	30-800	Excellent to good	Clay and sand, alternating
Early Paleozoic (?)		Wissahickon Formation		Poor	Schist or gneiss, commonly weathered



## CHEMICAL CHARACTER OF WATER

All ground water contains dissolved mineral matter derived from the sediments, rocks, and soils with which the water has been in contact. The amount and kind of dissolved minerals determine the hardness, corrosiveness, scale-forming properties, and other physical-chemical characteristics of the water. The constituents most likely to occur in objectionable concentrations in ground water are silica ( $\text{SiO}_2$ ), calcium (Ca), magnesium (Mg), iron (Fe), manganese (Mn), chloride (Cl), and nitrate ( $\text{NO}_3$ ). In addition, the ground-water properties of importance are dissolved solids, specific electrical conductance, hardness, and pH.

According to Rainwater and Thatcher (1960, p. 141-259), silica in water appears to have little effect on human beings, livestock, fishes, or plants. It is of chief concern to industrial users because it contributes to the formation of hard "boiler scale" on heat-exchange equipment and steam turbine blades. Iron and manganese cause stains on textiles and fixtures if their combined concentration exceeds 0.4 parts per million (ppm); they also cause an unpleasant taste. Chloride is noticeable to the taste at concentrations of above 250 ppm and renders water unpotable at concentrations above 1,000 ppm.

Calcium and magnesium cause most of the hardness of water (Hem, 1959, p. 145). Nitrate, an end product of decomposition of organic matter, when present in concentrations greater than 5 ppm may be indicative of pollution. Water containing more than 44 ppm of nitrate may cause infant cyanosis ("blue-baby disease"), if used in infant-feeding formulas (Maxcy, 1950). The highest concentration of nitrate given on table 4 is 21 ppm, but it generally is less than 1 ppm.

The dissolved-solids content is an index of the total mineralization of the water. Specific conductance is a relative measure of the presence of dissociated ions in solution and thus is also an index of the total mineralization of the water. The specific conductance multiplied by 0.6 to 0.7 gives an approximation of the dissolved-solid content in parts per million.

The property of hardness is difficult to define, in spite of its wide usage (Hem, 1959, p. 145-148). It can be said that hardness represents the soap-consuming power of a water. However, it is commonly expressed in an analysis as an equivalent quantity of calcium carbonate ( $\text{CaCO}_3$ ). The degree of acidity or alkalinity of water is denoted by pH, on a scale of 1 to 14. Water having a pH of 7.0 is neutral; a pH of less than 7.0 is acidic, and more than 7.0, alkaline. However, in water chemistry a pH of below 4.5 may indicate free mineral acids, whereas above 4.5 some alkaline ions may be present (Hem, 1959, p. 48). The pH value of water is primarily of concern to industrial users of water. Chemical analyses of water from wells in Burlington County, N. J. are given in table 4.

## REFERENCES

- Hem, J. D., 1959, Study and interpretation of the chemical characteristics of natural waters: U. S. Geol. Survey Water-Supply Paper 1473, 269 p.
- Kummel, Henry B., 1910, Annual report of the State Geologist: New Jersey Geol. Survey Ann. Rept. for 1909, Trenton, N. J., 123 p.
- Maxcy, K. F., 1950, Report on the relation of nitrate concentrations in well waters to the occurrence of methemoglobinemia: Nat. Research Council, Bull. Sanitary Eng., p. 265 App. D.
- Rainwater, F. H., and Thatcher, L. L., 1960, Methods for collection and analysis of water samples: U. S. Geol. Survey Water-Supply Paper 1454, 301 p.
- Richards, Horace G., 1948, Studies on the subsurface geology and paleontology of the Atlantic Coastal Plain: Philadelphia Acad. Nat. Sci. Proc., v. C, p. 39-76.
- 1954, A new gastropod and other fossils from the Cretaceous of New Jersey: Notulae Naturae, no. 258, Philadelphia Acad. of Nat. Sci. Proc., 5 p.
- Smock, John C., 1893, Annual report of the State Geologist: New Jersey Geol. Survey Ann. Rept. for 1892, Trenton, N. J., 367 p.
- 1894, Annual report of the State Geologist: New Jersey Geol. Survey Ann. Rept. for 1893, Trenton, N. J., 457 p.
- 1901, Annual report of the State Geologist: New Jersey Geol. Survey Ann. Rept. for 1900, Trenton, N. J., 231 p.

Table 2.--Records of selected wells in Burlington County, N. J.

Well no.: Numbers correspond to those in figure 2 and tables 3 and 4; A. indicates chemical analysis given in table 4, L. indicates driller's log given in table 3.

Water-bearing formation: Pw, Wissahickon Formation; Kmr, Magothy and Raritan Formation; Ket, Englishtown Formation; Km, Mount Laurel Sand and Wenonah Formation; Tvt, Vincentown Formation; Tk, Kirkwood Formation; Tch, Cohansy Sand; Qcm, Cape May Formation.

Use: CI, commercial and institutional; D, domestic; Ind, industrial; Irr, irrigation; O, observation; PS, public supply; U, unused.

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
1	Bordentown	Bordentown Water Dept. No. 1	Layne-New York Co., Inc.	1955	89	441	16 & 8	355	355-385	Kmr	86	572	17	PS		1
2	1/2 mi. E of U.S. Highway 206, at Crosswicks Creek	Bordentown Water Dept. White Horse well field No. A 40		1941	9	80	3	75	(none)	do.	(flowing)	25		PS	Approximately 43 flowing wells in field, all with similar construction, 2", 3" or 4" in diam., with total depths of 75'-80'.	2
3	Burlington	Burlington Water Dept. No. 3	Layne-New York Co., Inc.	1949	9	90	16	64	64-85	Kmr	10	517	48	PS	A., L.	3
4	1 mi. NE of Burlington on Burlington Island	Burlington Water Dept. Test Well No. 2 - Island	Artesian Well Drilling Co.	1952	19	49	6	40	10	Qcm	14	75	8	O	L.	4
5	1 mi. N of Burlington on Burlington Island	Burlington Water Dept. No. 2 - Island	do.	1952	20	49	17	33	33-49	Qcm	18	760	13	PS	Concrete casing and screen. A.	5
6	Columbus	Columbus Water Co. No. 2	William Stothoff Co.	1943	73	260	8	240	240-260	Kmr	60	125	40	PS	A.	6
7	1 mi. N. of Crosswicks	Crosswicks Water Co. No. 1	do.	1959	30	299	8	269	269-299	Kmr	15	500	44	PS	A.	7
8	Beverly	Delaware River Water Co. No. 15	A. C. Schultes & Sons	1951	12	59	12	49	47-57	Kmr	18	1200	14	PS	Drilled to bedrock. A.	8
9	Riverside	do. No. 17	do.	1955	21	140	12	97	97-118	Kmr	23	483	70	PS	do. L.	9
10	1 mi. SW of Riverside, at Cambridge	do. No. 19	do.	1959	25	143	12	91	89-130	Kmr	14	974	59	PS	do. L.	10
11	Marlton	Evesham Tp. Water Dept. No. 1	A. G. Dunphy	1897	115	212	6	--	-----	Ket	42	100	36	PS	A., L.	11
12	Marlton	do. No. 3	Layne-New York Co., Inc.	1956	85	512	12 & 8	369	369-389	Kmr	91	517	168	PS	L.	12
13	Florence	Florence Tp. Water Dept. No. 3	do.	1949	25	162	16	123	123-138	Kmr	24	620	34	PS	Drilled to bedrock. A., L.	13
14	2 mi. S of Wrightstown	Fort Dix No. 1	Artesian Well Drilling Co.	1941	138	980	20, 16, & 10	920	916-960	Kmr	132	800	96	PS	A.	14
15	1 mi. S of Wrightstown	do. No. 2	do.	1941	131	1051	do.	1030	1030-1051	Kmr	123	900	61	PS	A., L.	15
16	3 mi. S of Wrightstown	do. No. 4	Layne-New York Co., Inc.	1943	148	1096	do.	1056	1056-1086	Kmr	140	700	70	PS	A., L.	16
17	3 mi. NE of Browns Mills at Hanover Lake	do. No. 11	William Stothoff Co.	1943	85	485	6	470	470-980	Ket	20	45	193	PS	L.	17
18	1 mi. E of Wrightstown	McGuire Air Force Base - A	A. C. Schultes & Sons	1953	128	1100	24, 16, & 10	992	992-1055	Kmr	120	913	63	PS	Drilled to weathered bedrock. A., L.	18
19	2 mi. E of Wrightstown	do. - C	do.	1953	102	1123	do.	1036	1036-1089	Kmr	110	900	45	PS	Drilled to weathered bedrock. A., L.	19
20	2 mi. SE of Wrightstown	do. - D	do.	1953	112	1120	do.	1012	1012-1075	Kmr	110	925	34	PS	Drilled to weathered bedrock. A.	20
21	3 mi. SW of Medford, at Lake Pine	Lakes Water Co. No. 2	J. Henry Robbins	1950	52	200	6	180	(none)	Kmr	20	100	--	PS	A., L.	21
22	Levittown	Levitt and Sons, Inc. No. 2	A. C. Schultes & Sons	1955	29	363	20, 16, & 12	225	225-354	Kmr	22	1404	65	U	Screen blanks 246-292 and 315-344. L.	22
	do.	do. No. 3	C. W. Lauman Co., Inc.	1959	28	381	do.	203	203-303	Kmr	25	1530	67	PS	Screen blank 238-283.	23



Table 2.--Records of selected wells in Burlington County, N. J.--Continued

Map no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
24		No. 4	A. C. Schultes & Sons	1958	28	284	do.	177	196-279	Kmr	21	1438	88	PS	Screen blank 216 259. A., L.	24
25	do.	do. No. 5	do.	1958	39	269	do.	230	230-256	Kmr	37	1421	144	PS	A., L.	25
26	do.	do. No. 7	C. W. Lauman Co., Inc.	1958	19	306	do.	179	179-255	Kmr	17	1421	56	PS	A., L.	26
27	do.	do. No. 12	A. C. Schultes & Sons	1955	40	526	---	(none)	(none)	---	---	---	---	U	Abandoned. Drilled as a test hole. L.	27
28	3 mi. S of Mount Holly, at Lumberton	Lumberton Light, Water and Sewage Co. No. 1	Artesian Well Drilling Co.	1928	10	404	10, 8, & 6	360	360-401	Kmr	22	100	---	PS	A., L.	28
29	Maple Shade	Maple Shade Tp. Water Dept.	Uriah White	1893	55	375	---	---	---	Kmr	---	---	---	PS	Abandoned. L.	29
30	do.	do. No. 4	Artesian Well Drilling Co.	1955	10	282	20 & 12	211	211-272	Kmr	19	1020	42	PS	L.	30
31	Medford	Medford Water Co. No. 3	Layne-New York Co., Inc.	1957	48	590	12 & 8	506	506-536	Kmr	48	517	86	PS	L.	31
32	Moorestown	Moorestown Tp. Water Dept. No. 1	do.	1923	20	497	12, 8, & 6	350	350-372	Kmr	28	450	114	PS	Drilled to bedrock. A., L.	32
33	Mount Holly	Mount Holly Water Co. No. 3	Layne-New York Co., Inc.	1953	11	380	16 & 10	316	316-346	Kmr	64	1200	132	PS	A.	33
34	do.	Mount Holly Water Co.	Thomas B. Harper	1900	20	562	8	---	---	Kmr	---	---	---	PS	Abandoned. L.	34
35	Pemberton	Pemberton Water Dept. No. 1	A. L. Lyons	1939	80	206	12 & 8	155	155-185	Kmr	30	168	51	PS	A., L.	35
36	Browns Mills	Pemberton Tp. Water Dept. No. 2	Charles Bainbridge	1947	63	282	8	257	(none)	Kmr	18	187	27	PS	A.	36
37	do.	do. No. 3	Thomas Magee	1955	112	303	8	277	277-303	Kmr	47	250	51	PS	L.	37
38	do.	do. No. 4	Layne-New York Co., Inc.	1960	93	402	12 & 8	294	294-334	Kmr	60	500	141	PS	A., L.	38
39	4 mi. W of Mount Holly at Rancocas Woods	Rancocas Woods Water Co. No. 1	Charles L. Mollitor, Inc.	1954	50	248	6	237	237-248	Kmr	50	150	5	PS	L.	39
40	Palmyra	Riverton-Palmyra Water Co. No. 8	A. C. Schultes & Sons	1955	13	100	16	64	64-84	Kmr	26	503	32	PS	Drilled to weathered bedrock. A., L.	40
41	1 mi. E of Pemberton at Sunbury Village	Sunbury Village Water Co. No. 1	William Stothoff Co.	1953	62	198	8	178	178-198	Kmr	17	252	119	PS	A.	41
42	2 mi. SE of Bordentown	Norman F. Scattergood	Theodore Ziegenfuss	1952	72	362	6	348	348-359	Kmr	60	60	40	D	Farm use also. L.	42
43	1 mi. E of Bordentown	Mrs. Helen Swayze	William L. Bainbridge	1952	92	279	6	274	274-279	Kmr	68	15	---	D		43
44	1 mi. S of Bordentown on U.S. Hwy. 130	National Guard Shops	Louis Bainbridge	1952	83	230	6	---	---	Kmr	---	---	---	CI	L.	44
45	2 mi. NE of Burlington	Hercules Powder Co. No. 1	Layne-New York Co., Inc.	1946	22	146	10	105	105-135	Kmr	10	520	21	CI	Drilled to bedrock. A., L.	45
46	do.	do. No. 1-test	do.	1945	13	132	3	94	94-104	Kmr	5	---	---	O	Drilled to bedrock.	46
47	do.	do. No. 3-test	do.	1945	16	121	3	94	94-104	Kmr	14	---	---	O	do. A.	47
48	2 mi. SW of Burlington on U.S. Hwy. 130	Edgar B. Heisler	Charles L. Mollitor, Inc.	1958	30	215	10	185	185-215	Kmr	33	600	82	Irr	L.	48
49	1 mi. SW of Burlington on U.S. Hwy. 130	Carroll Murphy	do.	1950	35	83	8	64	64-82	Kmr	15	250	45	Irr	L.	49
50	2 mi. SE of Burlington	Masonic Home	American Well Drilling Co.	1921	79	395	8	174	174-194	Kmr	65	150	45	U	A., L.	50
51	3 mi. E of Burlington	William Donald	Charles L. Mollitor, Inc.	1955	42	111	10	72	72-106	Kmr	26	275	14	Irr	L.	51
52	3 mi. SE of Burlington on Oxmead Rd.	Russell H. Deacon	W. R. Clair	1958	62	208	4	203	203-208	Kmr	62	30	---	D		52
53	3 mi. SE of Burlington	Roy F. Main	E. Weik	1959	67	215	4	207	207-215	Kmr	60	30	5	D		53
54	3 mi. SE of Burlington on Oxmead Rd.	Cherry Hill Inn Farm No. 1	Charles L. Mollitor, Inc.	1950	72	208	6	199	199-206	Kmr	82	10	36	D	Farm use also. L.	54
55	2 mi. SW of Burlington	Frank Pino	Edward Robbins	1950	63	228	8 & 6	208	208-228	Kmr	60	300	30	Irr	Pump capacity is 400 gpm. L.	55

Table 2.--Records of selected wells in Burlington County, N. J.--Continued

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
56	1 mi. SW of Florence	National Gypsum Co. No. 2	Layne-New York Co., Inc.	1955	22	157	12 & 8	113	113-123	Kmr	21	183	21	Ind	Drilled to bedrock. L.	56
57	2 mi. SW of Burlington on U.S. Hwy. 130	Mechanics National Bank No. 2	Charles L. Mollitor, Inc.	1958	35	209	4	189	189-209	Kmr	30	180	27	CI	Air conditioning recharge well.	57
58	1 mi. S of Burlington	Charles Heal, Jr.	do.	1955	50	80	8	59	59-80	Kmr	36	400	50	Irr	Water pumped into irrigation pond. L.	58
59	2 mi. SW of Burlington	Anthony Cannuli	Edward Robbins	1950	66	173	6	153	153-173	Kmr	55	200	30	Irr	Water pumped into irrigation pond.	59
60	2 mi. SW of Burlington on U.S. Hwy. 130	Albert Heisler	do.	1950	46	252	8	212	212-252	Kmr	41	500	42	Irr	Pump capacity is 500 gpm.	60
61	11 mi. S of Chatsworth, at Harrisville	Harrisville well	--	1866	15	375	8	---	---	TKw (flowing)	---	---	---	U	Well is still flowing.	61
62	2 mi. NE of Bordentown	N.J. State Reformatory No. 4	Layne-New York Co., Inc.	1951	97	454	12 & 8	357	357-387	Kmr	86	503	36	U	A., L.	62
63	4 mi. SE of Bordentown, at Chesterfield	H. W. Briel	Howard Farmer	1959	93	272	4	266	266-272	Kmr	72	50	10	D		63
64	1 mi. SW of Chesterfield, on Whitepine Rd.	Ralph J. DeMarco	Louis Bainbridge	1958	82	393	4	388	388-392	Kmr	75	4	45	D		64
65	3 mi. SE of Bordentown, on Old York Rd.	Ernest Liptak	Greenhalgh & Kaye	1957	114	214	4	208	208-214	Kmr	100	10	20	D		65
66	2 mi. N of Wrightstown	Jacob Dana Packing Co. No. 3	do.	1958	208	276	6	255	255-276	Ket	100	60	80	CI	L.	66
67	7 mi. N of Wrightstown, on Ellisdale Rd.	Francis D. Gray	Louis Bainbridge	1960	92	329	6	319	319-325	Kmr	75	40	5	D	Used for swimming pool also. A., L.	67
68	8 mi. N of Wrightstown, on Ellisdale Rd.	John Marincas	Greenhalgh & Kaye	1954	97	208	6	204	204-208	Kmr	80	25	69	D	L.	68
69	4 mi. N of Wrightstown	Joseph S. Kucowski	do.	1957	104	149	6	143	143-149	Ket	38	10	7	D		69
70	2 mi. NE of Riverton	Hoegaens Sponge Iron Co.	Layne-New York Co., Inc.	1951	35	136	8	119	119-134	Kmr	16	326	68	CI	Pump capacity is 50 gpm. L.	70
71	3 mi. NE of Mount Holly, on State Route 537	Eastampton School	Charles L. Mollitor, Inc.	1957	73	417	--	407	407-417	Kmr	70	75	30	U	L.	71
72	1 mi. NE of Mount Holly, on North Pemberton Rd.	J. A. Methvin	do.	1953	39	364	6	352	352-363	Kmr	63	30	15	D	L.	72
73	1 mi. NW of Marlton, on Cropwell Rd.	Byron T. Roberts Farms Inc. No. 2	A. C. Schultes & Sons	1957	93	406	18 & 8	322	322-375	Kmr	100	750	80	Irr	Screen blank 348-359. A., L.	73
74	1 mi. W of Marlton, on Old Marlton Pike	David D. Griscom	J. Henry Robbins	1952	98	340	4	320	320-340	Kmr	100	40	---	D	L.	74
75	2 mi. SE of Marlton	U.S. Army Nike Base, control area well	Edward Robbins, Sr.	1954	84	158	8	138	138-158	Kmr	41	200	43	U		75
76	1 mi. SE of Beverly, on U.S. Highway 130	Earl R. Blyler	Charles L. Mollitor, Inc.	1950	39	117	10	90	90-114	Kmr	14	575	19	Irr	L.	76
77	1 mi. SE of Beverly, on Cooper St.	John W. Heal	do.	1951	36	135	10	107	107-135	Kmr	18	500	89	Irr	A.	77
78	1 mi. SW of Beverly, at Cambridge	Amico Sand and Gravel Co.	do.	1957	15	75	6	27	27-37	Kmr	7	120	11	CI	L.	78
79	1 mi. S of Riverside, on U.S. Hwy 130	Hillside Farms No. 3	A. C. Schultes & Sons	1956	20	112	12	90	90-106	Kmr	6	444	68	CI		79
80	1 mi. SE of Riverside, on U.S. Hwy 130	Holy Cross High School	Charles L. Mollitor, Inc.	1958	82	174	10	154	154-174	Kmr	55	185	20	Irr	L.	80
81	1 mi. E of Riverside, on U.S. Hwy 130	Holiday Lake	do.	1958	25	198	8	188	188-198	Kmr	15	180	---	CI	L.	81

Table 2.—Records of selected wells in Burlington County, N. J.—Continued

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
82	3 mi. W of Mount Holly	Alfred Thomas	Artesian Well Drilling Co.	1954	34	75	4	70	70-74	Ket	31	4	19	D		82
83	4 mi. SW of Mount Holly	Fred Worth	Edward Robbins, Sr.	1949	30	317	8	277	277-297	Kmr	23	200	97	Irr	Pump capacity is 200 gpm.	83
84	3 mi. NE of Medford	Charles H. Clark	Charles L. Mollitor, Inc.	1956	51	191	4	184	184-190	Ket	25	35	20	D		84
85	Mount Holly	Plasticon Corp.	do.	1955	45	107	6	97	97-107	Ket	40	75	5	CI		85
86	3 mi. E. of Burlington	John and Nick Cannuli	do.	1955	41	93	10	72	72-92	Kmr	29	250	30	Irr	Pump capacity is 500 gpm.	86
87	do.	James Workman	Edward Robbins, Sr.	1952	41	194	10	174	174-194	Kmr	52	800	32	Irr	do.	87
88	3 mi. NW of Wrightstown on State Route 545	John H. Cook	Charles L. Mollitor, Inc.	1950	115	148	6	136	136-148	Ket	36	50	4	D	Water temperature is 54°F. L.	88
89	2 mi. S of Bordentown on U.S. Hwy 206	Turnpike Junction Ind. Park	A. C. Schultes & Sons	1958	60	340	10	293	293-329	Kmr	55	402	31	U	Screen blank 307-318. Undeveloped industrial park. A., L.	89
90	4 mi. S of Bordentown	Franklin Tallman	Charles L. Mollitor, Inc.	1949	82	195	6	184	184-195	Kmr	75	38	5	D	Farm use also. L.	90
91	3 mi. S of Bordentown	Ed Phares	Louis Bainbridge	1949	92	505	6	364	364-370	Kmr	125	---	---	D	Farm use also. Pump capacity is 20 gpm. L.	91
92	do.	George H. Preidel	Edward Robbins, Sr.	1952	102	215	6	205	205-215	Kmr	100	50	5	D	Farm use also.	92
93	do.	M. P. Woodward	W. R. Clair	----	72	239	3	234	234-239	Kmr	54	20	--	D		93
94	4 mi. SE of Bordentown	North Burlington Co. Regional Jr.-Sr. High School	Russell D'Agostino	1959	83	350	8 & 6	290	290-310	Kmr	85	200	35	CI	Water contains iron. L.	94
95	6 mi. E of Burlington	Ralph R. Parker	W. R. Clair	1959	62	380	4	370	370-380	Kmr	50	50	5	D	Farm use also. L.	95
96	5 mi. E of Burlington	Franklin Wainwright	do.	----	62	209	3	204	204-209	Kmr	42	20	--	D		96
97	1 mi. N of Medford on Church Rd.	Peter Martin	John S. Wrobel	1949	62	126	3	123	123-126	Kmr	21	10	2	D	A., L.	97
98	2 mi. NW of Medford on Church Rd.	William E. Johnson, Jr.	Charles L. Mollitor, Inc.	1950	79	440	8 & 6	420	420-440	Kmr	65	200	45	Irr	L.	98
99	1 mi. S of Medford Lakes on Tuckerton Rd.	George Aaron	J. Henry Robbins	1952	116	340	4	320	(none)	Kmr	50	50	--	D	L.	99
100	3 mi. SW of Medford Lakes, at Centennial Lake	William G. Freeman	do.	1955	65	275	4	260	(none)	Kmr	20	40	--	D	L.	100
101	2 mi. W of Medford Lakes, at Taunton Lake	Robert M. Dickson	do.	1951	66	242	4	222	(none)	Kmr	20	100	--	D		101
102	1 mi SE of Maple Shade on State Hwy. 73	Albert E. Crisp	Haines and Moore	1951	62	197	4	190	190-196	Kmr	58	20	5	D	Water temperature is 56°F. Water contains iron. L.	102
103	State Highway 38, Maple Shade	Savar Corporation	Charles L. Mollitor, Inc.	1955	62	166	6	155	155-166	Kmr	65	120	10	CI	Water used at a drive-in theater.	103
104	3 mi. E of Moorestown, on Marne Hwy.	William Haines, Jr.	Thomas C. Magee, Jr.	1954	52	55	4	47	47-53	Ket	21	10	10	D	A., L.	104
105	2 mi. SE of Moorestown	Roy W. Conrow	J. Henry Robbins	1952	37	205	6	195	195-205	Kmr	60	50	--	D	Water temperature is 56°F. L.	105
106	2 mi. E of Moorestown, on Union Mill Rd.	John McNicholas	Charles L. Mollitor, Inc.	1955	30	190	4	184	184-190	Kmr	22	15	3	D		106
107	3 mi. E of Moorestown, on Hartford Rd.	Thomas P. McGinnis	J. Henry Robbins	1952	46	276	4	256	256-276	Kmr	50	50	--	D	L.	107



Table 2.--Records of selected wells in Burlington County, N. J.--Continued

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
108	3 mi. E of Moorestown	Dominic Tusciano	Gus Hauser	1954	28	196	3	182	182-192	Kmr	48	8	—	D	Water temperature is 58°F. L.	108
109	do. on New Jersey Turnpike	New Jersey Turnpike	do.	1951	50	230	8	192	192-222	Kmr	48	970	140	CI	Used for air conditioning also.	109
110	4 mi. E of Moorestown	J. S. Coverly	Charles L. Mollitor, Inc.	1954	32	250	4	244	244-250	Kmr	30	50	10	D		110
111	3 mi. NW of Medford, on Hartford Rd.	Eva Diamond	John S. Wrobel	1951	73	202	4	190	190-200	Ket	48	10	6	D	L.	111
112	4 mi. SE of Moorestown, on Hartford Rd.	Anthony Palumbo	J. Henry Robbins	1949	66	348	4	328	328-348	Kmr	60	50	—	D		112
113	4 mi. SE of Moorestown	Samuel Dickter	Charles L. Mollitor, Inc.	1957	50	300	4	292	292-300	Kmr	60	40	5	D	L.	113
114	do.	Gertrude Freeman	do.	1953	38	262	4	256	256-262	Kmr	33	15	8	D		114
115	4 mi. W of Mount Holly, at Rancocas Woods	Walter H. Kopp	do.	1952	43	237	6	227	227-237	Kmr	41	20	46	D		115
116	2 mi. SE of Maple Shade, on Church Rd.	Charles J. Ruhle	J. Henry Robbins	1951	57	240	4	230	230-240	Kmr	60	50	—	D	L.	116
117	2 mi. SE of Maple Shade, on State Highway 73	Felix Oliveto	Charles L. Mollitor, Inc.	1953	63	223	4	216	216-223	Kmr	55	60	10	D	L.	117
118	do.	Ralph Vasturo	J. Henry Robbins	1950	68	119	6	100	(none)	Ket	25	10	—	D	A.	118
119	3 mi. E. of Moorestown, on Hartford Rd.	John D. Hallet	Charles L. Mollitor, Inc.	1956	32	244	4	238	238-244	Kmr	35	30	20	D		119
120	3 mi. E of Moorestown, at Masonville	William Breur	do.	1958	78	272	4	266	266-272	Kmr	90	40	10	D	L.	120
121	2 mi. NW of Marlton, on State Hwy 73	W. R. Fought	do.	1953	85	284	4	276	276-281	Kmr	89	15	10	D		121
122	2 mi. NE of Marlton	Joseph E. Rudderow	do.	1954	48	457	8 & 6	441	441-457	Kmr	56	450	125	Irr	Pump capacity is 450 gpm. L.	122
123	3 mi. NE of Marlton	do.	William C. Barr	before 1898	82	306	3	—	—	—	50	—	—	D	Mrs. Samuel Shreeve was the owner at the time the well was drilled.	123
124	do.	Walter E. Cotter, Jr.	Edward Robbins, Sr.	1955	77	384	4	379	379-384	Kmr	70	60	10	D	A.	124
125	4 mi. SE of Moorestown, on Hartford Rd.	Saul Grossman	Charles L. Mollitor	1951	40	105	4	98	98-105	Ket	22	10	18	D	A.	125
126	New Albany Rd., Moorestown	Pittsburgh Plate Glass Co.	Layne-New York Co., Inc.	1947	65	318	16 & 10	263	263-283	Kmr	65	517	94	Ind	A., L.	126
127	Mount Holly	Hollyford Ice & Cold Storage No. 1	do.	1946	10	500	10	470	470-500	Kmr	4	350	29	CI		127
128	do.	Dunlap's Carpet Works	C. G. Orcutt	before 1893	12	675	—	—	—	—	—	—	—	U	Well is abandoned. L.	128
129	do.	Aome Market	Charles L. Mollitor, Inc.	1956	62	467	8	456	456-467	Kmr	75	125	10	CI	Well deepened to 546' in 1960. New screen setting is 520-545'. Used for air conditioning. L.	129
130	4 mi. NE of Wrightstown, at Jacobstown	Jacobstown Baptist Church	Andrew White	1951	175	140	4	136	136-140	Kmr	50	15	—	CI	L.	130

Table 2.--Records of selected wells in Burlington County, N. J.--Continued

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
131	4 mi. NE of Wrightstown, at Ocean Co. line.	Jersey Central Power & Light Co.	Greenhalgh & Kaye	1956	88	121	6	110	110-121	Kmw	3	30	105	CI	A., L.	131
132	5 mi. N of Wrightstown	Edward H. Gancars	do.	1954	121	157	4	148	148-152	Ket	35	10	17	D	L.	132
133	6 mi. N of Wrightstown	Stanley Horner, Jr.	do.	1958	118	154	4	151	151-154	Ket	40	15	25	D		133
134	Wrightstown	Spartan Village, Inc.	Louis Bainbridge and Charles L. Mollitor, Inc.	1955	154	395	6	128	128-150	Kmw	26	40	19	U	L.	134
135	5 mi. N of Wrightstown	John Giberson	Greenhalgh & Kaye	1956	140	135	4	---	(none)	Kmw	42	10	13	D		135
136	Riverside	B. F. Goodrich Co. No. 2	Artesian Well Drilling Co.	1951	34	162	10	95	95-120	Kmr	24	458	19	Ind	A., L.	136
137	1 mi. SW of Riverside, at Cambridge	Dredge Harbor Yacht Basin	Charles L. Mollitor, Inc.	1947	10	80	--	76	(none)	Pzw	--	25	--	CI		137
138	Riverside	Riverside-Alloy Metal Co.	Ridpath & Potter	1938	18	800	6	120	(none)	Pzw	--	--	--	Ind	L.	138
139	do.	Riverside Industrial Center	Charles L. Mollitor, Inc.	1952	18	80	8	54	54-67	Kmr	12	250	28	Ind	Water temperature reported as 49°F(?)	139
140	1 mi. NW of Browns Mills	Jane Tabone	do.	1954	130	140	4	134	134-140	Tvt	38	10	8	CI	L.	140
141	2 mi. S of Pemberton	A. F. Brunt	do.	1955	83	210	4	204	204-210	Kmw	45	15	--	D	L.	141
142	1 mi. S of Pemberton	Clarence Adams	J. Henry Robbins	1949	72	195	4	175	(none)	Kmw	25	6	--	D	Farm use also.	142
143	Browns Mills	Mr. Reilley	Ridpath & Potter	before 1910	70-80	430	6	---	---	Ket	(flowing)	48	--	-	Pumped at a rate of 225 gpm also. L.	143
144	2 mi. E of Browns Mills, on Ridge Rd.	Joseph O'Neill	A. C. Schultes and Sons	1949	91	337	6	322	322-337	Kmw	29	110	27	D	L.	144
145	2 mi. SE of Browns Mills, at Country Lakes	Country Lakes, Inc.	Charles L. Mollitor, Inc.	1959	82	345	4	329	329-345	Kmw	23	80	10	PS	L.	145
146	2 mi. SE of Pemberton	Harvey Gower	Edward Robbins, Sr.	1949	68	250	6	250	(none)	Kmw	14	150	86	Irr		146
147	1 mi. E of Pemberton, on State Rte 530	John W. Goodman	Harvey Beebe, Sr.	1951	73	231	2	140	(none)	Kmw	15	30	--	D		147
148	4 mi. W of Wrightstown	Helis Stock Farm	Wm. Blaisdell & Orcutt Bros.	before 1893	approx. 70	715	-	---	---	Kmr	--	--	--	U	Abandoned. Formerly called Lorillard well. L.	148
149	4 mi. W of Wrightstown, at Jobstown	Kauffman & Minter, Inc.	Charles L. Mollitor, Inc.	1960	74	107	6	95	95-105	Ket	14	60	4	CI	A., L.	149
150	3 mi. SW of Wrightstown, at Juliustown	Milton L. Havens, Sr.	Gus Hauser	1953	115	137	4	117	(none)	Kmw	20	110	2	D	Water temperature is 56°F.	150
151	3 mi. SW of Wrightstown	Helis Stock Farm	Ridpath & Potter	1928	96	988	12 & 8	757	757-800	Kmr	--	120	4	O	Water contains iron. Formerly called Walker-Gordon Dairy well. L.	151
152	3 mi. N of Mount Holly	Ridge Stables	Charles L. Mollitor, Inc.	1957	41	230	6	220	220-230	Kmr	45	100	65	D	Farm use also. L.	152
153	2 mi. NW of Wrightstown	Frank J. Ondrushek, Sr.	Greenhalgh & Kaye	1958	182	256	4	253	253-256	Ket	96	15	30	D		153
154	5 mi. NE of Mount Holly, on U.S. Highway 206	Tallman Brothers, Inc.	J. Henry Robbins	1956	42	260	8 & 6	240	240-260	Kmr	60	40	--	CI		154
155	do.	do.	W. R. Clair	1952	42	52	2	47	47-52	Ket	6	10	4	CI	L.	155

Table 2.--Records of selected wells in Burlington County, N. J.--Continued

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
156	5 mi. SE of Mount Holly, at Vincentown	Calvin G. Smith	Harvey Beebe, Sr.	1949	37	271	2	171	(none)	Kmw	7	35	9	D		156
157	6 mi. SE of Mount Holly, on U.S. Hwy. 206	Ross Monroe	do.	1949	46	248	2	157	(none)	Kmw	11	25	—	D	L.	157
158	5 mi. NE of Medford Lakes, at Hampton Lakes	Hampton Lake Development Co.	Charles L. Mollitor, Inc.	1956	49	268	10	247	247-268	Kmw	6	300	43	PS	A., L.	158
159	3 mi. W of Pemberton, on Highways 38 and 206	Vincentown Diner	Edward Robbins, Sr.	1953	72	250	4	240	(none)	Ket	45	100	15	CI	L.	159
160	5 mi. NW of Chatsworth, on State Highway 72	Frank Mayo	Greenhalgh & Kaye	1953	140	69	4	65	65-69	Toh	37	15	9	CI	L.	160
161	3 mi. N of Mount Holly	Ronald Smith	J. W. Jenkins and Sons	1960	78	208	4	202	202-208	Kmr	90	25	60	D		161
162	2 mi. N of Mount Holly, on Oxmead Rd.	A. Marone	Charles L. Mollitor, Inc.	1956	40	265	6	254	254-265	Kmr	40	150	5	D	Supplies water to two houses. L.	162
163	3 mi. N of Mount Holly	Clifford Seriner	do.	1955	104	248	6	238	238-248	Kmr	96	30	14	D		163
164	3 mi. W of Mount Holly	Walter Jessup	do.	1950	80	241	4	234	234-241	Kmr	90	4	50	D	Water temp. is 54°F.	164
165	do.	Alexander Construction Co.	do.	1955	57	215	6	204	204-214	Kmr	60	50	20	U	Abandoned. L.	165
166	3 mi. NW of Mount Holly, on Woodlane Rd.	John S. Pew	do.	1957	79	228	4	222	222-228	Kmr	60	20	10	D	L.	166
167	3 mi. S of Burlington	Eugene D. Buzzelli	do.	1956	45	211	4	204	204-210	Kmr	48	20	6	D		167
168	2 mi. N of Mount Holly, on Burr Rd.	Burlington Co. Country Club	Edward Robbins, Sr.	1950	102	309	8 & 6	269	269-309	Kmr	100	400	20	Irr	Supplies water to the clubhouse also. A.	168
169	1 mi. E of Mount Holly, on Powell Rd.	John W. Hampton	Charles L. Mollitor, Inc.	1955	55	160	4	154	154-160	Ket	27	45	18	D	L.	169
170	2 mi. E of Mount Holly, on Powell Rd.	Donald C. Foote	do.	1954	52	160	4	154	154-160	Ket	15	15	7	D		170
171	2 mi. W of Pemberton, at Birmingham	Ionac Chemical Co.	A. C. Schultes & Sons	1960	30	921	12	490	490-521	Kmr	39	1001	104	Ind	A., L.	171
172	2 mi. E of Pemberton, on State Route 530	Burlington Co. Institution No. 1	Layne-New York Co., Inc.	1931	59	776	12 & 3	353	353-381	Ket	42	—	—	CI	A., L.	172
173	4 mi. N of Medford	Lester Jones	Charles L. Mollitor, Inc.	1953	41	159	6	148	148-159	Ket	16	110	34	D	Farm use also. A.	173
174	Mount Holly	Mount Holly Water Co. No. 1	Layne-New York Co., Inc.	1925	9	376	18 & 10	322	322-372	Kmr	12	700	101	PS	A.	174
175	1 mi. SW of Florence	National Gypsum Co. No. 1	do.	1956	22	167	12 & 8	118	118-128	Kmr	19	154	38	Ind	A.	175
176	7 mi. N of Wrightstown, on Ellisdale Rd.	R. H. Clark	J. Henry Robbins	1952	87	95	4	85	85-95	Ket	40	20	—	D	A.	176
177	2 mi. N of Wrightstown, at Sykesville	Allen Herbert	Greenhalgh & Kaye	1956	170	129	4	115	(none)	Kmw	40	10	20	D	A.	177
178	do.	William C. Rahilly	do.	1955	195	158	4	155	155-158	Kmw	58	10	5	D	L.	178
179	2 mi. E of Wrightstown, on State Route 528 Spur	Dix Concrete Co.	Leon Gager	1953	100	117	6	100	100-110	Kmw	(flowing)	60	68	CI	Flows 8 gpm 2 feet above land surface. Static water level is 8 feet above land surface	179
180	Wrightstown	Clifford C. Holden	Greenhalgh & Kaye	1957	133	150	4	—	(none)	Kmw	20	15	20	D		180
181	do.	Wrightstown Municipal Utility Authority		----	120	268	8	—	—	Ket	(flowing)	—	—	U	A., L	181

Table 2.--Records of selected wells in Burlington County, N. J.--Continued

Well no.	Location	Owner or name and number	Driller	Year completed	Altitude of land surface (feet)	Total depth drilled (feet)	Well diameter (inches)	Casing length (feet)	Screen setting (feet)	Water-bearing formation	Static water level below land surface (feet)	Yield (gpm)	Drawdown (feet)	Use	Remarks	Well no.
182	1 mi. W of Wrightstown	John L. Bond	Greenhalgh & Kaye	1953	143	123	-	--	-----	Kmw	13	15	7	D		182
183	2 mi. NW of Wrightstown	Charles B. Hutchinson	Charles L. Mollitor, Inc.	1960	180	134	4	124	124-134	Kmw	54	25	5	D	L.	183
184	2 mi. NE of Browns Mills	Fort Dix No. 10	William Stothoff Co., Inc.	1943	100	449	6	--	-----	Ket	26	60	102	PS	A.	184
185	4 mi. SE of Browns Mills, at Whitesbog	Joseph J. White, Inc.	Ridpath & Potter	1914	95	801	10 & 8	498	(none)	Ket	5	45	--	U	L.	185
186	do.	do.	J. W. Jenkins	1959	95	388	6 & 4	376	376-388	Kmw	--	150	--	CI	Pumping level is 90 feet. L.	186
187	4 mi. SE of Pemberton	Francis Bush	Edward Robbins, Sr.	1954	65	273	6	250	250-256	Kmw	28	400	38	Irr		187
188	Chatsworth	Anthony DeMarco Cranberry Plant	Bernard A. Leek	----	99	67	3	--	-----	Tch	6	--	--	D	A.	188
189	3 mi. NE of New Gretna, at Lake Absegami	Bass River State Forest		----	20	31	1 1/2	27	(none)	Tch	3	--	--	PS	A.	189
190	6 mi. NW of New Gretna	Green Bank State Forest Nursery			13	73	8 & 6	53	53-73	Tch	-	150	--	Irr	A.	190
191	New Gretna	Chester Allen			9	232	3	--	-----	Tkw	(flowing)	10	4	CI	Static water level is 4 feet above land surface. A.	191
192	Atsion	Lesson Small			51	88	1 1/2	--	-----	Tch	6	--	--	D	A.	192
193	3 mi. N of Atsion, at Indian Mills	Leah Prickett		----	77	60	---	--	-----	Tch	-	--	--	D	A.	193
194	4 mi. SW of Pemberton, near Vincentown	Vincentown Water Co.		1923	40	153	8	--	-----	Kmw	(flowing)	175	--	PS	Flows 28 gpm at land surface in the spring season. A.	194
195	2 mi. S of Mount Holly, at Lumberton	Electronic Parts Specialty Co.	J. Henry Robbins	1953	33	121	6 & 4	101	101-121	Ket	30	50	--	Ind	A.	195
196	2 mi. W of Medford Lakes, at Taunton Lakes	Taunton Lakes Water Co. No. 1	do.	1950	57	252	6	230	(none)	Kmw	25	300	--	U	Abandoned. A., L.	196
197	Beverly	National Waterproof Papers, Inc.	Charles L. Mollitor, Inc.	1955	38	61	16	41	41-6	Kmr	22	---	---	Ind	Used for cooling.	197
198	2 mi. SE of Riverton	Riverton-Palmyra Water Co. No. 10	A. C. Schultes & Sons	1961	79	308	12	243	243-281	Kmr	75	1051	95	PS	Drilled to weathered bed-rock. A., L.	198
199	2 mi NE of Moorestown, on Hartford Rd.	Moorestown Tp. Water Dept. No. 4	Layne-New York Co., Inc.	1959	59	360	12	298	298-338	Kmr	58	715	52	PS	L.	199
200	2 mi NE of Chatsworth	The Superior Zinc Corp.	Artesian Well Drilling Co.	1941	140	153	12	121	121-146	Tch	36	300	50	Ind		200
201	6 mi. E of Chatsworth on State Hwy 72	Cedar Bridge Fire Tower	J. Henry Robbins	1938	200	110	6	---	-----	Tch	61	---	--	O	L.	201
202	3 mi. W of Mount Holly	Inductotherm Corp.	Charles L. Mollitor, Inc.	1961	65	444	6	411	411-443	Kmr	80	300	16	Ind	L.	202
203	6 mi. S of Chatsworth, on State Route 563	Arthur Sooy	(Owner)	----	45	30	2	---	-----	Tch	-	---	--	D	Driven well. A.	203
204	1 mi. SW of Bordentown at Fieldsboro	Bzura Chemical Co., Inc.	A. J. Connelly, Inc.	1960	9	79	---	---	-----	Kmr	-	---	--	Ind	A.	204
205	3 mi. NE of Pemberton	Pemberton Tp. High School	Layne-New York Co., Inc.	1959	75	220	6	140	140-150	Kmw	-	---	--	CI	A.	205
206	3 mi. E of Moorestown on Marne Hwy	William Haines, Jr.	Artesian Well Drilling Co.	1958	55	238	8	200	200-238	Ket	-	400	--	Irr	A.	206
207	5 mi. NE of Mount Holly on U.S. Hwy 206	Harry Rupp		----	62	52	4	---	-----	Ket	-	---	--	CI	A.	207
208	2 mi. S of Moorestown	Mount Laurel Water Co.	A. C. Schultes & Sons	1961	30	595	12, 8	558	558-589	Kmr	32	548	77	PS	L.	208

Table 3. — Selected drillers' logs of wells in Burlington County, N. J.

Well no. 3, Burlington Water Department  
(Log by Layne-New York Co., Inc.)

	Thickness (feet)	Altitude 9 feet Depth (feet)
Topsoil . . . . .	1	1
Fill . . . . .	4	5
<b>Quaternary:</b>		
<b>Cape May Formation:</b>		
Coarse gravel . . . . .	5	10
Gravel and clay . . . . .	7	17
Gravel . . . . .	5	22
Gravel and sand . . . . .	2	24
White sand and gravel . . . . .	2	26
Brown sand and clay . . . . .	7	33
<b>Cretaceous:</b>		
<b>Magothy (?) and Raritan (?) Formations:</b>		
White sand and clay . . . . .	10	43
Sand, gravel, and sandy clay . . . . .	8	51
<b>Magothy and Raritan Formations:</b>		
Blue clay . . . . .	10	61
Clay, gravel, and sand . . . . .	3	64
Gravel, sand and clay . . . . .	7	71
Clay, gravel, and sand . . . . .	2	73
Sand, gravel, and clay . . . . .	12	85
Quartz sand and blue clay . . . . .	5	90



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 4, Burlington Water Department  
(Log by Artesian Well Drilling Co.)

	Altitude 19 feet	
	Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Sand and loam . . . . .	5	5
Dirty, brown sand . . . . .	5	10
Coarse sand . . . . .	5	15
Coarse sand and mud, some gravel . . . . .	5	20
Coarse sand and medium gravel . . . . .	5	25
Coarse sand and gravel . . . . .	5	30
Coarse sand and fine gravel . . . . .	10	40
Coarse sand and gravel . . . . .	6	46
Medium and large gravel . . . . .	2	48
Cretaceous:		
Magothy and Raritan Formations:		
Gray sandy clay . . . . .	1	49

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 9, Delaware River Water Co.  
(Log by A. C. Schultes & Sons)

	Altitude 21 feet	
	Thickness	Depth
	(feet)	(feet)
Quaternary:		
Cape May Formation:		
Fine, brown sand . . . . .	23	23
Sand, gravel, and clay . . . . .	3	26
Cretaceous(?):		
Magothy (?) and Raritan (?) Formations:		
Gray clay . . . . .	13	39
Stones . . . . .	2	41
Interbedded brown sand, gravel, and clay . . . . .	8	49
Coarse, brown sand . . . . .	5	54
Cretaceous:		
Magothy and Raritan Formations:		
Coarse, brown sand and white clay, some gravel . . . . .	21	75
Gravel and yellow clay . . . . .	7	82
Red, white and yellow clay . . . . .	8	90
Red and brown clay . . . . .	6	96
Yellow and white clay . . . . .	2	98
Stones, gravel, and clay . . . . .	7	105
Fine sand and gravel . . . . .	4	109
Fine sand and gravel, some white clay . . . . .	10	119
Brown and white sand, clay . . . . .	7	126
Coarse, brown sand and clay . . . . .	6	132
Yellow and white clay . . . . .	3	135
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock . . . . .	5	140

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 10, Delaware River Water Co.  
(Log by A. C. Schultes and Sons)

	Altitude 25 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
No sample .....	20	20
Cretaceous (?):		
Magothy (?) and Raritan (?) Formations:		
Hard clay .....	15	35
Cretaceous:		
Magothy and Raritan Formations:		
White and red clay .....	2	37
Soft, gray clay .....	3	40
Yellow sandy clay .....	14	54
Coarse, white sand .....	5	59
Medium to coarse sand, some ironstone .....	5	64
Yellow clay and fine sand .....	10	74
Pebbles, fine sand, and yellow clay - water-bearing .....	7	81
Medium to coarse sandy clay, ironstone .....	5	85
Coarse sand and gravel .....	5	91
Medium to coarse sand .....	5	96
Medium to coarse sand, clay, and stones .....	18	114
Fine sand, stones, and yellow clay .....	5	119
Medium to coarse sand .....	10	129
Brown clay .....	7	136
Red clay .....	6	142
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock .....	1	143

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 11, Evesham Township Water Department  
(Log by A. G. Dunphey)

	Altitude 115 feet Thickness (feet)	Depth (feet)
Topsoil . . . . .	9	9
Quaternary:		
Pensauken Formation:		
Yellow gravel . . . . .	5	14
Tertiary:		
Kirkwood Formation:		
Yellow sand . . . . .	15	29
Tertiary and Cretaceous:		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Black mud . . . . .	20	49
Marl . . . . .	25	74
Brown mud . . . . .	19	93
Mud and shells . . . . .	3	96
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Gray sand with abundance of water, somewhat irony . . . . .	10	106
White sand . . . . .	1	107
Black sand . . . . .	1	108
Whitish or gray sand . . . . .	17	125
Black sandy mud . . . . .	At	125
Marshalltown (?) Formation:		
Black clay . . . . .	Below	155
Englishtown (?) Formation:		
Black, muddy sandy . . . . .	At	185
Englishtown Formation:		
Sandy marl . . . . .	At	196
Light-colored sand . . . . .	At	208
Water-bearing, yellowish-white sand . . . . .	At	212

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 12, Evesham Township Water Department  
(Log by Layne-New York Co., Inc.)

	Altitude 89 feet	
	Thickness (feet)	Depth (feet)
Topsoil . . . . .	1	1
<b>Tertiary:</b>		
Hornestown Sand:		
Yellow sandy clay, red and blue marl . . . . .	15	16
<b>Cretaceous:</b>		
Navesink Formation:		
Black zeolite (?) sand and marl with clay streaks . . . . .	57	73
Mount Laurel Sand, Wenonah Formation, and Marshalltown Formation (undifferentiated):		
Fine to coarse sand with shells . . . . .	37	110
Blue clay with sand streaks . . . . .	55	165
Englishtown Formation:		
Fine to medium sand and gravel . . . . .	35	200
Woodbury Clay and Merchantville Formation (undifferentiated):		
Blue clay and sand with shells . . . . .	76	276
Blue and white clay with sand streaks . . . . .	84	360
Magothy and Raritan Formations:		
Fine to coarse, black and white sand . . . . .	33	393
Blue clay with streaks of fine sand . . . . .	22	415
Fine to medium sand . . . . .	11	426
Fine to medium sand with blue and white clay streaks. . . . .	74	500
Blue, red, and white clay . . . . .	12	512

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 13, Florence Township Water Department  
(Log by Layne-New York Co., Inc.)

	Altitude 25 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Quaternary:</b>		
Cape May Formation:		
Fine, brown sand . . . . .	4	4
Coarse, brown sand . . . . .	4	8
Gravel . . . . .	23	31
<b>Cretaceous:</b>		
Magothy and Raritan Formations:		
Blue and white clay . . . . .	2	33
Tough red clay . . . . .	7	40
Tough red and white clay . . . . .	44	84
Blue and white sandy clay . . . . .	10	94
Medium gray sand and white clay . . . . .	19	113
Coarse sand, red and white clay . . . . .	12	125
Coarse, brown sand . . . . .	13	138
Yellow and blue clay . . . . .	12	150
Coarse sand and clay . . . . .	8	158
<b>Early Paleozoic (?):</b>		
Wissahickon Formation:		
Mica rock . . . . .	5	163



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 15, Fort Dix  
(Log by Artesian Well Drilling Co.)

	Altitude 131 feet	
	Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
Kirkwood Formation:		
Fine, yellow clayey sand . . . . .	22	22
Gray clay and fine silty sand . . . . .	18	40
Green, clayey, glauconitic sand . . . . .	25	65
Manasquan and Vincentown Formations, Hornerstown Sand (undifferentiated):		
Yellowish-green, clayey, glauconitic sand . . . . .	41	106
Bright-green, clayey, and sandy glauconite with shells . . .	14	120
<b>Cretaceous:</b>		
Navesink Formation:		
Dark-gray glauconitic clay with shells . . . . .	21	141
Medium, yellowish-green, glauconitic sand . . . . .	21	162
Mount Laurel Sand and Wenonah Formation:		
Gray, clayey, glauconitic sand with fossils . . . . .	At	162
Fine, gray, clayey micaceous sand . . . . .	At	180
Marshalltown Formation:		
Gray clay and fine, micaceous sand . . . . .	At	274
Dark greenish-gray glauconitic clay with some fine micaceous sand . . . . .	At	275
Englishtown Formation:		
Brown clay with a trace of lignite . . . . .	At	299
Gray clay . . . . .	At	328
Gray clay . . . . .	At	352
Woodbury Formation:		
Gray, glauconitic, fossiliferous clay . . . . .	At	370
Gray, glauconitic, fossiliferous clay . . . . .	77	447
Merchantville Formation:		
Greenish-gray to dark gray glauconitic clay . . . . .	83	530
Magothy and Raritan Formations:		
Fine to medium glauconitic sand . . . . .	At	530
Very fine sand . . . . .	At	540
Very fine sand and some clay . . . . .	40	580
Fine sand . . . . .	42	622
Fine to coarse sand and gravel . . . . .	6	628
Fine to coarse, gray slightly clayey sand . . . . .	2	630
Gray sand . . . . .	At	630
Gray sand . . . . .	At	656
Very fine, light gray silty sand . . . . .	At	663
Fine to coarse, light gray sand, gravel and white clay . . . .	At	668
Fine, brown, micaceous sand . . . . .	At	715
Fine sand and gray clay . . . . .	At	720
Fine sand and clay . . . . .	150	870
Red, white, and gray clay . . . . .	160	1030
Coarse water-bearing sand . . . . .	21	1051

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 16, Fort Dix  
(Log by Layne-New York Co., Inc.)

	Altitude 148 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Tertiary:</b>		
Undifferentiated:		
Sand and gravel . . . . .	82	82
Clay and black sand . . . . .	59	141
<b>Cretaceous:</b>		
Undifferentiated:		
Hard, black sand . . . . .	45	186
Silt and clay . . . . .	45	231
Gray sand and gravel (?) . . . . .	25	256
Marshalltown and Englishtown Formations (undifferentiated):		
Silt, sand, and clay . . . . .	42	298
Woodbury Clay and Merchantville Formation (undifferentiated):		
Black clay and sand . . . . .	68	366
Clay and sand . . . . .	56	422
Hard clay and sand . . . . .	101	523
Magothy (?) and Raritan (?) Formations:		
Sandy clay . . . . .	33	556
Tough clay . . . . .	12	568
Sandy clay . . . . .	68	636
Hard sandy clay . . . . .	58	694
Tough clay with hard streaks . . . . .	48	742
Clay and gravel . . . . .	7	749
Gravel and hard clay . . . . .	23	772
Sand and hard clay . . . . .	50	822
Magothy and Raritan Formations:		
Clay with red streaks . . . . .	87	909
Clay, sand, gravel, and wood . . . . .	20	929
Medium, gray sand . . . . .	19	948
Tough clay . . . . .	21	969
Muddy sand . . . . .	12	981
Tough clay . . . . .	22	1003
Medium, gray sand . . . . .	21	1024
Tough clay . . . . .	10	1034
Sand, clay, and gravel . . . . .	16	1050
Coarse, gray sand and gravel . . . . .	36	1086
Clay . . . . .	10	1096

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 17, Fort Dix  
(Log by H. G. Richards, Geologist)

	Altitude 85 feet Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
<b>Kirkwood Formation:</b>		
Fine, yellow, slightly clayey sand . . . . .	10	10
Fine to coarse, buff and light gray sand . . . . .	10	20
Fine sand and gray clay . . . . .	10	30
Very fine, gray, clayey sand . . . . .	30	60
Very fine, gray sand and clay . . . . .	10	70
Black, lignitic, silty clay . . . . .	10	80
<b>Manasquan and Vincentown Formations (undifferentiated):</b>		
Green clay . . . . .	50	130
Green glauconitic clay . . . . .	10	140
Hard, light green clay . . . . .	20	160
Green clay . . . . .	30	190
Green glauconitic clay . . . . .	10	200
Green clay . . . . .	10	210
<b>Hornerstown Sand:</b>		
Dark green to gray glauconitic clay . . . . .	20	230
Green glauconitic clay . . . . .	10	240
Green glauconitic sandy clay . . . . .	20	260
<b>Cretaceous:</b>		
<b>Navesink Formation:</b>		
Gray glauconitic clay . . . . .	10	270
Gray sandy clay . . . . .	30	300
<b>Mount Laurel (?) Sand and Wenonah (?) Formation:</b>		
Silty, slightly micaceous sand and clay . . . . .	10	310
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Green silty, slightly clayey sand . . . . .	30	340
Gray silty sand and clay . . . . .	50	390
<b>Marshalltown Formation:</b>		
Gray, silty, sandy clay . . . . .	50	440
Fine, green to gray, clayey, glauconitic sand . . . . .	10	450
<b>Englishtown Formation:</b>		
Gray clay and sand . . . . .	10	460
Fine to medium, light green to gray, slightly glauconitic sand with shell fragments . . . . .	25	485

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 18, McGuire Air Force Base (Log by A. C. Schultes and Sons)		Altitude 128 feet
	Thickness (feet)	Depth (feet)
Topsoil . . . . .	1	1
Tertiary:		
Kirkwood Formation:		
Fine, yellow sand . . . . .	5	6
Fine, yellow sand and gravel . . . . .	9	15
Fine, gray sand . . . . .	7	22
Medium, yellow and gray sand . . . . .	13	35
Vincentown Formation:		
Fine, gray sand . . . . .	15	50
Tertiary and Cretaceous:		
Undifferentiated:		
Gray and white sand, black marl with shells . . . . .	151	201
Blue clay and marl . . . . .	63	264
Cretaceous:		
Englishtown Formation:		
Streaks of hardpan and sand . . . . .	7	271
Gray sand with lignite, streaks of hardpan and shells . . . . .	88	359
Woodbury Clay:		
Black clay with some sand . . . . .	16	375
Black clay, fine sand, and shells . . . . .	53	428
Merchantville Formation:		
Dark green clay, shells, and fine sand . . . . .	4	432
Clay and medium sand . . . . .	13	445
Black clay, medium sand, and shells . . . . .	57	502
Magothy (?) and Raritan (?) Formations:		
Green clay, fine sand, and black clay . . . . .	48	550
Magothy and Raritan Formations:		
Red and black clay with hard sand streaks . . . . .	38	588
Gray sand and white clay . . . . .	53	641
Gray sand with streaks of black clay . . . . .	24	665
Black clay . . . . .	5	670
Black clay and fine sand . . . . .	5	675
Fine, gray sand and black clay . . . . .	5	680
Fine to medium, gray sand and black clay . . . . .	41	721
Black clay and very fine sand . . . . .	19	740
Black clay, red clay, and shells . . . . .	12	752
Red and black clay, shells, and fine sand . . . . .	23	775
Fine, gray and white sand . . . . .	50	825
Black and red clay, wood, and black and gray sand . . . . .	55	880
Gray sand, red clay, and wood . . . . .	39	919
Tough, variegated clay . . . . .	31	950
Hard, red clay and coarse sand . . . . .	5	955
Hard, variegated clay . . . . .	5	960
Very hard, red clay . . . . .	24	984
Medium to coarse sand . . . . .	44	1028
Fine to coarse sand . . . . .	32	1060
Early Paleozoic (?):		
Wissahickon (?) Formation:		
Clay, sand, and weathered rock . . . . .	37	1097
Red clay . . . . .	3	1100

Well no. 19, McGuire Air Force Base (Log by A. C. Schultes and Sons)		Altitude 102 feet	
		Thickness	Depth
		(feet)	(feet)
<b>Tertiary:</b>			
Kirkwood Formation:			
Yellow sand . . . . .	5	5	
Fine, gray sand . . . . .	15	20	
Yellow clay, sand, and wood . . . . .	5	25	
Vincentown Formation and Hornerstown Sand (undifferentiated):			
Hard, gray and white sand . . . . .	65	90	
<b>Cretaceous:</b>			
Navesink Formation:			
Black marl with shells . . . . .	40	130	
Red clay . . . . .	1	131	
Mount Laurel Sand and Wenonah Formation:			
Fine, black and white sand . . . . .	89	220	
Marshalltown Formation:			
Fine, gray and black sand . . . . .	32	252	
Gray clay . . . . .	8	260	
Englishtown Formation:			
Fine to medium, gray, micaceous sand . . . . .	25	285	
Medium to coarse sand, black marl, and shells . . . . .	20	305	
Coarse sand, black marl, and shells . . . . .	25	330	
Coarse sand with shells . . . . .	20	350	
Woodbury Clay:			
Fine to coarse sand (?), marl, and shells . . . . .	25	375	
Fine to medium sand (?), shells, and marl . . . . .	35	410	
Medium to coarse sand (?), shells, and marl . . . . .	15	425	
Woodbury Clay and Merchantville Formation (undifferentiated):			
Medium to coarse sand (?), red clay, marl and shells . . . . .	38	463	
Merchantville Formation:			
Marl, shells, and red clay . . . . .	67	530	
Magothy and Raritan Formations:			
Clay, shells, and wood . . . . .	40	570	
Clay, shells, white clay, and wood . . . . .	15	585	
Coarse sand, gray clay, and wood . . . . .	47	632	
Medium to coarse sand, lignite, and gray clay . . . . .	46	678	
Red and white clay and marl . . . . .	29	707	
Layers of gravel, wood, and micaceous clay . . . . .	14	721	
White clay . . . . .	65	786	
White and gray clay and coarse sand . . . . .	44	830	
Variegated clay with hard streaks . . . . .	20	850	
Variegated clay . . . . .	70	920	
Gray sand, streaks of red and white clay, and wood . . . . .	62	982	
Medium to coarse sand and white clay . . . . .	6	988	
Coarse sand and red clay . . . . .	12	1000	
Sand and red clay . . . . .	20	1020	
Red clay . . . . .	8	1028	
Variegated sand . . . . .	72	1100	
<b>Early Paleozoic (?):</b>			
Wissahickon Formation:			
Weathered rock . . . . .	23	1123	

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 21, Lakes Water Co.  
(Log by J. Henry Robbins)

	Altitude 52 feet Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
Kirkwood Formation:		
Yellow sand and clay . . . . .	30	30
Kirkwood and Vincentown Formations (undifferentiated):		
Black clay . . . . .	70	100
<b>Tertiary and Cretaceous:</b>		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Green clay . . . . .	70	170
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Fine, green sand . . . . .	10	180
Coarse, green and white sand . . . . .	20	200



Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 22, Levitt and Sons Inc.  
(Log by A. C. Schultes and Sons)

	Altitude 29 feet Thickness (feet)	Depth (feet)
Topsoil . . . . .	5	5
Quaternary:		
Cape May Formation:		
Fine white sand with mica . . . . .	5	10
Fine to medium, brown sand . . . . .	15	25
Coarse sand and marl . . . . .	7	32
Gravel . . . . .	8	40
Coarse gravel and black clay . . . . .	5	45
Cretaceous:		
Merchantville Formation:		
Black clay . . . . .	20	65
Magothy and Raritan Formations:		
Fine to medium, brown sand and clay . . . . .	5	70
Fine, gray sand . . . . .	15	85
Fine to medium gray sand with mica and wood . . . . .	45	130
Medium to coarse sand . . . . .	10	140
Fine to medium sand . . . . .	35	175
White clay . . . . .	19	194
Fine gray sand . . . . .	51	245
Fine to medium, white sand . . . . .	10	255
White clay and fine, white sand . . . . .	10	265
Gray clay . . . . .	11	276
Gray clay, fine to coarse sand, and gravel . . . . .	3	279
No sample . . . . .	11	290
Gray clay . . . . .	4	294
Fine to coarse, gray sand . . . . .	29	323
Gray and red clay . . . . .	15	338
Medium, gray sand . . . . .	23	361
Early Paleozoic (?):		
Wissahickon Formation:		
Gray weathered rock . . . . .	2	363

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 24, Levitt and Sons Inc.  
(Log by A. C. Schultes and Sons)

	Altitude 28 feet Thickness (feet)	Depth (feet)
Topsoil .....	5	5
Quaternary:		
Cape May Formation:		
Yellow sandy clay .....	10	15
Cretaceous:		
Magothy and Raritan Formations:		
Red and white clay .....	39	54
Sand and gravel .....	5	59
Dark gray clay .....	5	64
Red clay with gravel .....	16	80
Dark gray clay .....	4	84
Medium, gray sand with clay .....	11	95
Fine, brown sand .....	15	110
Gravel with some stones .....	5	115
Medium, gray sand and gravel .....	45	160
Medium to coarse sand and gravel .....	40	200
Medium to coarse, white sand .....	30	230
Gravel and red clay .....	10	240
Gray silty clay .....	3	243
Red clay .....	4	247
White silty clay .....	8	255
Fine to medium sand with clay .....	18	273
Fine, brown sand with clay .....	9	282
Hardpan .....	1	283
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock .....	1	284

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 25, Levitt and Sons Inc.  
(Log by A. C. Schultes and Sons)

	Altitude 39 feet Thickness (feet)	Depth (feet)
Fill .....	5	5
Cretaceous:		
Magothy and Raritan Formations:		
Fine, gray sand .....	29	34
Gray clay .....	7	41
Fine, gray sand .....	8	49
Light gray clay .....	38	87
Fine, gray sand .....	11	98
Red clay .....	3	101
Fine, gray sand .....	58	159
Red clay .....	28	187
Fine to coarse, gray sand .....	6	193
Sand .....	3	196
Light gray clay .....	14	210
Fine, gray, silty sand .....	12	222
Gray clay .....	10	232
Fine gray sand .....	24	256
Gray sandy clay .....	12	268
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock .....	1	269

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 26, Levitt and Sons Inc.  
(Log by C. W. Lauman and Co. Inc.)

	Altitude 19 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Brown sand and gravel . . . . .	11	11
Cretaceous:		
Merchantville Formation:		
Gray to black, sandy silt with lignite . . . . .	22	33
Magothy and Raritan Formations:		
Fine to medium, gray sand . . . . .	14	47
Gray and white clay . . . . .	15	62
White, sandy clay . . . . .	4	66
White sand . . . . .	5	71
Clay . . . . .	5	76
Red and white clayey sand . . . . .	10	86
Brown sandy clay . . . . .	5	91
Fine, brown sand with some gravel . . . . .	5	96
Fine, white, clayey sand and gravel . . . . .	5	101
Fine, light brown sand . . . . .	10	111
Fine to coarse, brown sand with coarse gravel . . . . .	5	116
Fine clayey sand . . . . .	7	123
Gravel . . . . .	1	124
Fine, white sand . . . . .	2	126
Gray clay . . . . .	4	130
Fine to coarse, brown sand . . . . .	10	140
Fine to medium sand . . . . .	2	142
Clay . . . . .	1	143
Fine to coarse, brown sand . . . . .	9	152
Gravel . . . . .	1	153
Red and gray clay . . . . .	11	164
Fine, white sand . . . . .	12	176
Fine to medium, brown sand . . . . .	5	181
Fine to coarse, white sand . . . . .	6	187
White sandy clay . . . . .	5	192
Fine white sand . . . . .	15	207
Fine gray sand with some clay . . . . .	10	217
Fine to coarse, gray sand . . . . .	7	224
Gray clay . . . . .	3	227
Fine to coarse, gray sand . . . . .	6	233
Clay . . . . .	1	234
Fine to medium, gray sand . . . . .	6	240
Fine to coarse, brown and white sand . . . . .	5	245
Fine to coarse, gray sand and gravel . . . . .	10	255
Gray clay . . . . .	5	260
Red clay . . . . .	6	266
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock . . . . .	40	306

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 27, Levitt and Sons Inc.  
(Log by E. A. Hickok and D. W. Miller)

	Altitude 40 feet Thickness (feet)	Depth (feet)
<b>Quaternary:</b>		
<b>Pensauken Formation:</b>		
Fine sand (and fill dirt) .....	15	15
Fine, gray sand .....	3	18
Fine, yellow sand.....	2	20
<b>Cretaceous:</b>		
<b>Woodbury Clay:</b>		
Black micaceous clay .....	30	50
<b>Merchantville Formation:</b>		
Black to green glauconitic clay .....	60	110
<b>Magothy (?) and Raritan (?) Formations:</b>		
Black to green glauconitic clay and some fine sand.....	10	120
<b>Magothy and Raritan Formations:</b>		
Gray and white clay .....	30	150
Red and gray clay .....	10	160
Gray silty clay .....	20	180
Gray clay and medium gravel .....	10	190
Red and gray clay.....	30	220
Dark gray clay with wood .....	10	230
Gray clay .....	10	240
Fine to medium sand with gray clay stringers .....	20	260
Fine to medium sand with some mica .....	10	270
White silt .....	10	280
Sand, silt, and clay .....	10	290
Gray clay .....	10	300
Fine to medium sand .....	30	330
Red and gray clay .....	10	340
Gray silt.....	10	350
Fine to coarse sand with gravel and clay.....	30	380
Red and gray clay.....	35	415
Gray silt and clay.....	25	440
Fine to medium sand .....	10	450
Sand, gravel, and clay .....	10	460
Gray clay and silt.....	20	480
Medium to coarse sand and some gray clay .....	10	490
Gray clay with lignite .....	9	499
Medium gravel .....	10	509
Gray clay and fine sand.....	1	510
<b>Early Paleozoic (?):</b>		
<b>Wissahickon Formation:</b>		
Green to black, weathered, micaceous schist .....	16	526

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 28, Lumberton Light, Water and Sewage Co.  
(Log by Artesian Well Drilling Co.)

	Altitude 10 feet Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>		
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Sand.....	15	15
<b>Undifferentiated:</b>		
Green marl and clay .....	69	84
<b>Marshalltown Formation:</b>		
Clay with mica .....	20	104
<b>Englishtown Formation:</b>		
Black sand.....	20	124
Water-bearing sand .....	20	144
Sand, no water (?).....	5	149
<b>Woodbury Clay:</b>		
Hardpan .....	2	151
No record .....	49	200
Black clay and mud .....	46	246
<b>Merchantville Formation:</b>		
Green marl .....	35	281
Black marl .....	67	348
<b>Undifferentiated:</b>		
Marl and sand .....	12	360
<b>Magothy and Raritan Formations:</b>		
Pink clay .....	2	362
Water-bearing sand .....	38	400
White, plastic clay .....	4	404



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 29, Maple Shade Water Department  
(Log by James M. McCrae <sup>1/</sup>)

	Altitude 55 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Surface gravel .....	5	5
Yellow clay .....	8	13
Clay and ironstone .....	4	17
Cretaceous:		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Fossiliferous, black sand .....	3	20
Black clay .....	44	64
Magothy and Raritan Formations:		
Fine to coarse, gray sand and gravel .....	33	97
Greensand marl (?) .....	6	103
Fine, gray sand and coarse gravel .....	27	130
White clay .....	100	230
Red clay .....	10	240
Sand and clay .....	20	260
White sand .....	40	300
Fine, white sand with clay streaks .....	15	315
Coarse, white sand .....	35	350
Coarse, white gravel .....	20	370
Very coarse, white gravel with pebbles and boulders .....	5	375

<sup>1/</sup> Smock, 1894, p. 409-410

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 30, Maple Shade Water Department  
(Log by Artesian Well Drilling Co.)

	Altitude 10 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Clay and gravel .....	8	8
Cretaceous:		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Black marl .....	63	71
Magothy and Raritan Formations:		
Hard, dirty sand .....	32	103
Sand and gravel .....	10	113
Sand .....	4	117
Sand and gravel .....	9	126
Sand and coarse gravel .....	8	134
Clay .....	38	172
Dirty sand .....	3	175
Coarse, dirty sand .....	3	178
Coarse sand .....	4	182
Clay .....	7	189
Fine dirty sand .....	7	196
Clay .....	8	204
Fine, dirty sand .....	1	205
Sandy clay .....	2	207
Coarse sand and gravel .....	2	209
Coarse sand, gravel, and clay .....	9	218
Sandy clay .....	27	245
Coarse sand .....	7	252
Clay .....	20	272
No sample .....	10	282

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 31, Medford Water Co.  
(Log by Layne-New York Co., Inc.)

	Altitude 48 feet Thickness (feet)	Depth (feet)
Topsoil.....	1	1
Quaternary:		
Cape May Formation:		
Yellow sandy clay and gravel.....	11	12
Tertiary:		
Vincentown Formation and Hornerstown Sand (undifferentiated):		
Green marl with shells and streaks of clay.....	33	45
Tertiary and Cretaceous:		
Undifferentiated:		
Black sandy clay with shells .....	39	84
Fine sand with shells and streaks of gray clay .....	27	111
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Sandstone and shells.....	7	118
Fine, gray sand with shells and streaks of clay.....	19	137
Fine, black and white sand with shells .....	13	150
Marshalltown Formation:		
Black sandy clay with streaks of tough gray clay .....	25	175
Marshalltown and Englishtown Formations (undifferentiated):		
Gray sandy clay with shells .....	84	259
Englishtown Formation:		
Tough, gray clay .....	6	265
Woodbury Clay:		
Gray sandy clay and shells .....	52	317
Tough, gray clay .....	15	332
Fine, black and white sand with shells .....	26	358
Merchantville Formation:		
Black marl with shells and streaks of gray clay.....	77	435
Magothy and Raritan Formations:		
Hard, black and white sand with shells and streaks of gray clay .....	59	494
Tough, gray clay.....	6	500
Fine to medium, white sand .....	18	518
Tough, gray clay .....	2	520
Medium to coarse, gray sand .....	20	540
White sandy clay .....	12	552
Tough, gray clay .....	7	559
Fine sand .....	6	565
Tough, gray clay .....	3	568
Hard sand with streaks of gray clay .....	22	590

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 32, Moorestown Township Water Department  
(Log by Layne-New York Co., Inc.)

	Altitude 20 feet Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Blue sandy clay .....	13	13
Blue clay .....	112	125
Magothy and Raritan Formations:		
Fine, gray sand .....	23	148
Blue clay .....	8	156
Coarse, gray sand .....	33	189
White clay .....	60	249
Coarse, gray sand .....	56	305
Blue clay .....	15	320
Fine, gray sand .....	9	329
Blue clay .....	8	337
Coarse, gray sand .....	5	342
Coarse, gray sand and gravel .....	33	375
Yellow clay .....	7	382
Red clay .....	34	416
Sandy clay .....	6	422
Fine, gray sand .....	19	441
Coarse sand .....	22	463
Sand, gravel, and boulders .....	31	494
Blue clay .....	2	496
<b>Early Paleozoic (?):</b>		
Wissahickon Formation:		
Rock .....	1	497

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 34, Mount Holly Water Company  
(Log by John C. Smock <sup>2/</sup>)

	Altitude 20 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Surface material .....	12	12
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Dark gray micaceous sand with some green sand (glaucinite) .....	At	12
Marshalltown Formation:		
Marly clay .....	45	57
Englishtown Formation:		
Water-bearing, yellow sand .....	40	97
Woodbury Clay:		
Dark sandy clay .....	25	122
Clay with gravel .....	At	122
Dark marly clay .....	71	193
Merchantville Formation:		
Green sand (glaucinite) marl .....	42	235
Coarse, dark sandy clay .....	40	275
Magothy (?) and Raritan Formations (?):		
Pink, very clayey sand .....	15	290
Green and white clayey sand .....	10	300
Red clay with green sand .....	5	305
Clayey sand and green sand (glaucinite) .....	15	320
Magothy and Raritan Formations:		
Water-bearing, white lignitic sand .....	20	340
Fine, yellow gravel .....	2	342
Dark, micaceous sandy clay .....	8	350
White micaceous sand .....	18	368
Dark gray sand and fine gravel .....	2	370
Green clayey sand and fine gravel .....	24	394
Sandstone .....	At	394
Pink sandy clay .....	106	500
Red clay, less sandy .....	39	539
Water-bearing sand and gravel .....	23	562

<sup>2/</sup> Smock, 1901, p. 144-147

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 35, Pemberton Water Department  
(Log by A. L. Lyons)

	Altitude 80 feet	
	Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
Manasquan Formation:		
Yellow sandy clay .....	17	17
Vincentown Formation:		
Gray sand .....	22	39
Marl and sand .....	20	59
Undifferentiated:		
Green sand and clay .....	52	111
<b>Tertiary and Cretaceous:</b>		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Blue clay and marl .....	39	150
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Dark gray sand .....	35	185
Marshalltown Formation:		
Dark sandy clay .....	21	206

Well no. 37, Pemberton Township Water Department  
(Log by Thomas Magee)

	Altitude 112 feet	
Tertiary:		
Cohansey Sand:		
Sand .....	12	12
Kirkwood Formation:		
White sandy clay .....	7	19
Yellow sandy clay .....	40	59
White sand .....	15	74
Black marl .....	24	98
Manasquan and Vincentown Formations, Hornerstown Sand (undifferentiated):		
Green marl .....	24	122
Green sandy marl .....	56	178
Cretaceous:		
Navesink Formation:		
Hard, green marl .....	18	196
Green sandy marl .....	14	210
Green sandy marl with shells .....	65	275
Mount Laurel Sand and Wenonah Formation:		
Hard sand .....	27	302
Hard sand and marl with shells .....	1	303

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 38, Pemberton Township Water Department  
(Log by Layne-New York Co., Inc.)

	Altitude 93 feet Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
<b>Kirkwood Formation:</b>		
Coarse, brown sand .....	30	30
Brown sand and clay .....	15	45
Blue clay .....	60	105
<b>Manasquan and Vincentown Formations, Hornerstown Sand (undifferentiated):</b>		
Green clay .....	80	185
<b>Cretaceous:</b>		
<b>Navesink Formation:</b>		
Black clay and silt .....	92	277
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Coarse sand with shells .....	60	337
<b>Marshalltown Formation:</b>		
Fine sand and silt .....	20	357
Black sandy clay .....	45	402

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 39, Rancocas Woods Water Co. (Log by Charles L. Mollitor Inc.)		
	Altitude 50 feet Thickness (feet)	Depth (feet)
Topsoil .....	1	1
Cretaceous:		
Englishtown Formation:		
Sand .....	20	21
Water-bearing, fine sand .....	12	33
Brown sand and gravel .....	27	60
Woodbury Clay:		
Black marl .....	60	120
Merchantville Formation:		
Green marl .....	60	180
Magothy and Raritan Formations:		
White clay .....	7	187
Fine sand .....	48	235
Sand and gravel .....	13	248
Well no. 40, Riverton and Palmyra Water Co. (Log by A. C. Schultes and Sons)		
	Altitude 13 feet	
Topsoil .....	1	1
Quaternary:		
Cape May Formation:		
Brown silt .....	8	9
Brown sand and gravel .....	9	18
Cretaceous:		
Magothy and Raritan Formations:		
Gravel, sand, and red clay .....	2	20
Large gravel and some brown sand .....	14	34
Coarse, white sand .....	1	35
Clay, gravel, and stones .....	6	41
Yellowish-white clay with some gravel .....	6	47
Coarse, brown sand and clay .....	11	58
Yellowish-white clay, gravel, and coarse sand .....	1	59
Yellowish-white clay .....	3	62
Coarse, brown sand and gravel .....	7	69
White sand and clay .....	4	73
White sand, clay, and gravel .....	1	74
White sand with some gravel .....	5	79
Coarse, brown sand .....	2	81
Large stones and gravel .....	3	84
Yellow clay .....	9	93
Yellow micaceous clay .....	3	96
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock .....	4	100



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 42, Norman F. Scattergood  
(Log by Theodore Ziegenfus)

	Altitude 72 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
White sand .....	20	20
Cretaceous:		
Undifferentiated:		
Blue clay .....	125	145
Magothy and Raritan Formations:		
Gray sand .....	1	146
White clay .....	15	161
Fine sand .....	4	165
Clay .....	20	185
Fine sand .....	5	190
Fine and coarse sand .....	5	195
Blue clay .....	19	214
Blue clay and coarse sand .....	6	220
Clay .....	27	247
Clay and fine to coarse sand .....	6	253
Red clay .....	10	263
White clay .....	16	279
Coarse sand .....	1	280
White clay .....	35	315
Red clay .....	18	333
Coarse sand and gravel .....	2	335
Blue and white clay .....	13	348
Fine to coarse sand and gravel .....	14	362

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 44, National Guard Maintenance Shop  
(Log by Louis M. Bainbridge and Son)

	Altitude 83 feet	
	Thickness	Depth
	(feet)	(feet)
Topsoil .....	3	3
Quaternary:		
Cape May Formation:		
Yellow clay and sand .....	17	20
Cretaceous:		
Merchantville Formation:		
Green glauconitic marl .....	90	110
Magothy and Raritan Formations:		
Glauconitic marl and sand with wood .....	10	120
Black clay with pyrite and wood .....	5	125
Black sandy clay .....	5	130
Gray sand with black and white clay .....	15	145
Sand and gravel with black clay .....	10	155
Red and white clay .....	10	165
White clay .....	2	167
Gray sand and clay .....	6	173
Medium sand .....	12	185
Medium, gray sand .....	7	192
Black clay .....	7	199
Gray sand and black clay .....	1	200
Coarse sand and black clay .....	10	210
Coarse, gray sand .....	5	215
Coarse, white and gray sand .....	10	225
Red and white clay .....	5	230

Well no. 45, Hercules Powder Co.  
(Log by Layne-New York Co., Inc.)

	Altitude 22 feet	
Quaternary:		
Cape May Formation:		
Sand .....	6	6
Gravel and boulders .....	32	38
Cretaceous:		
Magothy and Raritan Formations:		
Red clay .....	39	77
White clay and sand .....	19	96
Coarse sand and fine gravel .....	24	120
Clay .....	7	127
Sand and gravel .....	8	135
Early Paleozoic (?):		
Wissahickon Formation:		
Mica rock .....	11	146

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 48, Edgar B. Heisler  
(Log by Charles L. Mollitor Inc.)

	Altitude 30 feet	
	Thickness	Depth
	(feet)	(feet)
Fill .....	1	1
Quaternary:		
Cape May Formation:		
Dry sand .....	18	19
Cape May (?) Formation:		
Yellow sand .....	29	48
Cretaceous:		
Magothy and Raritan Formations:		
Yellow sand and gravel .....	14	62
Brown sand .....	9	71
Muddy sand .....	12	83
Gray clay .....	9	92
Yellow clay .....	10	102
White clay .....	15	117
Yellow sand .....	3	120
Yellow sand and clay .....	13	133
Coarse sand and clay .....	4	137
White clay .....	20	157
Yellow sand and gravel .....	7	164
White clay .....	17	184
Sand and gravel .....	31	215

Well no. 49, Carroll Murphy  
(Log by Charles L. Mollitor Inc.)

	Altitude 35 feet	
Quaternary:		
Cape May Formation:		
Medium to coarse, brown sand and loam .....	6	6
Medium to coarse sand .....	10	16
Cretaceous:		
Magothy and Raritan Formations:		
Gray clay .....	2	18
White clay with sand .....	7	25
Red clay with sand .....	5	30
Red and gray clay .....	8	38
Gray clay with sand and gravel .....	3	41
Brown sandstone .....	10	51
Medium to coarse, gray sand .....	25	76
Fine, white sand .....	6	82
White clay and white sand .....	1	83

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 50, Masonic Home  
(Log by American Well Drilling Co.)

	Altitude 79 feet	
	Thickness	Depth
	(feet)	(feet)
Quaternary:		
Pensauken Formation:		
Yellow sandy clay .....	25	25
Cretaceous:		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Black silty clay .....	72	97
Magothy and Raritan Formations:		
Dirty, fine sand .....	12	109
Red clay .....	42	151
Dirty medium sand .....	9	160
Clean, fine to coarse, gray sand .....	13	173
Fine to coarse, gray sand with wood fragments .....	1	174
Hardpan .....	1	175
Clay .....	1	176
Clean, medium sand .....	8	184
Clean, coarse sand and fine gravel .....	2	186
Clay .....	1	187
Clean, fine gravel .....	5	192
Clean, medium to coarse gravel .....	5	197
Clay .....	1	198
Clean, coarse sand .....	4	202
Clay .....	1	203
Clean, coarse sand and fine gravel .....	30	233
Red and white clay .....	49	282
White sand .....	12	294
White clay .....	22	316
Water-bearing sand and gravel .....	20	336
Red clay .....	59	395

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 51, William A. Donald  
(Log by Paul Steffens)

	Altitude 42 feet	
	Thickness (feet)	Depth (feet)
Loamy topsoil .....	6	6
Quaternary:		
Cape May Formation:		
Clay .....	5	11
Dirty sand .....	18	29
Cretaceous (?):		
Magothy (?) and Raritan (?) Formations:		
Sand and gravel .....	10	39
Cretaceous:		
Magothy and Raritan Formations:		
Gray clay .....	1	40
Sand and gravel .....	15	55
White clay .....	2	57
Coarse sand .....	3	60
Fine gravel .....	11	71
Gray clay .....	1	72
Sand and gravel .....	7	79
Gray clay .....	4	83
Fine sand .....	2	85
White clay .....	1	86
Sand and fine gravel .....	20	106
White clay .....	5	111

Well no. 54, Cherry Hill Inn Farm  
(Log by Charles L. Mollitor Inc.)

	Altitude 72 feet	
Topsoil .....	6	6
Cretaceous:		
Englishtown Formation:		
Brown loam .....	4	10
Loam with gray clay .....	8	18
Sand and red clay .....	8	26
Woodbury Clay and Merchantville Formation (undifferentiated):		
Black and gray clay .....	85	111
Green marl with hardpan .....	27	138
Gray clay with hardpan .....	32	170
Magothy and Raritan Formations:		
Hardpan, gravel, and fine, gray sand .....	21	191
Water-bearing, fine, gray sand .....	17	208

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 55, Frank Pino  
(Log by Edward Robbins)

	Altitude 63 feet	
	Thickness (feet)	Depth (feet)
<b>Cretaceous (?):</b>		
Merchantville (?) Formation:		
Sand .....	30	30
Magothy (?) and Raritan (?) Formations:		
Light-gray clay .....	50	80
<b>Cretaceous:</b>		
Magothy and Raritan Formations:		
Sand .....	1	81
Clay .....	1	82
No sample .....	8	90
Water-bearing sand .....	10	100
Sand and clay .....	49	149
Water-bearing sand and clay streaks .....	20	169
Clay and sand .....	31	200
Red clay .....	7	207
White sand and gravel .....	20	227

Well no. 56, National Gypsum Co.  
(Log by Layne-New York Co., Inc.)

	Altitude 22 feet	
	Thickness (feet)	Depth (feet)
Topsoil .....	1	1
<b>Quaternary:</b>		
Cape May Formation:		
Red sand and clay .....	9	10
Brown sand and gravel .....	11	21
Coarse sand and gravel .....	10	31
<b>Cretaceous:</b>		
Magothy and Raritan Formations:		
White and yellow sandy clay .....	6	37
Coarse, yellow sand and gravel with streaks of yellow clay .....	11	48
Yellow and white sandy clay with some gravel .....	12	60
Red and white clay .....	28	88
White sandy clay .....	22	110
Brown sand and coarse gravel .....	14	124
White clay .....	2	126
Coarse sand .....	4	130
White and yellow clay .....	10	140
<b>Early Paleozoic (?):</b>		
Wissahickon Formation:		
Mica rock .....	17	157

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 58, Charles Heal, Jr.  
(Log by Charles L. Mollitor Inc.)

	Altitude 50 feet	
	Thickness (feet)	Depth (feet)
<b>Quaternary:</b>		
Cape May Formation:		
Sand .....	1	1
Dry sand .....	4	5
Sand .....	14	19
Gravel .....	2	21
<b>Cretaceous:</b>		
Merchantville Formation:		
Black marl .....	33	54
Magothy and Raritan Formations:		
Sand .....	17	71
Gravel .....	9	80

Well no. 61, Harrisville  
(Log by Mahlon Broon <sup>3/</sup>)

	Altitude 15 feet	
Tertiary:		
Cohansey Sand:		
Sand. ....	77	77
Sand. ....	8	85
Blue mud .....	13	98
Clay (with pyrite?) .....	10	108
Marly mud .....	16	124
Strata with wood (?) .....	7	131
Mud with shells .....	15	146
Hard sediments .....	50	196
Water-bearing sand .....	35	231
Dark, slushy sand .....	14	245
Yellow sand .....	16	261
Coarse, red sand .....	45	306
Dark sand .....	12	318
White clay .....	13	331
Kirkwood Formation:		
Water-bearing, green marl .....	37	368
Slate stone (?) .....	7	375

<sup>3/</sup> Smock, 1893, p. 288-289

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 62, New Jersey State Reformatory  
(Log by Layne-New York Co., Inc.)

	Altitude 97 feet	
	Thickness	Depth
	(feet)	(feet)
Topsoil .....	1	1
Quaternary:		
Pensauken Formation:		
Red sand, gravel, and clay .....	12	13
Cretaceous:		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Tough, black clay .....	22	35
Soft, green clay .....	40	75
Soft, black clay .....	40	115
Magothy (?) and Raritan (?) Formations:		
Soft, gray clay .....	25	140
Magothy and Raritan Formations:		
Fine, white sand.....	35	175
Sand, gravel, and clay .....	15	190
Soft clay .....	15	205
Coarse, white sand .....	15	220
Soft clay .....	10	230
Sand with streaks of clay .....	22	252
Clay .....	12	264
Sand and clay .....	12	276
Sand .....	30	306
Red clay .....	39	345
Coarse sand .....	52	397
Clay .....	25	422
Coarse sand .....	22	444
Clay .....	10	454



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 66, Dana Packing Co.  
(Log by Greenhalgh and Kaye)

	Altitude 208 feet	
	Thickness	Depth
	(feet)	(feet)
Quaternary (?):		
Bridgeton (?) Formation:		
Yellow gravel .....	38	38
Tertiary and Cretaceous:		
Hornerstown and Red Bank Sands, Navesink Formation		
(undifferentiated):		
Dense, brown clay .....	22	60
Green and black marl .....	46	106
Cretaceous (?):		
Mount Laurel (?) Sand and Wenonah (?) Formation:		
Green clay .....	24	130
Dark gray clay .....	25	155
Cretaceous:		
Marshalltown Formation:		
Clay and fine sand .....	95	250
Englishtown Formation:		
Coarse, yellow sand .....	26	276

Well no. 67, Francis D. Gray  
(Log by Louis M. Bainbridge and Son)

	Altitude 92 feet	
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Yellow sand and clay .....	30	30
Marshalltown Formation:		
Green clay .....	15	45
Gray clay .....	25	70
Englishtown Formation:		
Gray sand with pyrite .....	16	86
Gray clay .....	5	91
Gray sand and clay .....	10	101
Englishtown Formation, Woodbury Clay and Merchantville		
Formation (undifferentiated):		
Gray clay .....	94	195
Green marl .....	40	235
Gray clay .....	25	260
Grayish-green clay .....	5	265
Black clay .....	14	279
Magothy and Raritan Formations:		
Gray sand and clay .....	34	313
Black clay .....	At	313
Gray sand .....	4	317

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 68, John Marincas  
(Log by Greenhalgh and Kaye)

	Altitude 97 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Quaternary and Cretaceous:</b>		
Pensauken and Englishtown Formations (undifferentiated):		
Yellow gravel and clay .....	38	38
<b>Cretaceous:</b>		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Dense, black clay .....	12	50
Gray and black clay .....	70	120
Dark green marl with shells .....	45	165
Black clay .....	15	180
<b>Magothy and Raritan Formations:</b>		
Clay and sand .....	5	185
Coarse sand and sandstone with wood .....	23	208

Well no. 70, Hoeganaes Sponge Iron Co.  
(Log by Layne-New York Co., Inc.)

	Altitude 35 feet	
<b>Quaternary:</b>		
Cape May (?) Formation:		
Sand .....	10	10
<b>Cretaceous:</b>		
Magothy and Raritan Formations:		
Sand and sandy clay .....	6	16
Muddy sand .....	5	21
Sand, gravel, and clay .....	5	26
Sand .....	8	34
Clay .....	11	45
Sand, gravel, and clay .....	14	59
Yellow sand, gravel, and clay .....	13	72
Gray sand and white clay .....	10	82
Gravel, sand, and clay .....	16	98
Muddy gravel and sand .....	4	102
Gravel with streaks of clay .....	7	109
Coarse sand and gravel .....	26	135

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 71, Eastampton School  
(Log by Charles L. Mollitor, Inc.)

	Altitude 73 feet Thickness (feet)	Depth (feet)
Loamy topsoil .....	6	6
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Sand .....	8	14
Undifferentiated:		
Marl .....	117	131
Englishtown Formation:		
Sand .....	15	146
Clay .....	23	169
Sand .....	13	182
Woodbury Clay and Merchantville Formation (undifferentiated):		
Marl .....	161	343
Magothy and Raritan Formations:		
Sand .....	4	347
Red and white clay .....	22	369
Sand and hardpan .....	5	374
Gray clay .....	21	395
Muddy sand .....	6	401
Sand .....	16	417

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 72, J. A. Methvin  
(Log by Paul Steffens)

	Altitude 39 feet Thickness (feet)	Depth (feet)
Loamy topsoil .....	11	11
Cretaceous:		
Mount Laurel Sand and Wenonah and Marshalltown Formations (undifferentiated):		
Black sandy marl .....	87	98
Englishtown Formation:		
Water-bearing, fine sand .....	6	104
Dark gray clay .....	3	107
Water-bearing, coarse sand .....	5	112
Hardpan and dirty, coarse sand .....	4	116
Black marl .....	15	131
Fine sand with wood .....	4	135
Black marl .....	13	148
Hard, fine sand .....	2	150
Woodbury Clay and Merchantville Formation (undifferentiated):		
Very stiff, black marl .....	46	196
Marl with streaks of hardpan .....	44	240
Green marl .....	9	249
Hardpan .....	4	253
Streaks of clay and marl .....	12	265
Sandy marl with thin streaks of sand .....	17	282
Hardpan .....	3	285
Marl .....	14	299
Gray clay .....	5	304
Magothy and Raritan Formations:		
Hard sand .....	4	308
Red clay .....	4	312
Dirty, black sand .....	5	317
Gray and red clay .....	16	333
Hard, coarse sand .....	5	338
Black clay .....	14	352
Sand .....	11	363
Red clay .....	1	364

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 73, Byron T. Roberts Farms, Inc.  
(Log by A. C. Schultes and Sons)

	Altitude 93 feet Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>		
<b>Navesink Formation:</b>		
Fine to medium, brown sand .....	6	6
Brown marly sand .....	21	27
Gray marl .....	5	32
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Fine, gray sand .....	53	85
<b>Marshalltown Formation:</b>		
Soft, gray clay .....	51	136
Gray, laminated clay .....	4	140
<b>Englishtown Formation:</b>		
Fine, gray sand .....	6	146
Hard, gray clay .....	30	176
Fine sand and clay .....	17	193
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Gray clay .....	79	272
Hard, gray clay .....	43	315
<b>Magothy and Raritan Formations:</b>		
Fine, gray sand .....	5	320
Hard, gray clay .....	2	322
Fine to coarse, gray sand .....	32	354
Clay .....	6	360
Fine sand .....	22	382
Hard, gray clay .....	24	406

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 74, David D. Griscom  
(Log by J. Henry Robbins)

	Altitude 92 feet	
	Thickness (feet)	Depth (feet)
Quaternary:		
Pensauken Formation:		
Yellow clay .....	20	20
Cretaceous:		
Navesink Formation:		
Black mud .....	50	70
Undifferentiated:		
Green marl .....	80	150
Hard, black clay .....	120	270
Green clay .....	30	300
Magothy (?) and Raritan (?) Formations:		
Black and gray clay .....	20	320
Magothy and Raritan Formations:		
White sand .....	20	340

Well no. 76, Earl R. Blyler  
(Log by Charles L. Mollitor, Inc.)

	Altitude 39 feet	
	Thickness (feet)	Depth (feet)
Topsoil .....	1	1
Quaternary:		
Cape May Formation:		
Medium to coarse, yellow sand .....	7	8
Medium, brown sand .....	5	13
Water-bearing, brown sand .....	7	20
Coarse, brown sand and gravel .....	8	28
Very coarse sand and gravel .....	10	38
Cretaceous:		
Magothy and Raritan Formations:		
Grayish-white sandy clay .....	10	48
Fine, yellow sand, clay, and hardpan .....	2	50
Fine, yellow sand .....	11	61
Medium to coarse, dark yellow sand .....	7	68
White clay with iron-stained sand .....	2	70
White clay, sand, and hardpan .....	6	76
Water-bearing, medium to coarse, yellow sand .....	6	82
Yellowish-white sand .....	8	90
Medium to coarse, light tan sand .....	2	92
Coarse, light tan sand with clay .....	10	102
Water-bearing, medium to coarse, light tan sand and gravel .....	15	117

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 78, Amico Sand and Gravel Co.  
(Log by Paul Steffens)

	Altitude 15 feet Thickness (feet)	Depth (feet)
<b>Quaternary:</b>		
<b>Cape May Formation:</b>		
Sand .....	3	3
Coarse gravel .....	3	6
<b>Cretaceous:</b>		
<b>Magothy (?) and Raritan (?) Formations:</b>		
Black clay .....	9	15
Dirty sand .....	3	18
Sand and gravel .....	12	30
Gravel .....	6	36
Clay .....	13	49
<b>Early Paleozoic (?):</b>		
<b>Wissahickon Formation:</b>		
Soft rock .....	26	75

Well no. 80, Holy Cross High School  
(Log by Charles L. Mollitor Inc.)

	Altitude 82 feet	
Topsoil .....	2	2
<b>Cretaceous:</b>		
<b>Magothy and Raritan Formations:</b>		
Yellow sand .....	4	6
Brown sand .....	20	26
White clay .....	6	32
Red clay .....	11	43
Gray and white clay .....	6	49
Gray clay .....	22	71
Dirty, gray sand .....	48	119
White clay .....	19	138
White sand .....	7	145
White clay .....	8	153
Sand and gravel .....	21	174

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 81, Holiday Lake  
(Log by Charles L. Mollitor, Inc.)

	Altitude 25 feet Thickness (feet)	Depth (feet)
Quaternary (?):		
Cape May (?) Formation:		
Yellow sand .....	9	9
Sand .....	12	21
Gravel .....	4	25
Cretaceous:		
Magothy and Raritan Formations:		
Yellow sand .....	21	46
Sand .....	15	61
White clay .....	4	65
White sand .....	7	72
Gray clay .....	13	85
White clay .....	14	99
White sand .....	8	107
Muddy sand .....	6	113
Hard clay .....	6	119
Muddy sand .....	10	129
White sand .....	9	138
White clay .....	4	142
White sand .....	4	146
White clay .....	10	156
Fine sand .....	5	161
White clay .....	6	167
Dirty sand .....	3	170
Sand and gravel .....	7	177
White clay .....	10	187
Sand and gravel .....	11	198



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 88, John H. Cook  
(Log by Charles L. Mollitor Inc.)

	Altitude 115 feet	
	Thickness	Depth
	(feet)	(feet)
Topsoil .....	2	2
Quaternary:		
Pensauken Formation:		
Brown sandy loam .....	6	8
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Hard, coarse, brown sand with some clay .....	7	15
Hard, brown sandy clay .....	16	31
Marshalltown (?) Formation:		
Green marl .....	14	45
Black clay .....	25	70
Marshalltown Formation:		
Black to green marl .....	6	76
Gray sandy clay .....	16	92
Black clay with some fine sand .....	40	132
Englishtown Formation:		
Water-bearing, green sand with fossils and wood .....	16	148

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 89, Turnpike Junction Industrial Park  
(Log by A. C. Schultes and Sons)

	Altitude 60 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Cretaceous:</b>		
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Brown and gray clay .....	11	11
Dark gray clay .....	88	99
Gray clay .....	7	106
Dark gray clay .....	4	110
<b>Magothy and Raritan Formations:</b>		
Fine, gray sand .....	49	159
Dark gray clay .....	7	166
Fine to medium, gray sand .....	15	181
Dark gray clay .....	8	189
Fine to medium, gray sand .....	12	201
Dark gray clay .....	4	205
Fine, gray sand .....	7	212
Red clay .....	3	215
Fine to medium, gray sand .....	24	239
Red clay .....	5	244
Fine to medium, gray sand .....	5	249
Red clay .....	4	253
Fine, gray sand .....	2	255
Red clay .....	4	259
Fine to medium, gray sand .....	5	264
Red clay .....	21	285
Fine, gray sand .....	28	313
Red clay .....	3	316
Fine to coarse, gray sand .....	18	334
Red clay .....	6	340

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 90, Franklin Tallman  
(Log by Charles L. Mollitor Inc.)

	Altitude 82 feet Thickness    Depth (feet)        (feet)	
<b>Quaternary:</b>		
Cape May Formation:		
Fine, brown sand and loam .....	6	6
<b>Cretaceous (?):</b>		
Englishtown (?) Formation:		
Hardpan .....	1	7
Medium, yellow sand .....	7	14
Water-bearing, medium sand with a thin layer of hardpan ....	7	21
Brown clay .....	9	30
<b>Cretaceous:</b>		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Very sticky, gray clay .....	42	72
Soft, gray clay .....	18	90
Green marl .....	18	108
Light gray clay and hardpan .....	14	122
Black clay .....	18	140
Fine, gray sandy clay with mica .....	20	160
Hard, black clay .....	20	180
<b>Magothy and Raritan Formations:</b>		
Red and white sandy clay .....	7	187
Sand .....	8	195

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 91, Ed Phares  
(Log by Louis Bainbridge)

	Altitude 92 feet Thickness (feet)	Depth (feet)
Topsoil .....	18	18
Cretaceous:		
Woodbury Clay:		
Blue and brown clay .....	27	45
Undifferentiated:		
White clay .....	50	95
Merchantville Formation:		
Green clay .....	30	125
Magothy and Raritan Formations:		
White clay .....	99	224
Red clay .....	6	230
No sample .....	27	257
Brown and white clay .....	10	267
White clay .....	73	340
Black sand .....	2	342
White clay .....	3	345
Gray and white clay with wood .....	5	350
Gray and white clay .....	20	370
Blue clay .....	10	380
White sand .....	2	382
Soft, white clay .....	28	410
Red and white clay .....	1	411
Red clay .....	11	422
Hardpan .....	1	423
Red and white clay with hardpan .....	82	505

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 94, North Burlington Co. Regional Jr.-Sr. High School  
(Log by Russell D'Agostino)

	Altitude 83 feet Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>		
<b>Marshalltown Formation:</b>		
Medium, brown sand and clay .....	30	30
Brown clay .....	5	35
<b>Englishtown Formation:</b>		
Medium, brown sand .....	17	52
Brown sand and clay .....	11	63
<b>Undifferentiated:</b>		
Dark clay .....	92	155
Gray gravel and clay .....	5	160
Dark clay .....	60	220
<b>Merchantville (?) Formation:</b>		
Dark clay with some stones .....	12	233
<b>Magothy and Raritan Formations:</b>		
Sand .....	24	256
Fine, brown sand .....	8	264
Clay .....	12	276
Fine to coarse sand .....	34	310
Sand and clay .....	10	320
Sand and clay with wood .....	At	320
Sand and clay .....	16	336
Sand .....	6	342
Dark clay .....	8	350

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 95, Ralph R. Parker  
(Log by W. R. Clair)

	Altitude 62 feet	
	Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>		
<b>Englishtown Formation:</b>		
Clayey sand .....	15	15
Water-bearing sand .....	10	25
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Black clay .....	105	130
Water-bearing, green marl .....	20	150
<b>Undifferentiated:</b>		
Black clay .....	130	280
<b>Magothy and Raritan Formations:</b>		
Fine to medium sand .....	5	285
White clay .....	75	360
Fine sand .....	10	370
Coarse sand .....	10	380

Well no. 97, Peter Martin  
(Log by John S. Wrobel)

	Altitude 62 feet	
<b>Tertiary and Cretaceous:</b>		
Vincentown Formation, Hornerstown Sand, and Navesink Formation (undifferentiated):		
Marl .....	100	100
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Sandstone .....	20	120
Water-bearing, gray sand .....	6	126

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 98, William E. Johnson Jr.  
(Log by Charles L. Mollitor Inc.)

	Altitude 79 feet	
	Thickness	Depth
	(feet)	(feet)
Topsoil .....	2	2
Tertiary:		
Hornerstown Sand:		
Dry sand .....	5	7
Water-bearing, green sand .....	36	43
Cretaceous:		
Navesink Formation:		
Sandy marl .....	32	75
Mount Laurel Sand and Wenonah Formation:		
Fine sand .....	50	125
Marshalltown Formation:		
Black sandy marl: .....	70	195
Englishtown Formation and Woodbury Clay (undifferentiated):		
Green marl .....	20	215
Black sandy marl .....	25	240
Gray marl .....	5	245
Hardpan and clay .....	10	255
Sandy marl .....	8	263
Hardpan and clay .....	2	265
Black marl .....	73	338
Merchantville Formation:		
Green marl .....	37	375
Black marl .....	12	387
Green marl .....	13	400
Magothy and Raritan Formations:		
Fine sand .....	5	405
Clay .....	15	420
Sand .....	20	440

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 99, George Aaron  
(Log by J. Henry Robbins)

	Altitude 116 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Tertiary:</b>		
Kirkwood Formation:		
Yellow clay and fine sand .....	80	80
<b>Tertiary and Cretaceous:</b>		
Manasquan and Vincentown Formations, Hornerstown Sand and		
Navesink Formation (undifferentiated):		
Black clay .....	70	150
Green clay .....	170	320
<b>Cretaceous:</b>		
Mount Laurel (?) Sand and Wenonah (?) Formation:		
Green sand .....	20	340

Well no. 100, William G. Freeman  
(Log by J. Henry Robbins)

	Altitude 65 feet	
<b>Tertiary:</b>		
Kirkwood Formation:		
Yellow clay and sand .....	50	50
<b>Tertiary and Cretaceous:</b>		
Manasquan and Vincentown Formations, Hornerstown Sand, and		
Navesink Formation (undifferentiated):		
Black clay .....	100	150
Green clay .....	110	260
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Green and white sand .....	15	275



Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 102, Albert E. Crisp  
(Log by Haines and Moore)

	Altitude 62 feet	
	Thickness (feet)	Depth (feet)
Quaternary:		
Pensauken Formation:		
Sand .....	8	8
Cretaceous:		
Marshalltown Formation:		
Black clay .....	24	32
Englishtown Formation, Woodbury Clay and Merchantville Formation (undifferentiated):		
Fine, black sand .....	7	39
Black clay .....	86	125
Sand, marl, and clay .....	24	149
Black clay .....	28	177
Magothy and Raritan Formations:		
Sand .....	20	197

Well no. 104, William Haines, Jr.  
(Log by Thomas C. Magee, Jr.)

	Altitude 52 feet	
Quaternary:		
Cape May Formation:		
Sand and loam .....	3	3
Sand and gravel .....	19	22
Cretaceous:		
Marshalltown Formation:		
Green marl .....	3	25
Englishtown Formation:		
White sand .....	30	55

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 105, Roy W. Conrow  
(Log by J. Henry Robbins)

	Altitude 37 feet	
	Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Yellow clay .....	10	10
Cretaceous:		
Englishtown Formation, Woodbury Clay, and Merchantville Formation (undifferentiated):		
Black clay .....	100	110
Green marl .....	80	190
Magothy and Raritan Formations:		
White clay and sand .....	5	195
Sand .....	10	205

Well no. 107, Thomas P. McGinnis  
(Log by J. Henry Robbins)

	Altitude 46 feet	
Cretaceous:		
Marshalltown Formation:		
Yellow clay and sand .....	20	20
Black clay and sand .....	50	70
Englishtown Formation:		
White sand .....	40	110
Woodbury Clay and Merchantville Formation (undifferentiated):		
Black clay .....	140	250
Magothy and Raritan Formations:		
Fine sand .....	6	256
White sand .....	20	276

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 108, Dominic Tuscano  
(Log by Gus Hauser)

	Altitude 28 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Cretaceous:</b>		
Englishtown Formation:		
Brown, loamy sand . . . . .	3	3
Green and brown sand . . . . .	4	7
Green and brown loamy sand . . . . .	10	17
Tan loamy sand . . . . .	7	24
Englishtown Formation, Woodbury Clay, and Merchantville Formation (undifferentiated):		
Gray clay . . . . .	24	48
Black clay . . . . .	24	72
Brown and black clay . . . . .	24	96
Black clay . . . . .	64	160
Gray and black clay . . . . .	30	190
Magothy and Raritan Formations:		
Fine, gray sand . . . . .	2	192
Fine, gray sand with some fine gravel . . . . .	4	196

Well no. 111, Eva Diamond  
(Log by John Wrobel)

	Altitude 73 feet	
Cretaceous:		
Navesink Formation, Mount Laurel Sand, and Wenonah Formation and Marshalltown Formation (undifferentiated):		
No sample .....	100	100
Marl .....	20	120
Clay .....	12	132
Marl .....	11	143
White sand with some clay .....	7	150
Marl .....	6	156
Hard sandstone .....	4	160
Clay .....	19	179
Englishtown Formation:		
Clay and black sand .....	7	186
Water-bearing, black sand with some shells .....	16	202

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 113, Samuel Dickter  
(Log by Charles L. Mollitor Inc.)

	Altitude 50 feet Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>		
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Sand .....	7	7
Black marl .....	38	45
<b>Marshalltown Formation:</b>		
Green sandy marl .....	80	125
<b>Englishtown Formation:</b>		
Sand .....	10	135
<b>Englishtown Formation, Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Black marl .....	20	155
Green sand .....	5	160
Black marl .....	50	210
Black sand .....	2	212
Green marl .....	36	248
Hardpan .....	2	250
Green marl .....	35	285
<b>Magothy and Raritan Formations:</b>		
Sand .....	15	300

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 116, Charles J. Ruhle  
(Log by James L. Ruhle and J. H. Robbins <sup>4/</sup>)

	Altitude 57 feet	
	Thickness	Depth
	(feet)	(feet)
Quaternary:		
Pensauken Formation:		
Sand and gravel .....	3	3
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Yellow sandy clay .....	17	20
Marshalltown Formation:		
Glaucinitic sandy clay .....	30	50
Englishtown Formation:		
Glaucinitic sandy clay, but lighter in color .....	80	130
Woodbury Clay and Merchantville Formation (undifferentiated):		
Dark clay with fossils .....	45	175
Lighter-colored clay with fossils .....	55	230
Magothy and Raritan Formations:		
Water-bearing sand .....	10	240

<sup>4/</sup> Richards, 1954

Well no. 117, Felix Oliveto  
(Log by Paul Steffens)

	Altitude 63 feet	
	Thickness	Depth
	(feet)	(feet)
Topsoil .....	2	2
Quaternary:		
Pensauken Formation:		
Loam .....	10	12
Gravel .....	2	14
Cretaceous:		
Marshalltown Formation:		
Sandy marl .....	38	52
Englishtown Formation:		
Hardpan and sandy marl .....	35	87
Woodbury Clay and Merchantville Formation (undifferentiated):		
Marl .....	130	217
Magothy and Raritan Formations:		
Sand .....	6	223

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 120, William Breur  
(Log by Charles L. Mollitor Inc.)

	Altitude 78 feet Thickness (feet)	Depth (feet)
Topsoil .....	1	1
Cretaceous:		
Englishtown Formation:		
Yellow sand .....	8	9
Gravel .....	4	13
Wet sand .....	9	22
Green clay .....	7	29
Dry sand .....	3	32
Yellow sand .....	46	78
Woodbury Clay:		
Black marl .....	26	104
Gray clay .....	27	131
Sandy marl .....	7	138
Gray clay .....	13	151
Merchantville Formation:		
Black marl .....	6	157
Green marl .....	29	186
Black marl .....	28	214
Magothy and Raritan Formations:		
White, sandy clay .....	29	243
Dirty sand .....	9	252
White sand .....	7	259
Coarse sand .....	5	264
White clay .....	1	265
Coarse, white sand .....	7	272

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

(Well no. 122, Joseph E. Rudderow)  
(Log by Charles L. Mollitor Inc.)

	Altitude 48 feet	
	Thickness	Depth
	(feet)	(feet)
Topsoil .....	2	2
Quaternary and Cretaceous:		
Cape May Formation, Mount Laurel Sand, and Wenonah Formation (undifferentiated):		
Hard sand .....	9	11
Sandy marl with streaks of sand and shells .....	33	44
Cretaceous:		
Marshalltown Formation:		
Green marl .....	47	91
Englishtown Formation:		
Sand .....	2	93
Hardpan .....	2	95
Sand .....	5	100
Gray clay .....	13	113
Water-bearing sand .....	4	117
Englishtown Formation, Woodbury Clay, and Merchantville Formation (undifferentiated):		
No sample .....	121	238
Sand .....	3	241
Marl .....	27	268
Magothy and Raritan Formations:		
Coarse sand .....	3	271
Red clay .....	10	281
Sand .....	3	284
Red and gray clay .....	43	327
Sand and gravel .....	3	330
Gray clay .....	4	334
Hardpan .....	25	359
Gray clay .....	6	365
Hardpan .....	11	376
Gray clay .....	5	381
Hardpan .....	2	383
Gray clay .....	5	388
Hardpan and sand with wood .....	16	404
Hardpan .....	2	406
Streaks of sand and hardpan .....	7	413
Red and gray clay .....	26	439
Water-bearing, coarse sand with wood .....	17	456
Gray clay .....	1	457

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 126, Pittsburgh Plate Glass Co.  
(Log by Layne-New York Co.)

	Altitude 65 feet Thickness (feet)	Depth (feet)
Quaternary:		
Pensauken Formation:		
Red sand .....	12	12
Cretaceous:		
Woodbury Clay and Merchantville Formation (undifferentiated):		
Blue clay .....	126	138
Magothy and Raritan Formations:		
Medium to coarse, gray sand .....	9	147
Tough, blue and gray clay .....	103	250
Coarse, gray sand and gravel .....	33	283
Clay .....	6	289
Coarse, gray sand and gravel .....	26	315
Clay, coarse, gray sand, and gravel .....	3	318



Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 128, Dunlap Carpet Works <sup>5/</sup>  
(Log by C. G. Orcutt)

	Altitude 12 feet Thickness Depth (feet) (feet)	
<b>Cretaceous:</b>		
<b>Marshalltown Formation:</b>		
Dark (marl) .....	43	43
<b>Englishtown Formation:</b>		
Sand and gravel .....	3	46
Dark (marl) .....	61	107
Fine sand .....	4	111
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Dark (marl) or clay .....	134	245
<b>Magothy and Raritan Formations:</b>		
Gray sand .....	5	250
Dark sandy clay .....	12	262
Red clay .....	11	273
White clay .....	1	274
Brown sandy clay .....	20	294
White clay .....	2	296
Light sandy clay .....	69	365
Red clay .....	2	367
Light sandy clay .....	43	410
Red clay .....	2	412
Brown clay .....	18	430
Red clay .....	6	436
Light sandy clay .....	6	442
Fine, gray sand .....	15	457
Sand and clay .....	52	509
Light sandy clay .....	37	546
Red clay .....	14	560
Sandy clay and fine sand .....	48	608
Yellow clay .....	3	611
Brown clay .....	9	620
Red clay .....	40	660
Fine sand with some brown clay .....	7	667
Fine and coarse sand and some gravel .....	8	675

<sup>5/</sup> Smock, 1893, p. 303-304

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 129, Acme Market (Log by Charles L. Mollitor Inc.)			Altitude 62 feet	
	Thickness (feet)	Depth (feet)		
Topsoil .....	1	1		
<b>Cretaceous:</b>				
<b>Mount Laurel Sand and Wenonah Formation:</b>				
Yellow sand .....	25	26		
Yellow clay .....	8	34		
<b>Marshalltown Formation:</b>				
Gray marl .....	11	45		
Soft marl .....	4	49		
Brown sand .....	3	52		
Black marl .....	21	73		
<b>Englishtown Formation:</b>				
Muddy sand and clay .....	12	85		
Fine gray sand .....	8	93		
Gray clay .....	12	105		
Fine sand .....	7	112		
Gray clay .....	9	121		
Fine sand .....	7	128		
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>				
Gray clay .....	80	208		
Green marl .....	35	243		
Sticky, hard, black marl .....	35	278		
<b>Magothy and Raritan Formations:</b>				
White clay .....	2	280		
Gray marl .....	3	283		
Hard, red clay .....	19	302		
White clay .....	32	334		
Gray clay .....	40	374		
Hard, white clay .....	18	392		
Sandy clay .....	21	413		
Sticky, red clay .....	8	421		
Gray clay .....	14	435		
Sandy clay .....	16	451		
Sand .....	6	457		
Sand .....	13	470		
Gray clay .....	2	472		
Sand .....	2	474		
Gray clay .....	16	490		
Sand .....	1	491		
Clay .....	5	496		
Sand .....	1	497		
Clay .....	1	498		
Sand .....	7	505		
White clay .....	10	515		
Sand .....	7	522		
Clay .....	2	524		
Sand .....	21	545		
Clay .....	1	546		

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 130, Jacobstown Baptist Church  
(Log by Andrew White)

	Altitude 175 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Quaternary, Tertiary, and Cretaceous:</b>		
Pensauken, Kirkwood, and Vincentown Formations (undifferentiated):		
Yellow sandy clay .....	48	48
Hornerstown and Red Bank Sands, and Navesink Formation		
(undifferentiated):		
Light-brown marl .....	17	65
Dark-green marl .....	22	87
Black clay and sand .....	13	100
Gray clay and black sand with some shells .....	20	120
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Black sand with some fine gravel and shells .....	8	128
Gray to black sand with some fine gravel and shells .....	12	140

Well no. 131, Jersey Central Power and Light Co.  
(Log by Greenhalgh and Kaye)

	Altitude 88 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Tertiary:</b>		
Vincentown Formation:		
Yellow sand .....	8	8
Gray clay and sand .....	32	40
<b>Tertiary and Cretaceous:</b>		
Hornerstown and Red Bank (?) Sands, and Navesink Formation		
(undifferentiated):		
Dark-green and black marl .....	60	100
Light-green marl .....	10	110
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Sand with shells .....	11	121

Table 3 – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 132, Edward H. Gancarz  
(Log by Greenhalgh and Kaye)

	Altitude 121 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Cretaceous:</b>		
Mount Laurel Sand, Wenonah and Marshalltown Formations (undifferentiated):		
Black and green marl .....	30	30
Yellow sandy clay .....	10	40
Black clay .....	80	120
<b>Englishtown Formation:</b>		
Fine, gray sand .....	25	145
Coarse, gray sand with limestone and ironstone .....	12	157

Well no. 134, Spartan Village, Inc.  
(Log by Louis Bainbridge and Charles L. Mollitor Inc.)

	Altitude 154 feet	
Quaternary and Tertiary:		
Bridgeton and Kirkwood Formations.(undifferentiated):		
Dry sand .....	25	25
Tertiary:		
Vincentown Formation:		
Brown clay .....	3	28
Black marl .....	15	43
Black sandy marl .....	17	60
Tertiary and Cretaceous:		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Green marl with shells .....	70	130
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Black and white sand .....	20	150
Wenonah and Marshalltown Formations (undifferentiated):		
Green clayey marl .....	130	280
Englishtown Formation:		
Brown clay with some pyrite, wood, and brown sand .....	7	287
Hard, brown clay .....	18	305
Englishtown Formation and Woodbury Clay (undifferentiated):		
Hard, blue to brown clay .....	60	365
Hard, gray sand and sandy clay .....	30	395

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 136, B. F. Goodrich Co.  
(Log by Paul Steffens)

	Altitude 34 feet Thickness (feet)	Depth (feet)
<b>Quaternary:</b>		
Cape May Formation:		
Sand .....	10	10
Sand and gravel .....	4	14
<b>Cretaceous:</b>		
Magothy and Raritan Formations:		
Gray clay .....	17	31
Black hardpan .....	2	33
Muddy sand .....	8	41
White clay .....	11	52
Muddy sand and gravel .....	6	58
White clay .....	17	75
Sand and gravel .....	3	78
Brown clay .....	1	79
Fine gravel .....	2	81
Dirty sand .....	4	85
Dirty sand and gravel .....	21	106
Brown sandy clay .....	1	107
Sand and gravel .....	17	124
Sandy clay .....	5	129
White clay .....	5	134
Sand and gravel .....	3	137
White clay .....	4	141
Sand .....	6	147
Sandy clay .....	8	155
Reddish-brown clay .....	2	157
<b>Early Paleozoic (?):</b>		
Wissahickon Formation:		
Soft rock .....	5	162

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 138, Riverside-Alloy Metal Co.  
(Log by Ridpath and Potter)

	Altitude 18 feet	
	Thickness	Depth
	(feet)	(feet)
Quaternary and Cretaceous:		
Cape May, Magothy and Raritan Formations (undifferentiated):		
Fine to coarse sand, clay, and some gravel .....	120	120
Early Paleozoic (?):		
Wissahickon Formation:		
Mica gneiss .....	680	800

Well no. 140, Jane Tabone  
(Log by Charles L. Mollitor Inc.)

	Altitude 130 feet	
Quaternary and Tertiary:		
Bridgeton and Kirkwood Formations (undifferentiated):		
Yellow sand and gravel .....	22	22
Red sand and gravel .....	26	48
Tertiary:		
Kirkwood Formation:		
Yellow sand .....	22	70
Black mud .....	5	75
Muddy sand .....	30	105
Manasquan Formation:		
Green marl .....	25	130
Vincentown Formation:		
Green and black sand .....	10	140

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 141, A. F. Brunt  
(Log by Charles L. Mollitor Inc.)

	Altitude 83 feet	
	Thickness (feet)	Depth (feet)
Topsoil .....	1	1
<b>Tertiary:</b>		
Kirkwood Formation:		
Yellow sand .....	4	5
Gray sand .....	24	29
<b>Tertiary and Cretaceous:</b>		
Manasquan and Vincentown Formations, Hornerstown Sand and Navesink Formation (undifferentiated):		
Sandy marl .....	14	43
Soft, black marl .....	9	52
Sandy marl .....	6	58
Green marl .....	22	80
Soft marl .....	2	82
Green sandy marl .....	5	87
Hard, green clay .....	9	96
Green marl .....	42	138
Black sand with some shells .....	4	142
Green marl .....	17	159
Black marl .....	42	201
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Sand with shells .....	9	210

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 143, Mr. Reilley  
(Log by Henry B. Kummel <sup>6/</sup>)

	Altitude 70-80 feet	
	Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
Kirkwood (?) Formation:		
Sand and gravel .....	10	10
Kirkwood Formation:		
Brown sandy clay .....	50	60
<b>Tertiary and Cretaceous:</b>		
Manasquan and Vincentown Formations, Hornerstown Sand and Navesink Formation (undifferentiated):		
Greenish clay with some marl and a trace of lignite and pebbles .....	80	140
Greensand (glauconite) marl .....	80	220
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Fine, gray sand with some marl at the top and bottom .....	110	330
Coarse, yellow sand with shells, clay, and very micaceous dark sand .....	20	350
Marshalltown Formation:		
Black sandy clay .....	20	370
Very fine, greenish-black micaceous sand with glauconite .....	10	380
No sample .....	10	390
Englishtown Formation:		
Fine, gray quartz sand .....	10	400
Fine quartz sand with some glauconite .....	20	420
Water-bearing, coarse quartz sand .....	10	430

<sup>6/</sup> Kummel, 1910, p. 85-90



Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 144, Joseph O'Neill  
(Log by A. C. Schultes and Sons)

	Altitude 91 feet Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
<b>Kirkwood Formation:</b>		
Sand and gravel .....	60	60
Yellow sandy clay .....	20	80
<b>Manasquan Formation:</b>		
Green marl .....	38	118
<b>Tertiary and Cretaceous:</b>		
<b>Vincentown Formation, Hornerstown Sand, and Navesink   Formation (undifferentiated):</b>		
Hard sand .....	2	120
Green clay .....	25	145
White sand .....	4	149
Clay and green marl .....	26	175
Green marl .....	6	181
Clay and green marl .....	52	233
Green marl and sand with shells .....	27	260
Green marl .....	15	275
Green marl with some sand .....	16	291
<b>Cretaceous:</b>		
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Sand and gravel .....	9	300
Clayey sand .....	19	319
Medium and coarse sand .....	18	337

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 145, Country Lakes Inc.  
(Log by Charles L. Mollitor Inc.)

	Altitude 82 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Tertiary:</b>		
<b>Kirkwood Formation:</b>		
White sand .....	2	2
Brown sand .....	26	28
Sandy marl .....	23	51
Fine, dirty sand .....	11	62
Sandy marl .....	7	69
Clay .....	16	85
Sandy clay .....	21	106
<b>Tertiary and Cretaceous:</b>		
<b>Manasquan and Vincentown Formations, Hornerstown Sand, and Navesink Formation (undifferentiated):</b>		
Green marl .....	63	169
Sand .....	11	180
Sticky, green marl .....	80	260
Green marl with shells .....	61	321
<b>Cretaceous:</b>		
<b>Mount Laurel (?) Sand and Wenonah (?) Formation:</b>		
Gray sandy clay .....	8	329
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Sand with shells .....	16	345

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 148, Helis Stock Farm (Log by William Blaisdell and Orcutt Bros. <u>7/</u> )		Altitude approx. 70 feet	
		Thickness (feet)	Depth (feet)
<b>Cretaceous:</b>			
<b>Mount Laurel Sand and Wenonah Formation:</b>			
Water-bearing, yellowish loamy sand .....	14	14	
Fine sand with dark mud .....	34	48	
<b>Marshalltown Formation:</b>			
Stiff, black sandy clay .....	24	72	
Water-bearing, fine, muddy sand .....	1	73	
Stiff, black sandy clay .....	9	82	
<b>Englishtown Formation:</b>			
Water-bearing, fine sand with scattered layers of sandstone, clay, and shells .....	34	116	
Black sandy clay .....	1	117	
Water-bearing, fine sand .....	7	124	
Black sandy clay .....	1	125	
Water-bearing, fine sand .....	3	128	
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>			
Dark sandy clay with scattered layers of sandstone and shells .....	50	178	
Dark sandy clay and scattered layers of sandstone, shells, and wood .....	128	306	
<b>Magothy and Raritan Formations:</b>			
Fine sand with some gravel and brown clay .....	8	314	
Red and white clay .....	24	338	
Sand and sandstone with considerable wood and some clay .....	18	356	
No sample .....	11	367	
Fine sand .....	3	370	
Coarse sand which yields some water .....	17	387	
Coarse gravel .....	3	390	
Fine sand .....	5	395	
White clay .....	5	400	
White sand and some coarse gravel .....	10	410	
Fine, white sand .....	22	432	
Dark micaceous sand .....	8	440	
Hard clay .....	11	451	
Coarse, dark sand and red clay .....	8	459	
Fine, white sand .....	20	479	
Dark clay .....	4	483	
White and red clay .....	12	495	
Sand .....	2	497	
White, red and dark clay .....	78	575	
Sand .....	13	588	
Coarse, white sand with clay .....	12	600	
Coarse sand and gravel .....	3	603	
Fine sand .....	16	619	
Clay .....	25	644	
Sand .....	7	651	
White clay .....	13	664	
Sand .....	15	679	
White clay .....	2	681	
Gravel and coarse sand .....	9	690	
Red and white clay .....	25	715	
Sand .....	At	715	

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 149, Kauffman and Minter Inc.  
(Log by Charles L. Mollitor Inc.)

	Altitude 74 feet Thickness (feet)	Depth (feet)
Fill .....	1	1
Topsoil .....	1	2
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Dirty sand .....	17	19
Mount Laurel (?) Sand and Wenonah (?) Formation:		
Sandy marl .....	39	58
Marshalltown Formation:		
Marl .....	36	94
Englishtown Formation:		
Sand and gravel .....	13	107

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 151, Helis Stock Farm (Log by Ridpath and Potter)		
	Altitude 96 feet	
	Thickness	Depth
	(feet)	(feet)
Tertiary and Cretaceous:		
Undifferentiated:		
Green sandy marl with some cemented sand .....	342	342
Soft, dark clay .....	69	411
Cretaceous:		
Magothy and Raritan Formations:		
Hardpan with layers of light-colored sand .....	29	440
Hardpan with wood .....	18	458
Hardpan with some thin streaks of sand .....	40	498
Water-bearing, light-colored sand .....	10	508
Hardpan .....	14	522
Hard, dark clay .....	39	561
Very fine, light-colored sand .....	12	579
Tough, dark clay .....	4	583
Fine, brown sand .....	4	587
Clay and hardpan .....	26	613
Dark sand .....	2	615
Very hard, white clay .....	5	620
Hard, red clay .....	87	707
Red and white clay .....	7	714
Fine, white sand .....	3	717
Hard, white clay .....	6	723
Hardpan .....	21	744
Clay .....	27	771
Hardpan .....	11	782
Clay and hardpan .....	27	809
Light-colored sand .....	5	814
White clay .....	3	817
Red clay .....	78	895
Sand .....	1	896
Hardpan .....	7	903
Clay .....	25	928
Fine sand .....	6	934
Hardpan .....	16	950
Sandy clay .....	10	960
Hardpan .....	3	963
No sample .....	24	987
Early Paleozoic (?):		
Wissahickon (?) Formation:		
Hard sandstone, probably weathered bedrock .....	1	988

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 152, Ridge Stables  
(Log by Paul Steffens)

	Altitude 41 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Loam .....	22	22
Cretaceous:		
Englishtown Formation:		
Sand .....	29	51
Woodbury Clay and Merchantville Formation (undifferentiated):		
Marl .....	165	216
Magothy and Raritan Formations:		
Sand with wood and streaks of clay .....	14	230

Well no. 155, Tallman Brothers, Inc.  
(Log by W. R. Clair)

	Altitude 42 feet	
Quaternary:		
Cape May Formation:		
Fine sand .....	18	18
Yellow clay .....	6	24
Cretaceous:		
Englishtown Formation:		
Fine sand .....	28	52

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 157, Ross Monroe  
(Log by Harvey Beebe Sr.)

	Altitude 46 feet Thickness (feet)	Depth (feet)
Yellow topsoil .....	6	6
Quaternary:		
Cape May Formation:		
Gravel .....	6	12
Tertiary:		
Kirkwood Formation:		
Black clay .....	9	21
Tertiary and Cretaceous:		
Manasquan and Vincentown Formations, Hornerstown Sand, and Navesink Formation (undifferentiated):		
Marl .....	136	157
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Black and white sand .....	80	237
Marshalltown Formation:		
Hardpan .....	11	248

Well no. 158, Hampton Lakes Development Co.  
(Log by Charles L. Mollitor Inc.)

	Altitude 49 feet	
Loamy topsoil .....	2	2
Quaternary:		
Cape May Formation:		
Sand .....	8	10
Tertiary:		
Kirkwood Formation:		
Brown sandy marl .....	36	46
Black marl .....	13	59
Manasquan and Vincentown Formations and Hornerstown Sand (undifferentiated):		
Light-green marl .....	83	142
Dark-green marl .....	~	163
Cretaceous:		
Navesink Formation:		
Black marl with shells .....	6	169
Black marl .....	29	198
Navesink (?) Formation:		
Sandy marl with sand streaks .....	47	245
Mount Laurel Sand and Wenonah Formation:		
Sand with shells and clay streaks .....	23	268

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 159, Vincentown Diner  
(Log by Edward Robbins Sr.)

	Altitude 72 feet Thickness (feet)	Depth (feet)
<b>Tertiary and Cretaceous:</b>		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Black clay .....	80	80
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Sand with iron oxide stains .....	40	120
Marshalltown Formation:		
Black clay .....	110	230
Englishtown Formation:		
Sand .....	20	250

Well no. 160, Frank Mayo  
(Log by Greenhalgh and Kaye)

	Altitude 140 feet	
Tertiary:		
Cohansey Sand:		
Yellow sand and gravel .....	20	20
Clay and gravel .....	10	30
Black clay .....	15	45
Gravel .....	10	55
Clay .....	5	60
Yellow and red sand and clay .....	9	69



Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 162, A. Marone  
(Log by Charles L. Mollitor Inc.)

	Altitude 40 feet	
	Thickness (feet)	Depth (feet)
Topsoil .....	2	2
Cretaceous:		
Englishtown Formation:		
Yellow sand .....	14	16
Englishtown (?) Formation:		
Soft, gray marl .....	41	57
Woodbury Clay and Merchantville Formation (undifferentiated):		
Gray clay .....	53	110
Green marl .....	84	194
Magothy (?) and Raritan (?) Formation:		
Gray marl .....	41	235
Magothy and Raritan Formation:		
Sand .....	30	265

Well no. 165, Alexander Construction Co.  
(Log by Paul Steffens)

	Altitude 57 feet	
Loamy topsoil .....	4	4
Cretaceous:		
Englishtown Formation:		
Dry sand .....	3	7
Water-bearing sand and gravel .....	9	16
Sandy marl .....	32	48
Brown sand .....	7	55
Woodbury Clay and Merchantville Formation (undifferentiated):		
Marl .....	136	191
Magothy and Raritan Formations:		
Dark-gray clay .....	11	202
Coarse sand and gravel .....	13	215

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 166, John S. Pew  
(Log by Charles L. Mollitor Inc.)

	Altitude 79 feet Thickness (feet)	Depth (feet)
Topsoil .....	1	1
Cretaceous:		
Englishtown Formation:		
Yellow sand .....	42	43
Woodbury (?) Clay:		
Gray marl .....	32	75
Woodbury Clay and Merchantville Formation (undifferentiated):		
Hard, black marl .....	49	124
Gray marl .....	12	136
Green marl .....	13	149
Soft, gray marl .....	5	154
Green marl .....	9	163
Gray marl .....	48	211
Magothy and Raritan Formations:		
Sand .....	17	228

Well no. 169, John W. Hampton  
(Log by Charles L. Mollitor Inc.)

	Altitude 55 feet	
Topsoil .....	1	1
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Sand .....	34	35
Marshalltown Formation:		
Marl .....	93	128
Hardpan .....	2	130
Englishtown Formation:		
Muddy sand .....	18	148
Sand .....	12	160

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 171, Ionac Chemical Co.  
(Log by A. C. Schultes and Sons)

	Altitude 30 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Fine to coarse, brown sand and gravel .....	10	10
Tertiary and Cretaceous:		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Green marl .....	40	50
Dark-gray clay .....	15	65
Cretaceous:		
Mount Laurel Sand and Wenonah Formation:		
Sand with shells .....	7	72
Dark-gray clay .....	5	77
Fine to coarse, green sand .....	15	92
Dark-gray clay .....	11	103
Fine to medium, gray sand .....	33	136
Dark-gray silty clay .....	12	148
Fine, greenish-gray sand .....	7	155
Marshalltown Formation:		
Gray silty clay .....	37	192
Dark-gray clay .....	11	203
Englishtown Formation:		
Fine, gray sand .....	7	210
Hardpan .....	1	211
Fine, gray sand .....	8	219
Gray silty clay .....	13	232
Fine, gray sand .....	6	238
Gray, silty clay .....	19	257
Woodbury Clay:		
Dark gray clay .....	65	322
Gray silty clay .....	34	356
Merchantville Formation:		
Hardpan .....	2	358
Dark gray clay .....	16	374
Hardpan .....	15	389
Dark gray clay .....	25	414
Magothy and Raritan Formations:		
Fine, gray clay .....	15	429
Gray silty clay .....	9	438
Hard, red clay .....	24	462
Sand .....	4	466
Red clay .....	17	483
Sand .....	7	490
Fine to medium, variegated sand .....	31	521
Red clay .....	1	522
Sand .....	1	523

(Continued)

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

**Well no. 171, Ionac Chemical Co.  
(Log by A. C. Schultes and Sons)  
(Continued)**

<b>Cretaceous — Continued:</b>	<b>Thickness</b>	<b>Depth</b>
<b>Magothy and Raritan Formations — Continued:</b>	<b>(feet)</b>	<b>(feet)</b>
Hardpan .....	1	524
Red clay .....	5	529
Hardpan .....	11	540
Sand .....	9	549
Red and white clay .....	14	563
Fine, gray clay .....	6	569
Sandy clay .....	5	574
Red clay .....	2	576
Gray sandy clay with shells .....	2	578
Sand .....	10	588
Fine, gray sand .....	6	594
Sand .....	6	600
Fine, gray sand .....	18	618
Gray clay with shells .....	2	620
Sand .....	4	624
Fine, variegated sand .....	7	631
Hardpan .....	1	632
Gray clay with shells .....	21	653
Fine sand .....	3	656
Gray clay with shells .....	15	671
Hardpan .....	2	673
Dark gray silty clay .....	29	702
Hardpan .....	1	703
Sandy clay .....	13	716
Fine, gray sand .....	5	721
Sandy clay .....	4	725
Gray sand .....	2	727
Red and gray clay .....	21	748
Fine sand .....	26	774
Red clay .....	8	782
Hardpan .....	1	783
Gray and red sandy clay .....	7	790
Sand .....	4	794
Sandy clay .....	1	795
Sand .....	2	797
Hard, red clay .....	1	798
Sand .....	10	808
Fine to medium sand .....	19	827
Red clay .....	13	840
Fine sand .....	11	851
Red clay .....	38	889
Sand .....	1	890
Hardpan .....	1	891
Red clay .....	19	910
Fine sand .....	11	921

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 172, Burlington Co. Institution No. 1  
(Log by Layne-New York Co., Inc.)

	Altitude 59 feet Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
<b>Kirkwood Formation:</b>		
Topsoil (?) .....	23	23
Sand .....	21	44
<b>Tertiary and Cretaceous:</b>		
<b>Manasquan and Vincentown Formations, Hornerstown Sand and Navesink Formation (undifferentiated):</b>		
Sand and clay .....	22	66
Blue clay .....	144	210
<b>Cretaceous:</b>		
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Black sand .....	42	252
<b>Marshalltown Formation:</b>		
Black sand and clay .....	107	359
<b>Englishtown Formation:</b>		
Sand and gravel .....	41	400
Clay .....	21	421
Clay .....	42	463
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Blue marl .....	151	614
<b>Magothy and Raritan Formation:</b>		
Gravel and clay .....	20	634
Tough clay .....	41	675
Clay .....	27	702
Gray sand .....	10	712
Clay .....	58	770
<b>Early Paleozoic (?):</b>		
<b>Wissahickon Formation:</b>		
Rock .....	6	776

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 178, William C. Rahilly  
(Log by Greenhalgh and Kaye)

	Altitude 195 feet	
	Thickness	Depth
	(feet)	(feet)
Quaternary and Tertiary:		
Bridgeton Formation, Cohansey Sand, and Kirkwood Formation (undifferentiated):		
Hard, fine, yellow sand and clay .....	49	49
Tertiary:		
Vincentown Formation and Hornerstown Sand (undifferentiated):		
Yellow clay .....	11	60
Green marl .....	40	100
Cretaceous:		
Navesink Formation:		
Black marl .....	18	118
Black clay with lignite and shells .....	22	140
Mount Laurel Sand and Wenonah Formation:		
Green sand with shells and clay .....	18	158

Well no. 181, Wrightstown Municipal Utility Authority  
(Log by (unknown) 8/)

	Altitude 120 feet	
Tertiary:		
Kirkwood and Vincentown Formations (undifferentiated):		
Sandy clay .....	60	60
Tertiary and Cretaceous:		
Undifferentiated:		
Sand and clay with shells .....	130	190
Clay .....	56	246
Cretaceous:		
Englishtown Formation:		
Water-bearing, medium glauconitic sand .....	8	254
Clayey sand .....	14	268

8/ Richards, 1948, p. 39-76

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 183, Charles B. Hutchinson  
(Log by Charles L. Mollitor Inc.)

	Altitude 180 feet Thickness    Depth (feet)        (feet)	
<b>Tertiary:</b>		
<b>Cohansey Sand and Kirkwood Formation (undifferentiated):</b>		
Loam .....	13	13
Sand with large gravel .....	14	27
Sandy clay .....	8	35
<b>Vincentown Formation and Hornerstown Sand:</b>		
Green marl .....	55	90
<b>Cretaceous:</b>		
<b>Navesink Formation:</b>		
Black marl .....	30	120
Green marl .....	2	122
<b>Mount Laurel Sand and Wenonah Formation:</b>		
Green sand with shells .....	12	134

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 185, Joseph J. White Inc.  
(Log by Ridpath & Potter)

	Altitude 95 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Tertiary and Cretaceous:</b>		
Cohansey Sand and Kirkwood Formation (undifferentiated):		
Soil (?) .....	15	15
Fine, yellow sand with some coarse, angular grains .....	10	25
Coarse, angular quartz sand .....	5	30
Fine, gray micaceous sand .....	At	75
Very fine, gray sand with much mica .....	At	100
Fine, dark-gray sand with some dark-brown clay .....	At	115
Manasquan and Vincentown Formations, Hornerstown Sand, and Navesink Formation (undifferentiated):		
Glauconite marl with some quartz sand and mica .....	At	132
Glauconite marl .....	At	150
Glauconite marl with some white clay and white quartz sand .....	At	207
Light-green sandy clay with a little glauconite .....	At	310
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Fine, gray sand with a little lignite .....	At	341
Coarse, black and white glauconitic sand with shell fragments .....	At	380
Marshalltown Formation:		
Yellowish gray glauconitic sand .....	At	400
Fine, gray glauconitic sand with some mica .....	At	440
Fine, gray glauconitic sand with some mica .....	At	450
Englishtown Formation:		
Medium-yellow sand with some glauconite and mica .....	At	535
Medium-yellow sand with some glauconite and mica .....	15	550
Woodbury Clay and Merchantville Formation (undifferentiated):		
Very fine, dark-gray sand with glauconite and mica .....	At	700
Very fine, dark-gray sand with glauconite and mica .....	25	725
Very fine, dark-gray sand and sandy micaceous clay .....	At	730
Fine, dark-gray, sand with much glauconite and mica .....	At	800



Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 186, Joseph J. White Inc.  
(Log by J. W. Jenkins)

	Altitude 95 feet Thickness (feet)	Depth (feet)
<b>Tertiary:</b>		
Cohansey Sand:		
Fine, yellow sand .....	52	52
Kirkwood Formation:		
Fine, gray sand with clay .....	86	138
Manasquan Formation:		
Hard, dark-green clay .....	14	152
Dark-green clay with black sand .....	18	170
Vincentown Formation:		
Light-green clay .....	58	228
<b>Tertiary and Cretaceous:</b>		
Hornerstown Sand and Navesink Formation (undifferentiated):		
Dark-green clay with black sand and shells .....	11	239
Dark-brown clay with shells and black sand .....	57	296
Green clay with some black sand .....	24	320
Dark-green and black clay, and black sand with shells .....	25	345
<b>Cretaceous:</b>		
Mount Laurel Sand and Wenonah Formation:		
Sand .....	20	365
Water-bearing, fine to medium sand .....	23	388

Well no. 196, Taunton Lakes Water Co.  
(Log by J. Henry Robbins)

	Altitude 57 feet	
<b>Tertiary:</b>		
Kirkwood Formation:		
Yellow clay .....	25	25
Black clay .....	55	80
Manasquan and Vincentown (?) Formations and Hornerstown Sand (undifferentiated):		
Green clay .....	120	200
<b>Cretaceous:</b>		
Navesink Formation:		
Black clay .....	20	220
Undifferentiated:		
Green clay and fine sand .....	10	230
Mount Laurel Sand and Wenonah Formation:		
Green and white sand .....	22	252

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 198, Riverton-Palmyra Water Company  
(Log by A. C. Schultes and Sons)

	Altitude 79 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Quaternary:</b>		
Cape May Formation:		
Fine, brown sand .....	4	4
Fine, brown clayey sand .....	2	6
Fine, brown sand .....	6	12
<b>Cretaceous:</b>		
Merchantville Formation:		
Clay .....	7	19
Magothy and Raritan Formations:		
Fine to medium, brown sand .....	6	25
Medium to coarse, brown sand .....	5	30
Sand, clay, and stones .....	7	37
Sand and clay .....	2	39
Gravel and clay .....	1	40
Medium to coarse sand .....	4	44
Gravel .....	2	46
Fine to coarse sand .....	4	50
Gravel and clay .....	4	54
Sandy clay .....	5	59
Fine sand .....	4	63
Coarse sand and gravel .....	2	65
Clay and sand .....	11	76
Medium sand .....	1	77
Clay and sand .....	8	85
Coarse gravel and sand .....	3	88
Sandy clay .....	10	98
Clay and gravel .....	4	102
Clay and sand .....	4	106
Brown sand .....	12	118
Coarse gravel .....	5	123
Clay .....	1	124
Brown sand and gravel .....	25	149
Sand and clay .....	6	155
Gravel and sand .....	5	160
Sandy clay .....	5	165
Sand and gravel .....	14	179
Gravel and sand .....	9	188
Gravel and clay .....	2	190
White clay .....	4	194
Sand and gravel .....	6	200

(Continued)

Table 3. - Selected drillers' logs of wells in Burlington County, N. J. - Continued

Well no. 198, Riverton-Palmyra Water Company  
(Log by A. C. Schultes and Sons)  
(Continued)

Cretaceous - Continued:	Thickness	Depth
Magothy and Raritan Formations - Continued:	(feet)	(feet)
Clay and sand .....	11	211
White clay .....	6	217
Sandy clay .....	1	218
Sand and clay .....	14	232
Sand .....	5	237
Sand and clay .....	12	249
Sand and gravel .....	9	258
Clay and gravel .....	1	259
Sand and gravel .....	6	265
Medium to coarse, brown sand with some gravel .....	6	271
Medium to coarse, brown sand and gravel .....	11	282
Red clay .....	5	287
Very hard, brown and white clay .....	2	289
Coarse gravel .....	13	302
Early Paleozoic (?):		
Wissahickon Formation:		
Stones, gravel, and weathered, micaceous rock .....	2	304
Rock with some sand .....	2	306
Weathered rock .....		308

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 199, Moorestown Township Water Department  
(Log by Layne-New York Co., Inc.)

	Altitude 59 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Cretaceous:</b>		
<b>Englishtown Formation:</b>		
Yellow sand .....	8	8
<b>Undifferentiated:</b>		
Gray clay with hard streaks .....	84	92
<b>Magothy and Raritan Formations:</b>		
Gray clay .....	48	140
Red, white, and gray clay .....	10	150
Gray clay with hard streaks .....	70	220
Coarse, gray sand .....	12	232
Tough clay .....	4	236
Coarse, gray sand .....	19	255
Coarse, gray clayey sand .....	20	275
Medium to coarse sand with pebbles .....	68	343
Red and white variegated clay .....	17	360

Well no. 201, Cedar Bridge Fire Tower  
(Log by J. Henry Robbins)

	Altitude 200 feet	
Tertiary:		
Cohansey Sand:		
Yellow clayey sand with small pebbles .....	30	30
Layers of light-gray to buff clay and sand .....	30	60
Fine to coarse, yellow clayey sand with small pebbles .....	10	70
Fine to coarse, yellow clayey sand .....	30	100
Light-gray clay with thin sand layers .....	10	110

Table 3. — Selected drillers' logs of wells in Burlington County, N. J. — Continued

Well no. 202, Inductotherm Corporation  
(Log by Charles L. Mollitor Inc.)

	Altitude 65 feet	
	Thickness	Depth
	(feet)	(feet)
<b>Cretaceous:</b>		
<b>Englishtown Formation:</b>		
Clay .....	4	4
Sand .....	39	43
<b>Woodbury Clay and Merchantville Formation (undifferentiated):</b>		
Gray marl .....	37	80
Black clay .....	58	138
Green marl .....	70	208
<b>Magothy and Raritan Formations:</b>		
Gray sandy clay .....	4	212
Water-bearing sand .....	6	218
Red clay .....	16	234
Gray clay and gravel .....	13	247
White clay .....	1	248
Black clay and gravel .....	36	284
White clay .....	6	290
Black clay .....	15	305
White clay .....	4	309
Red clay .....	16	325
White clay with sand .....	20	345
White clay .....	11	356
Gray clay .....	7	363
White clay with sand .....	22	385
Water-bearing sand .....	1	386
Gray clay .....	3	389
Sand .....	4	393
Clay .....	12	405
Sand .....	39	444
Gray clay .....	1	445

Table 3. – Selected drillers' logs of wells in Burlington County, N. J. – Continued

Well no. 208, Mount Laurel Water Company  
(Log by A. C. Schultes and Sons)

	Altitude 30 feet Thickness (feet)	Depth (feet)
Quaternary:		
Cape May Formation:		
Topsoil, sand, and gravel .....	6	6
Cretaceous:		
Marshalltown Formation:		
Gray sandy clay .....	21	27
Englishtown Formation:		
Fine, gray sand .....	19	46
Woodbury Clay:		
Gray clay and fine, hard sand.....	52	98
Merchantville Formation:		
Black marl .....	94	192
Magothy and Raritan Formations:		
Fine to medium sand .....	24	216
Fine to coarse, gray sand .....	10	226
Black clay .....	28	254
Fine to coarse, gray sand .....	5	259
White clay .....	4	263
Fine to coarse gray sand and gravel .....	34	297
Clay .....	8	305
Sandy clay .....	18	323
Clay .....	6	329
Sand and gravel .....	30	459
Red clay .....	5	464
Sandy clay .....	16	480
Coarse sand and gravel .....	13	493
Red clay .....	2	495
Sand and gravel .....	13	508
Red clay .....	1	509
Sand, gravel, and clay .....	8	517
Coarse sand and gravel .....	38	555
Red clay .....	3	558
Coarse sand and gravel .....	21	579
Red clay .....	3	582
Coarse sand and gravel .....	12	594
Early Paleozoic (?):		
Wissahickon Formation:		
Weathered rock .....	1	595

Map No.	Date of collection	Temperature (°F)	Silica (SiO <sub>2</sub> )	Total iron (Fe)	Manganese (Mn)	Calcium (Ca)	Magnesium (Mg)	Sodium (Na)	Potassium (K)	Bicarbonate (HCO <sub>3</sub> )	Sulfate (SO <sub>4</sub> )	Chloride (Cl)	Fluoride (F)	Nitrate (NO <sub>3</sub> )	Dissolved solids (residue on evaporation at 180°C)	Hardness as CaCO <sub>3</sub>		Specific conductance (micro-mhos at 25°C)	pH	Source of data	Map no.
																Calcium magnesium	Noncarbonate				
3	5-24-51	55	10	0.01	3.6	23	10	14	10	80	38	20	0.2	5.9	203	98	33	318	7.2	USGS	3
5	9-9-52	----	----	.10	.00	----	----	----	----	12	----	5.0	.0	----	54	28	18	---	5.7	D	5
6	5-22-51	57	8.4	.02	.00	.28	4.5	2.6	5.5	103	12	1.2	.1	.5	113	88	4	190	7.6	USGS	6
7	5-31-61	----	----	15	.18	----	----	----	----	52	----	----	----	----	---	38	0	---	6.9	NJDH	7
8	9-14-54	56	----	.06	----	----	----	14	----	35	42	9.8	----	16	---	68	39	206	6.3	USGS	8
11	3-22-51	56	27	.17	.10	43	4.0	2.2	5.7	149	7.0	2.6	.4	.3	166	124	2	251	8.2	USGS	11
13	5-3-51	55	12	.00	.00	8.0	4.4	4.8	1.4	8	14	8.0	.0	21	84	38	32	112	7.1	USGS	13
14	6-19-59	----	9.8	2.4	.05	16	2.8	9.0	----	68	12	1.8	.1	.2	105	52	0	137	6.5	USGS	14
15	6-19-59	----	11	2.2	.04	16	3.3	8.3	----	68	11	2.8	.1	.2	85	54	0	140	6.8	USGS	15
16	6-18-59	----	9.8	2.1	.09	14	4.7	6.9	----	66	12	1.4	.1	.4	100	55	1	134	6.4	USGS	16
18	3-1-61	----	7.8	2.0	.06	15	2.1	8.2	3.8	64	8.2	5.1	.1	.4	84	46	0	139	7.1	USGS	18
19	3-1-61	----	8.3	2.8	.09	16	2.6	6.0	3.8	67	8.2	2.5	.0	.8	82	51	0	133	6.8	USGS	19
20	3-1-61	----	9.8	3.1	.08	17	2.6	6.0	3.8	69	7.6	2.4	.0	.5	84	53	0	135	6.9	USGS	20
21	5-22-51	56	14	.21	.00	26	2.9	2.8	3.9	92	7.0	2.0	.1	.2	100	77	1	167	8.3	USGS	21
23	4-18-58	----	----	9.3	.35	18	6.8	----	----	56	----	6.9	.2	----	104	73	27	---	6.2	O	23
24	6-13-61	56	10	5.2	.05	3.3	1.7	2.2	1.0	16	5.5	2.2	.0	.2	40	15	2	47	6.1	USGS	24
28	6-4-51	60	9.6	1.1	.00	28	5.2	3.5	6.8	105	20	1.2	.0	2.4	129	91	5	216	7.3	USGS	28
32	5-3-51	58	11	----	.13	19	3.2	4.5	4.1	47	32	2.4	.1	.4	94	61	22	154	6.6	USGS	32
33	11-9-54	----	----	.83	.00	75	17	----	----	95	----	2.0	----	----	---	258	180	---	7.2	O	33
35	5-28-51	56	16	.35	.00	44	3.0	2.2	4.3	153	5.0	3.4	.1	.3	156	122	0	253	7.8	USGS	35
36	5-4-51	59	11	.30	.00	21	5.4	3.3	8.4	107	4.5	2.2	.0	.9	109	75	0	187	7.7	USGS	36
38	1960	----	----	.08	.00	----	----	----	----	113	----	2.0	----	----	112	82	0	---	8.0	D	38
40	1960	----	----	----	.04	----	----	----	----	15	----	9.0	----	----	---	52	40	---	5.6	O	40
41	1-10-61	----	----	.04	----	----	----	----	----	100	----	3.0	.0	----	---	80	0	---	7.8	NJDH	41
45	5-3-51	55	12	.71	----	3.6	1.5	2.8	.5	3	14	3.2	.0	.8	41	15	13	49	5.6	USGS	45
47	5-24-51	----	----	2.0	----	23	10	14	10	88	38	20	----	5.9	---	98	26	---	7.2	O	47
50	11-5-54	----	----	1.1	.00	41	13	----	----	98	----	2.0	----	----	---	156	76	---	7.2	O	50
61	8-14-51	56	26	.10	.00	.8	.9	2.9	2.2	1	10	3.1	.0	.1	49	6	5	49	4.7	USGS	61
66	6-13-61	75	18	.60	.03	39	2.9	2.0	6.0	138	5.9	2.8	.1	.4	145	110	0	232	7.7	USGS	66
67	6-13-61	62	9.3	.18	.14	17	2.8	2.5	2.5	60	11	2.6	.1	.1	77	54	5	128	6.5	USGS	67
73	11-20-57	----	----	.10	----	33	3.9	7.0	----	113	23	3.0	----	----	---	110	18	---	7.9	D	73
77	11-6-54	----	----	.06	.00	12	15	----	----	1	----	9.0	----	----	---	92	91	---	5.9	O	77
89	12-9-58	----	----	4.5	.00	18.4	3.4	7.0	----	57	19	4.0	----	----	---	60	14	---	7.4	D	89
97	6-13-61	65	24	3.7	.06	59	1.3	1.8	2.5	168	18	4.0	.4	.2	205	153	15	305	7.3	USGS	97
118	10-10-57	63	13	.65	.02	38	5.6	2.0	6.7	151	3.5	2.6	.4	.2	141	118	0	239	8.2	USGS	118
124	6-13-61	62	8.8	.72	.02	29	6.4	5.5	13	111	25	1.6	.2	1.0	160	99	8	234	7.5	USGS	124
125	10-10-57	64	17	.87	.06	34	3.2	1.6	4.1	117	7.7	2.4	.4	.0	139	98	2	205	7.3	USGS	125
126	1-4-60	----	----	16	----	22	24	----	----	46	2.8	2.0	----	----	---	46	8	---	6.3	O	126

Table 4.--Chemical analyses, in parts per million, of water from wells in Burlington County, N. J.--Continued

Map no.	Date of collection	Temperature (°F)	Silica (SiO <sub>2</sub> )	Total iron (Fe)	Manganese (Mn)	Calcium (Ca)	Magnesium (Mg)	Sodium (Na)	Potassium (K)	Bicarbonate (HCO <sub>3</sub> )	Sulfate (SO <sub>4</sub> )	Chloride (Cl)	Fluoride (F)	Nitrate (NO <sub>3</sub> )	Dissolved solids (residue on evaporation at 180°C)	Hardness as CaCO <sub>3</sub>		Specific conductance (micro-mhos at 25°C)	pH	Source of data	Map no.
																Calcium magnesium	Noncarbonate				
131	6-13-61	70	16	.50	0.02	36	2.6	1.8	3.2	120	5.0	3.2	0.1	0.2	140	101	2	203	7.8	USGS	131
136	7-20-59	---	---	---	.00	5.0	3.0	---	---	2	15	5.0	---	---	---	24	23	---	5.8	O	136
149	6-13-61	60	32	11	.21	27	3.2	2.5	4.8	34	18	31	.2	.4	170	81	53	210	6.0	USGS	149
158	1- 7-61	---	---	.16	.00	---	---	---	---	142	---	6.0	.2	.2	136	70	0	---	8.1	NJDH	158
168	11- 8-54	---	---	.02	.00	75	18	---	---	109	---	4.0	---	---	---	260	171	---	7.2	O	168
171	4-27-60	---	9.0	.70	.00	20	4.9	12	---	87	7.0	6.0	---	---	---	70	0	---	8.1	O	171
172	4-23-56	58	9.7	.24	.01	27	4.6	2.7	7.4	112	6.4	2.2	0.1	0.9	118	86	0	196	8.0	USGS	172
173	10- 9-57	59	31	9.8	.11	19	1.1	1.6	4.3	47	18	3.5	.8	.4	116	52	14	129	6.5	USGS	173
174	5-24-51	58	9.8	.68	.45	26	4.6	3.2	6.3	98	15	1.2	.1	.6	113	86	3	190	7.9	USGS	174
175	10- -55	---	---	.15	.01	---	---	---	---	11	---	11	---	---	84	50	41	---	6.2	O	175
176	6-13-61	61	45	1.9	.05	33	3.5	2.5	2.5	108	8.4	3.0	.5	.4	154	97	9	197	6.8	USGS	176
177	6-13-61	64	32	7.2	.09	36	3.0	2.0	7.0	62	28	25	.6	.0	180	103	52	247	6.9	USGS	177
181	5-28-51	55	21	.64	.12	37	1.8	2.0	3.3	121	8.4	2.5	.1	.2	139	100	1	214	8.0	USGS	181
184	6-17-59	---	9.0	.64	.03	23	4.6	8.0	---	104	7.1	.8	.0	.4	150	77	0	176	7.1	USGS	184
188	5- 2-51	59	6.0	.00	.00	5.2	.0	5.0	.3	10	.0	3.2	1.9	1.9	44	13	5	46	6.9	USGS	188
189	8-14-51	56	4.3	.12	.00	1.3	1.9	2.4	.7	3	4.5	8.2	.0	.3	26	11	9	79	5.0	USGS	189
190	6-21-51	65	6.1	10	.52	2.4	.4	3.0	.9	5	4.2	5.0	.1	.2	25	8	4	35	5.5	USGS	190
191	8-14-51	56	32	.32	.00	1.0	.8	2.5	2.0	2	7.0	3.6	.0	.2	54	6	4	47	4.7	USGS	191
192	8-14-51	59	5.2	.04	.00	.2	.7	2.6	.8	3	.0	4.2	.0	2.1	16	3	1	23	5.5	USGS	192
193	6-21-51	58	2.7	5.7	.38	1.2	1.6	3.1	.6	0	7.5	8.5	.0	.3	29	10	10	63	4.4	USGS	193
194	6-21-51	57	13	.28	.00	22	5.7	5.3	8.5	111	5.0	2.0	.1	.0	115	78	0	192	8.1	USGS	194
195	10- 9-57	55	20	11	.20	5.9	1.0	1.6	1.8	14	10	2.3	.7	.4	50	19	7	57	6.1	USGS	195
196	6-21-51	58	15	.19	.00	26	2.4	2.7	3.7	93	6.5	1.8	.1	.2	106	75	0	165	8.0	USGS	196
198	9-10-59	---	12	.10	.00	---	---	4.0	---	7	.0	5.0	.1	---	---	8	2	---	6.6	D	198
203	6-13-61	69	1.5	.47	.04	1.6	1.0	1.8	.5	2	6.7	2.2	.1	.2	23	8	7	33	4.8	USGS	203
204	6-13-61	56	13	11	.40	13	5.0	4.4	3.0	18	28	7.6	.2	.2	125	53	24	197	5.8	USGS	204
205	6-13-61	57	14	.25	.10	32	2.1	2.0	3.0	106	4.6	2.8	.4	.2	120	89	2	184	7.7	USGS	205
206	6-13-61	59	10	1.2	.03	33	8.9	2.5	7.5	118	28	1.2	.1	.2	160	119	23	257	7.6	USGS	206
207	6-13-61	70	7.8	11	.29	6.5	3.3	1.8	3.2	4	21	7.6	.3	.2	73	30	26	92	5.1	USGS	207