

DEPARTMENT OF THE INTERIOR
UNITED STATES GEOLOGICAL SURVEY

TO ACCOMPANY
WRI REPORT 83-4118-B

MAPS SHOWING GROUND-WATER LEVELS, SPRINGS, AND
DEPTH TO GROUND WATER,
BASIN AND RANGE PROVINCE, NEW MEXICO

by

B.T. Brady, Deborah A. Mulvhill, Donald Hart, Jr.,
and William H. Langer

INTRODUCTION

This report on ground-water levels, springs and depth to ground water in the Basin and Range province of New Mexico (see index map) was prepared as part of a program of the U.S. Geological Survey to identify prospective regions for further study relative to isolation of high-level nuclear waste (Bedinger, Sargent, and Reed, 1984), utilizing program guidelines defined in Sargent and Bedinger (1984). Also included in this report are selected references on pertinent geologic and hydrologic studies of the region. Other map reports in this series contain detailed data on ground-water quality, surface distribution of selected rock types, tectonic conditions, areal geophysics, Pleistocene lakes and marshes, and mineral and energy resources.

In the Basin and Range province, ground water occurs in basin-fill deposits and consolidated rocks. The basin fill consists mostly of unconsolidated to semi-indurated sedimentary deposits. The material ranges from poorly sorted to moderately sorted mixtures of gravel, sand, silt, and clay that were derived from the consolidated rocks in the nearby mountains. Evaporite deposits, limestone, conglomerate, and volcanic rocks are present in places in the unit. Some of the basins may contain as much as 9,000 feet of basin fill, but the most permeable rocks and most of the recoverable ground water is in the upper 1,000 feet of the unit.

The consolidated rocks consist mostly of sedimentary and volcanic rocks, with lesser amounts of metamorphic and intrusive rocks. The consolidated rocks make up the mountain ranges that border the basins and are the principal source of sedimentary material to the basin fill.

Few wells exist in the consolidated rocks compared to the greater number of wells in the basin fill. The yield of wells tapping many consolidated rock units is due to interception of water in fracture zones. In some areas in the Basin and Range, carbonate rock is extensive in the subsurface and provides interconnection between alluvial basins through fractures and solution channels. Although the consolidated rock commonly has very low permeability, and very low rates of ground-water flow, the entire ground-water system, including basin fill and bedrock, must be treated as one integral system.

GROUND-WATER LEVELS AND SPRINGS

The accompanying map shows the altitude of the ground-water level and the discharge and temperature of selected springs. This information was obtained primarily from reports listed in the "Selected references" and from files of the U.S. Geological Survey. The altitude of the ground-water level is shown by contours in those areas where sufficient data exist. In other areas, where regional data are inadequate to contour, data on selected wells are shown.

Ground-water-level changes occur daily, seasonally, and yearly in response to natural and man-induced hydrologic conditions. The ground-water-level map is a compilation of water-level measurements taken over a number of years. However, for any one ground-water study area the measurements generally were made during a relatively short time period. Collectively, the ground-water levels show the regional configuration of the ground-water surface.

Similarly, water temperatures and discharges of springs may vary with time. The reported values for individual springs were taken at one time; however, collectively the springs were measured over a number of years. Springs may issue from perched water tables or may be under artesian pressure; under these conditions the altitude of springs do not correspond to the altitude of the regional water table.

DEPTH TO GROUND WATER

The accompanying map shows the approximate depth to ground water below the land surface. In the basin fill, ground water moves in a continuous system, although perched ground water may occur in places. In areas with good topographic control, the depth to ground water can be predicted with reasonable accuracy between widely-spaced water-level data points, except where geologic discontinuities, such as faults, may alter the flow pattern. Where data are sufficiently closely spaced, lines of equal depth to water are shown; where data are inadequate to draw lines, data from selected wells are shown. In consolidated rocks, indicated by shaded patterns on the map, ground water occurs in poorly interconnected joints and fractures; water levels in wells may vary in depth within short distances, and numerous zones of perched water may exist.

Line patterns approximately delineate zones within which the depth to ground water is inferred to be greater than 500 feet. These are highly generalized areas based on very limited data on depth to ground water in wells tapping consolidated rock and basin fill and mine shafts; and on information gained from the location and discharge rates of springs, locations of gaining and losing reaches of streams; and estimates of geohydrologic conditions in areas. The areas delineated as possibly having depths to ground water of greater than 500 feet are recognized as having potential for evaluating prospective environments for isolation of high-level nuclear waste in the unsaturated zone.

SELECTED REFERENCES

- Ballance, W. C., 1976, Ground-water resources of the Holloman Air Force Base well-field area, 1967, New Mexico, with a section on Geophysical exploration, by Robert Mattick: U.S. Geological Survey Open-File Report 76-807, 128 p.
- Bedinger, M. S., Sargent, K. A., and Reed, J. E., 1984, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part I, Introduction and guidelines: U.S. Geological Survey Circular 904-A, [in press].
- Bjorklund, L. J., 1957, Reconnaissance of ground-water conditions in the Crow Flats area, Otero County, New Mexico: New Mexico State Engineer Technical Report 8, 26 p.
- Bjorklund, L. J., and Maxwell, B. W., 1961, Availability of ground water in the Albuquerque area, Bernalillo and Sandoval Counties, New Mexico: New Mexico State Engineer Technical Report no. 21, 117 p.
- Bushman, F. X., 1955, Ground-water data for Dwyer Quadrangle, Grant and Luna Counties, New Mexico: New Mexico Bureau of Mines and Mineral Resources Circular 37, 19 p.
- Bushman, F. X., and Valentine, C. P., 1954, Water well records and well water quality in southwestern San Agustin Plains, Catron County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Circular 26, 21 p.
- Clark, N. J., and Summers, W. K., 1971, Records of wells and springs in the Socorro and Magdalena areas, Socorro County, New Mexico, 1968: New Mexico Bureau of Mines and Mineral Resources Circular 115, 51 p.
- Conover, C. S., 1954, Ground-water conditions in the Rincon and Mesilla Valleys and adjacent areas in New Mexico: U.S. Geological Survey Water-Supply Paper 1230, 200 p.
- Cooper, J. B., 1965, Ground-water resources of the northern Tularosa basin near Carrizozo, Lincoln County, New Mexico: U.S. Geological Survey Hydrologic Investigations Atlas, HA-193, scale 1:125,000.
- Davie, William, Jr., and Spiegel, Zane, 1967, Geology and water resources of Las Animas Creek and vicinity, Sierra County, New Mexico: New Mexico State Engineer Hydrographic Survey Report, 44 p.
- Doty, G. C., 1960, Reconnaissance of ground water in Playas Valley, Hidalgo County, New Mexico: New Mexico State Engineer Technical Report, no. 15, 40 p.
- _____, 1963, Water-supply development at the National Aeronautics and Space Agency--Apollo propulsion system development facility, Dona Ana County, New Mexico: U.S. Geological Survey Open-File Report, 40 p.
- _____, 1969, Availability of groundwater near Arena, Luna County, New Mexico: U.S. Geological Survey Open-File Report, 21 p.
- Garza, Sergio, and McLean, J. S., 1977, Fresh - water resources in the southeastern part of the Tularosa basin: New Mexico State Engineer Technical Report 40, 67 p.

- Herrick, E. H., and Davis, L. V., 1965, Availability of ground water in Tularosa basin and adjoining areas, New Mexico and Texas: U.S. Geological Survey Hydrologic Investigations Atlas HA-191, scale 1:500,000.
- Hood, J. W., and Herrick, E. H., 1965, Water resources of the Three Rivers area, Otero and Lincoln Counties, New Mexico: U.S. Geological Survey Hydrologic Investigations Atlas, HA-192, scale 1:126,720.
- King, W. E., Hawley, J. W., Taylor, A. M., and Wilson, R. P., 1969, Hydrogeology of the Rio Grande Valley and adjacent intermontaine areas of southern New Mexico: Las Cruces, New Mexico, New Mexico State University, Water Resources Research Institute Report no. 6, 141 p.
- _____, 1971, Geology and ground-water resources of central and western Dona Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Hydrologic Report no. 1, 64 p.
- Leggat, E. R., Lowry, M. E., and Hood, J. W., 1963, Ground-water resources of the lower Mesilla Valley, Texas and New Mexico: U.S. Geological Survey Water-Supply Paper 1669-AA, 49 p.
- McLean, J. S., 1977, Hydrologic maps and data in the Mimbres Basin, New Mexico: U.S. Geological Survey Open-File Report 77-314, 531 p.
- Meinzer, O. E., 1911, Geology and water resources of Estancia Valley, New Mexico, with notes on ground-water conditions in adjacent parts of central New Mexico: U.S. Geological Survey Water Supply Paper 275, 89 p.
- Meinzer, O. E., and Hare, R. F., 1915, Geology and water resources of Tularosa basin, New Mexico: U.S. Geological Survey Water-Supply Paper 343, 317 p.
- Mourant, W. A., 1980, Hydrologic maps and data for Santa Fe County, New Mexico: New Mexico State Engineer Basic Data Report, 180 p.
- O'Brien, K. M., and Stone, W. J., 1981, Water-level data compiled for hydrogeologic study of Animas Valley, Hidalgo County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Open-File Report 130, 64 p.
- Reeder, H. O., 1957, Groundwater in the Animas Valley, Hidalgo County, New Mexico: New Mexico State Engineer Technical Report no. 11, 101 p.
- Sargent, K. A., and Bedinger, M. S., 1984, Geologic and hydrologic characterization and evaluation of the Basin and Range province relative to the disposal of high-level radioactive waste--Part II, Geologic and hydrologic characterization: U.S. Geological Survey Circular 904-B, [in press].
- Sloan, C. E., and Garber, M. S., 1971, Groundwater hydrology of the Mescalero Apache Indian Reservation, south-central New Mexico: U.S. Geological Survey Hydrologic Investigations Atlas HA-349, scale 1:125,000.
- Smith, R. E., 1957, Geology and ground-water resources of Torrance County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Ground-water Report no. 5, 186 p.
- Sorensen, E. F., 1982, Water use by categories in New Mexico counties and river basins, and irrigated acreage in 1980: New Mexico State Engineer Technical Report 44, 51 p.

- Spiegel, Zane, 1955, Geology and groundwater resources of northeastern Socorro County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Ground-water Report no. 4, 99 p.
- Titus, F. B., 1980, Ground-water in the Sandia and northern Manzano Mountains, New Mexico: New Mexico Bureau of Mines and Mineral Resources Hydrologic Report no. 5, 66 p.
- Trauger, F. D., 1972, Water resources and general geology of Grant County, New Mexico: New Mexico State Bureau of Mines and Mineral Resources Hydrologic Report 2, 211 p.
- Trauger, F. D., and Herrick, E. H., 1962, Ground water in central Hachita Valley, northeast of the Big Hatchet Mountains, Hidalgo County, New Mexico: New Mexico State Engineer Technical Report no. 26, 21 p.
- Weir, J. E., Jr., 1965, Geology and availability of ground water in the northern part of the White Sands Missile Range and vicinity, New Mexico: U.S. Geological Survey Water-Supply Paper 1801, 78 p.
- West, S. W., and Broadhurst, W. L., 1975, Summary appraisals of the nation's ground-water resources--Rio Grande Region: U.S. Geological Survey Professional Paper 813-D, 39 p.
- Wilson, C. A., White, R. R., Orr, B. R., and Roybal, R. G., 1981, Water resources of the Rincon and Mesilla Valleys and adjacent areas, New Mexico: New Mexico State Engineer Technical Report 43, 514 p.