

Maps showing distribution of dissolved solids  
and dominant chemical type in ground water,  
Basin and Range province, Oregon

by

Thomas H. Thompson, Richard Chappell,  
J. B. Gonthier, and W. D. McFarland

Introduction

This map report is one of a series of geologic and hydrologic maps of States in the Basin and Range Province. These map reports contain information on ground-water hydrology, ground-water quality, surface distribution of selected rock types, data on tectonic conditions, areal geophysical data, Pleistocene lakes and marshes and natural resources. This information is the basis for a summary report that characterizes the Basin and Range Province. The geologic and hydrologic characterization will be used to evaluate the Province for prospective regions for further study relative to isolation of high-level radioactive waste. (Bedinger, Sargent and Reed, 1984).

## Chemical quality of ground water

Ground-water quality in the Basin and Range Province of Oregon was characterized by dissolved-solids concentration and dominant anions and cations in solution. Water-quality information was compiled from the U.S. Geological Survey water-quality files (WATSTORE), published reports of the U.S. Geological Survey, and the Oregon Water Resources Department.

The data on dissolved-solids concentration and water-quality type were plotted and their distributions mapped in the basin-fill deposits from the data compiled for samples collected from non-geothermal springs and wells of less than 501 feet in depth. In areas for which data were not compiled, other data sources and published reports were consulted where available. In lieu of data, the parameters in the basin-fill were estimated from the position in the ground-water flow system and the lithology of the adjacent bedrock. Data from wells greater than 500 feet in depth are included as supplemental data, but are not used in mapping dissolved-solids concentration or water-quality type. Data from wells greater than 500 feet in depth are not reflected in the mapped areas where the data are not believed to be representative of a significantly large area.

### Distribution of dissolved solids

Dissolved-solids concentration of major anions and cations was calculated from analyses using the following equation:

Dissolved solids =

$$\text{Ca} + \text{Mg} + \text{Na} + (0.4917)(\text{HCO}_3) + \text{SO}_4 + \text{Cl} + \text{K} + \text{Sr},$$

where the concentration of each component, calcium, Ca;

magnesium, Mg; sodium, Na; bicarbonate,  $\text{HCO}_3$ ; sulfate,  $\text{SO}_4$ ;

chloride, Cl; potassium, K; and strontium, Sr; is in milligrams per liter (mg/L).

### Water-quality type

The dissolved-solids composition of a water consists primarily of six major ions: the three cations - calcium, magnesium, and sodium; and the three anions - bicarbonate, sulfate, and chloride. Relative proportions of these cations and anions can be displayed on trilinear diagrams: Waters of similar ionic proportions, and thus of similar chemical character, plot near one another. Therefore, areas of chemically similar waters can be delineated on the trilinear diagrams. For the purposes of this study, each trilinear diagram is divided into seven areas as shown in the accompanying illustration.

Water quality in basin fill is characterized by dominant cations and anions in solution and is shown in the accompanying map. The explanation lists dominant cation(s) and anion(s) combinations mapped in each area.

### Selected References

- Bedinger, M. S., Sargent, K. A., and Reed, J. E., 1984, Geologic and hydrologic characterization and evaluation of the Basin and Range Province relative to the disposal of high-level radioactive waste--Part I, Introduction and guidelines: U.S. Geological Survey Circular 904-A, [in press].
- Brown, D. E., and Peterson, N. V., 1980, Preliminary geology and geothermal resource potential of the Alvord Desert area, Oregon: Oregon Department of Geology and Mineral Industries, Open-File Report 0-80-10, scale 1:250,000, 57 p.
- Gonthier, J. B., Collins, C. A., and Anderson, D. B., 1977, Groundwater data for the Drewsey Resource area, Harney and Malheur Counties, Oregon: U.S. Geological Survey Open-File Report 77-741, 28 p.
- Hampton, E. R., 1964, Geologic factors that control the occurrence and availability of ground water in the Fort Rock Basin, Lake County, Oregon: U.S. Geological Survey Professional Paper 383-B, 29 p.
- Huxel, C. J., Jr., Parker, J. E., and Everett, D. E., 1966, Effects of irrigation development on the water supply of Quinn River Valley area, Nevada and Oregon, 1950-64: Nevada Department of Conservation and Natural Resources, Water Resources Bulletin no. 34, 80 p.
- Leonard, A. R., 1970, Ground-water resources in Harney Valley, Harney County, Oregon: Oregon Water Resources Department Ground Water Report no. 16, 85 p.

- Leonard, A. R., and Harris, A. B., 1974, Ground-water in selected areas in the Klamath Basin, Oregon: Oregon Water Resources Department Ground Water Report, no. 21, 104 p.
- Phillips, K. N., Newcomb, R. C., Swenson, H. A., and Laird, L. B., 1965, Water for Oregon: U.S. Geological Survey Water-Supply Paper 1649, 150 p.
- Phillips, K. N., and Van Denburgh, A. S., 1971, Hydrology and geochemistry of Abert, Summer and Goose Lakes and other closed-basin lakes in south-central Oregon: U.S. Geological Survey Professional Paper 502-B, 86 p.
- Piper, A. M., Robinson, T. W., and Park, C. F., 1939, Geology and ground-water resources of the Harney Basin, Oregon: U.S. Geological Survey Water-Supply Paper 841, 189 p.
- Sammel, E. A., 1980, Hydrogeological appraisal of the Klamath Falls geothermal area, Oregon: U.S. Geological Survey Professional Paper 1044-G, 45 p.
- Sammel, E. A., and Craig, R. W., 1981, The geothermal hydrology of Warner Valley, Oregon--A reconnaissance study: U.S. Geological Professional Paper 1044-I, p. 11-151.
- Trauger, F. D., 1950, Basic ground-water data in Lake County, Oregon: U.S. Geological Survey Open-File Report, 287 p.