

**Table 10.** Summary of organic compound concentrations in water samples collected during baseline, 1995–98, and artificial recharge, 1996–98, conditions—Continued

Data-collection site (figs. 1, 2, 3, or 5)	Water-quality conditions (dates of collection)	Diuron ( $\mu\text{g/L}$ )	EPTC ( $\mu\text{g/L}$ )	Ethoprop ( $\mu\text{g/L}$ )	Ethal-fluralin ( $\mu\text{g/L}$ )	Linuron ( $\mu\text{g/L}$ )	Metol-achlor ( $\mu\text{g/L}$ )	Metribuzin ( $\mu\text{g/L}$ )	Naprop- amide ( $\mu\text{g/L}$ )
<b>Ground-water or directed-water monitoring sites—Continued</b>									
<b>Bals lead recharge site—deep monitoring wells</b>									
	Baseline (May 13, 1997)	<0.02 (2) —	<0.002 (2) —	<0.003 (2) —	<0.004 (2) —	<0.002 (2) —	<0.002 (2) —	<0.004 (2) —	<0.003 (2) —
	Recharge (May 29, 1997 through July 1998)	<0.02 (7) —	<0.002 (17) —	<0.003 (17) —	<0.004 (17) —	<0.002 (17) —	<0.002–<0.05 (36) 0.016*	<0.004–<0.05 (36) —	<0.004–<0.003 (17) —
<b>Sedgwick recharge site—treated directed source water</b>									
	Recharge (April 1998 through July 1998—only sampled during recharge activities)	<0.02 (2) —	<0.002 (3) —	<0.003 (3) —	<0.004 (3) —	<0.002 (3) —	<0.05–3.46 (24) 0.14*	<0.004–0.06 (24) —	<0.003 (3) —
<b>Sedgwick recharge site—shallow monitoring wells</b>									
	Baseline (June 1997 through February 1998)	<0.02 (2) —	<0.002–0.014 (4) —	<0.003 (4) —	<0.004 (4) —	<0.002 (4) —	$\leq$ 0.0029–7.0 (12) 2.5*	<0.004–<0.05 (12) —	<0.003 (4) —
	Recharge (April 1998 through July 1998)	<0.02 (2) —	<0.002–0.0065 (4) —	<0.003 (4) —	<0.004 (4) —	<0.002 (4) —	0.0079–3.0 (9) 1.2*	<0.004–<0.05 (9) —	<0.003 (4) —
<b>Sedgwick recharge site—deep monitoring wells</b>									
	Baseline (June 1997 through April 1998)	<0.02 (3) —	<0.002 (3) —	<0.003 (3) —	<0.004 (3) —	<0.002 (3) —	$\leq$ 0.0016–<0.05 (9) —	<0.004–<0.05 (9) —	<0.003 (3) —
	Recharge (April 1998 through July 1998)	<0.02 (2) —	<0.002 (4) —	<0.003 (4) —	<0.004 (4) —	<0.002 (4) —	<0.002–<0.05 (6) —	<0.004–<0.05 (6) —	<0.003 (4) —